

Harvesting, Storing, and Feeding Forages as Round Bale Silage¹

Matt Hersom and William E. Kunkle²

Introduction

Perennial warm-season grasses grown in Florida are often dried and harvested as hay. Florida weather data shows less than a 20% probability of 3 consecutive drying days from mid-June through August. These frequent rains often delay harvest and/or cause considerable field losses of forages. Maintaining a regular harvest schedule (every 4 to 5 weeks) during the summer months is essential for good forage quality.

Forages are often harvested at high moisture and ensiled when field drying is not feasible or possible. High costs for equipment and structures needed to chop, haul, and store silage have limited its use for perennial warm-season forages, especially for the medium and small beef and dairy producers. Forages have additional challenges because yields at each harvest are lower than corn or sorghums, they are cut several times during the summer, and they need to be wilted prior to ensiling, which is not compatible with storing in bunker silos or in piles.

Development of systems for ensiling chopped forage in long polyethylene tubes was very applicable to timely harvest of warm-season perennial forages in Florida. However, this system required costly specialized equipment for chopping and bagging the forage that limited its application to small

and medium beef and dairy producers. Round bale silage was developed in Europe and has been used since the mid-1980s in the U.S. Round bale silage is harvested with the same equipment used for making hay, requiring only the addition of a plastic wrapper for applying the “stretch wrap” to the bales. Round bale silage offers flexibility in harvesting small quantities at one time, requires less additional equipment to purchase than chopped silage systems, offers flexibility of making hay when weather permits and silage when the weather necessitates, and does not require much additional equipment operation and repair skills. This paper will review the research, experiences, and challenges of the round bale harvesting, storing, and feeding system.

Forage Quality Declines with Maturity

Bermuda and star grasses produce large yields during the warm season but quality declines rapidly after 4 weeks of regrowth (Table 1). These grasses are not high in quality at four weeks regrowth, and allowing them to grow past six weeks does not increase yearly dry matter production and forage quality is near maintenance (QI - 1.0) or below. Growing or lactating cattle fed these forages will require more supplement to attain desired performance. Digit and limpo grasses are higher in quality than bermuda and star

1. This document is AN145, one of a series of the Department of Animal Sciences, Florida Cooperative Extension Service, UF/IFAS. Originally published October 2003. Revised September 2011. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Matt Hersom, associate professor, Department of Animal Sciences; and William E. Kunkle, professor (deceased), Department of Animal Sciences; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean

grasses at similar maturity, but they also decline in quality during regrowth as the proportion of stem increases.

Research from Louisiana further emphasizes the importance of regrowth interval on forage quality (Table 2). The decline in quality reflected in gains of steers is more rapid in bermudagrass grown during the summer than in bermudagrass grown during the spring. Summer regrowth may have lower quality because higher temperatures increase lignin deposition and high rainfall increases growth rates and maturation. This further emphasizes the need to cut warm-season perennial grasses every 4 weeks to retain quality.

Round bale silage allows for harvest of forage every 4 to 5 weeks when the quality is better. This decreases or eliminates the need for supplements when feeding the forage to livestock. We suggest developing a plan to cut warm-season grasses by the calendar; make hay when the weather allows, and make round bale silage when the weather will not allow you to make hay.

Wrapping, Storing, and Feeding

Allowing air into ensiled forage will reduce the quality of round bale silage. Preservation of wet forages requires that air (oxygen) be kept out. Molds and other organisms that rot wet forages need air to grow. If air enters, the forage molds initially, and then rots over time. This has been the biggest problem for producers using the round bale system.

Round bale silage was first stored in individual bale bags that were put on bales lifted off the ground then gathered and tied with string to seal (Cromwell et al. 1994a). The bag cost (\$6 to \$8/bag) and labor to put bales in bags limited adoption of this approach. Another approach tried was to store bales in long plastic tubes (a long bag), and mechanized bale tube stuffers were developed. The long tube was more cost effective (\$4 to \$5/bale) and is still used by some producers. Our experience was that small hole(s) in one end allowed air to move freely in the tube, resulting in molding of bales in too many instances. A more recent version of this system has tubes that are stretched in diameter while loading bales, which reduces air space and movement.

The “stretch-wrap” round bale silage system has been used for over 20 years and has become the most widely used round bale silage storage system in Florida and across the U.S. The following discussion will address experiences with the stretch-wrap storage system.

Stretch-wrap plastic quality. The intense sunlight for many months in Florida can cause the plastic to lose its pliability and break up in pieces. Conditions in Florida require more additives in stretch-wrap plastic to resist sunlight degradation compared to other areas of the U.S. In the early 1990s, four tests (each a year long) were conducted to evaluate stability of plastic in sunlight (Cromwell et al. 1997b). Some samples of plastic were stable in sunlight for less than a month, while others were stable for over a year. Sunlight stability was a noted problem when we started the tests but problems decreased over the years. Most problems occurred with companies or employees new to the industry, poor mixing of ingredients prior to making stretch-wrap plastic, or problems with the manufacturing equipment. The frequency of failures has declined over the years, but several producers no longer make round bale silage because of their bad experience with plastic. If you notice sunlight degradation of stretch-wrap plastic, it is suggested to feed this forage before it rots or rewrap the bales. Many problems can be avoided by testing new sources of stretch wrap on a few bales for one or more years before risking a considerable portion of your harvested forage.

Other problems with stretch-wrap plastics have been air entry and spoilage without visible sunlight breakdown or visible holes. This problem has not been frequent and was usually traced to too few layers of stretch wrap, inappropriate stretching of the film (50%–60% usually suggested), or reprocessed resins used in manufacturing of stretch wrap. Six layers of stretch wrap are recommended to reduce holes and keep the air out. Although four layers have been used successfully, spoilage has been found in a few instances when no holes were present. These problems may have been caused by using reprocessed resin to make stretch-wrap plastic. Stretch-wrap plastic manufacturers that used this cost-cutting approach usually have not continued to market stretch-wrap plastic.

Wrapping machine. The most common system in Florida is in-line wrapping machines that wrap bales with stretch wrap in a long line. Round bales are loaded on this machine end to end, and wrapped with stretch wrap continuously. The wrapped bales move over rollers that gradually lower them to the ground as the machine moves ahead. In-line wrappers are less likely to damage stretch wrap after wrapping (they don’t move wrapped bales or drop them to the ground), and they use about half as much stretch wrap for each bale as machines that wrap bales individually. Although in-line wrappers cost a bit more initially, the savings in plastic and labor may offset this in many situations.

Although most of the older bale wrappers applied 20-inch-wide stretch wrap, most of the newer bale wrappers apply 20- and 30-inch-wide stretch wrap, and 30-inch is used widely by producers for cost and time reasons. The pretensioner must evenly stretch the plastic at the specified amount, usually 50%–60%. Occasionally pretensioners will under- or overstretch plastic wrapping machines, tilting their wrapping table for the bale to roll off the machine after wrapping. An uneven soil surface, rocks, stubble, etc., can puncture the stretch wrap or cause excessive stretching in a small area, which can lead to a hole after a few months of weathering. A table attachment on the wrapper that lowers the bale to the ground and decreases the distance a bale will roll is an option that can reduce plastic damage and reduce problems under some conditions.

Time of wrapping. How long can I wait from baling to wrapping bales? This has been a common question. Producers have observed that wet bales will heat if allowed to sit a few hours before wrapping. Air exposure and high moisture allow plant respiration and fungus (mold) growth, producing heat that is not dissipated once the wet forage is in a bale. Both plant respiration and fungus growth use water-soluble sugars that are the primary substrates for producing organic acids that lower the pH and ensile the forage. With this reasoning, it seems obvious that wrapping soon after baling is recommended. However, forage producers know that labor is limited, equipment breaks down, and rains and lightning are common during the summer. A study evaluating the effects of delaying wrapping up to 48 hours was conducted over 2 years to help answer this question (Garces-Yopez et al. 1992). Bale temperatures reached over 150°F when wrapping was delayed 48–52 hours (Table 3). Bale temperatures declined and within 1 week after wrapping temperatures were below 140°F, which is considered critical to avoid lowering protein digestibility caused by the high temperatures. Low acid detergent fiber nitrogen (ADFN) concentrations and similar concentrations across all time delays indicate higher temperatures did not increase ADFN, which is indirectly related to reduced protein digestibility. Acid detergent fiber nitrogen concentrations averaged 8% and 6% of total nitrogen in the two trials. Differences in dry matter recovery percentages, in vitro digestibilities, and pH values for delaying wrapping up to 48 hours were not found in this trial. A visual evaluation of molding on the bales did not show differences either. This trial did not evaluate animal preferences, intakes, or performance, but large differences in forage preservation due to delaying wrapping up to 48 hours were not found. However, it is recommended to wrap bales the same day.

Storage issues: holes in the plastic resulting in molding and rotting of the forage have been the biggest problem with round bale silage. Animals and birds can and sometimes do make holes in the plastic. Problems with cattle, deer, hogs, raccoons, groundhogs, cats, rats, mice, and various birds have been reported. Choosing a storage area where problems with these animals are less likely is the most workable solution. Tree limbs, hail, and soldier fly larva can also cause holes. Ants may not make the hole but have been observed to enter the bales and increase spoilage. Weekly monitoring of plastic for holes and patching with tape designed for this purpose is highly recommended. Bales with holes should be fed soon after holes are made (even if patched) or rewrapped if feeding time is a few months away. Bales damaged during wrapping should be rewrapped or separated so they can be fed first (patched areas tend to fail before undamaged plastic). Selecting bales with damaged plastic to feed first can help reduce losses.

Stacking bales has been tried but it is not recommended for most situations. Only bales wrapped individually can be stacked. Bales containing forage baled when too wet often shrink, and these change in shape after a few months of storage. Piling wet bales results in additional distortion in shape that causes air leakage and bales falling off the pile. If storage space is limited and piling is necessary, consider stacking on end. Moving the wrapped bales with equipment that does not damage the plastic is essential.

Feeding issues. Round bale silage can be handled and fed much like hay. Plastic is usually removed and left in the storage area. Using a round bale feeder or unwrapping bales helps to reduce trampling and waste. Bales may start to heat after 2 to 4 days in the open air, but this heating is usually not as much of a problem as with chopped silage. Presumably, the dense bale slows the air entry that causes the heating. Offering round bale silage that will be consumed in 4 days or less has worked well. However, longer feeding periods are possible during cooler weather. Molding and rotting can occur from holes in plastic, and some of this forage will not be eaten. Molded forage that is consumed has not caused problems in most situations. White, pink, gray, and blue mold have been observed, but problems with cattle deaths or noticeable reductions in performance have not been reported. Horse deaths from botulism have been reported when round bale silage was fed. This was associated with dead animals (rats, mice, etc.) trapped during baling.

Plastic and wrapper costs. Round bale silage can be cost-effective in some situations. Typical costs of equipment depreciation and plastic are itemized for the individual bale

and in-line wrapping systems at different annual uses (Table 4).

These are extra costs above those for making hay. Feedback from producers with automated wrappers (wrappers with power supplies and automatic start and stop) that already haul their hay to storage areas indicated labor required to fluff hay prior to raking was similar to labor required to wrap silage bales. Costs associated with extra labor to wrap forage need to be added in some situations. Part of added costs for round bale silage will be offset by reduced field and storage losses compared to hay stored outside. Added costs range from \$3.50 to \$8.50/bale or \$11 to \$27/ton of dry forage.

Forage Management Is Important

Harvesting hay or round bale silage preserves the forage for feeding at a later time. Neither system improves the quality of harvested forage; therefore, harvesting high quality forage is essential. Ensiling forage requires an anaerobic environment (no oxygen) that allows fermentation of sugars to organic acids. Organic acids such as lactic acid lower the pH, and this inhibits bacteria growth and fermentation. If fermentable sugars are gone before the pH is low enough, clostridia bacteria can grow, breaking down lactic acid and protein to undesirable end products. Bermudagrass is difficult to ensile because of low sugar concentration (usually less than 4% fermentable sugars; Bates et al. 1989a), high buffering capacity (more lactic acid needed to lower pH), and fermentation of some substrates (malic and oxaloacetic acids) to weak acids (acetic) that buffer near pH 5. Wilting forage to 35% or higher dry matter is a strategy used to preserve forages like bermudagrass that are difficult to ensile. Higher dry matter inhibits the growth of the clostridia bacteria.

Wilting. Bermudagrass cut at 4 to 6 weeks regrowth is typically 18%–28% dry matter at cutting. Wilting bermudagrass for 2 to 4 hours during good summer drying conditions increased dry matter 20 percentage units to 46% dry matter, reduced bale weights over 100 pounds, and increased dry matter in each bale over 200 pounds (Table 5). Wilting forage 2 to 4 hours resulted in 19% higher dry matter intake and 0.4 pounds per day greater gains in growing cattle. Bermudagrass wilted and stored as round bale silage had bale dry matter weight and cattle gains similar to those associated with forage harvested as hay, but storage and feeding losses were 4 to 6 percentage units better for hay. Wilting compared to direct cut round bale silage also reduces number of bales and storage costs by over 35%.

Most perennial forages (bermudagrass, stargrass, limpo-grass, perennial peanut) should benefit from wilting prior to baling. Annual forages such as millet and sudangrass are very high in moisture when cut, and wilting to increase dry matter is essential for good quality silage.

Additives. Microbial inoculants, enzymes, sugars, ammonia, and other additives can be added to enhance fermentation and preservation of silage. Limited research has been conducted with these additives on round bale silage. Microbial inoculants have been shown to increase the rate of fermentation and result in a lower final pH with chopped silage, and similar results have been found with round bale silage (Bates et al. 1989a). Trials evaluating microbial inoculants that provide fast-growing, lactic-acid-producing bacteria often show advantages in silage fermentation, but trials showing advantages in dry matter recovery and cattle performance are limited. Research evaluating effects of microbial inoculation of bermudagrass round bale silage produced mixed results. Growing cattle fed microbial-inoculated (lactic acid bacteria) bermudagrass round bale silage had 0.16 pounds per day greater gains (0.08 versus 0.24 pounds per day) than those fed uninoculated direct cut silage (70%–75% moisture), but microbial inoculation did not improve performance (0.50 versus 0.44 pounds per day) in wilted (50%–60% moisture) bermudagrass round bale silage. This research showed greater intakes and performance for inoculated direct cut bermudagrass silage, but gains of these cattle were still below gains of cattle fed wilted silage, indicating wilting was more important than inoculation. An evaluation of several experiments suggests that microbial inoculants often give improvements when ensiling conditions are marginal.

Ammonia is a mold inhibitor and has been shown to reduce visible molds on the outside of bales (Bates et al. 1989b). Unfortunately, ammoniation (1% of dry matter) has been associated with an undesirable fermentation in direct cut bermudagrass round bale silage characterized by more acetic acid and a higher pH. Enzyme additives to bermudagrass silage have shown promising results by increasing the fermentable substrate (Bates et al. 1989b); however, the specific enzymes and quantities (activities) needed for the optimal effect have not been determined.

Summary

The round bale storing system allows harvesting warm-season grasses every 4 to 5 weeks when the quality is good. This will reduce or eliminate the need for supplements when feeding the forage to beef cattle. Plan to cut warm-season grass by the calendar, make hay when the weather

allows, and make round bale silage when the weather will not allow drying for hay. Quality of round bale silage is dependent on excluding air from the ensiled forage. Holes in the plastic resulting in molding and rotting of the forage have been the biggest problem with round bale silage. Wilting warm-season grasses to increase dry matter 40%–50% before baling reduces the number of bales and storage costs, decreases storage losses, and improves cattle performance.

Literature Cited

Bates, D.B., W.E. Kunkle, C.G. Chambliss, and R.P.

Cromwell. 1989a. “Effect of dry matter and additives on bermudagrass and rhizoma peanut round bale silage.” *J. Prod. Agric.* 2:91-96.

Bates, D.B., W.E. Kunkle, T.E. Dawson, A. Berthe, S.C. Denham, C.G. Chambliss, R. C. Cromwell, J.G. Wasdin, and D.L. Wakeman. 1989b. “Round bale silage - A forage harvesting alternative.” *Proc. 38th Annual Beef Cattle Shortcourse*, pp 45-55, Univ. of Florida.

Berthe, A., W.E. Kunkle, and D.B. Bates. 1991. “Effects of harvesting system and wilting on bermudagrass intake and liveweight change of growing heifers.” *J. Anim. Sci.* 69:283 (Suppl. 1).

Brown, W.F., and R.S. Kalmbacher. 1998. “Nutritional value of native range and improved forages: A perspective from central and south Florida.” *Proc. 47th Annual Beef Cattle Shortcourse*, pp 79-87, Univ. of Florida, Gainesville.

Cromwell, R.P., W.E. Kunkle, and C.G. Chambliss. 1994a. “Equipment for preserving forage as round-bale silage.” Cooperative Extension Service, Circular 1071, 7 pp, Univ. of Florida, Gainesville.

Cromwell, R.P., W.E. Kunkle, G.D. Sadler, and C.G. Chambliss. 1994b. “The plastic wrapper is the key to making high quality round bale silage.” Cooperative Extension Service, Circular 1072, 6 pp, Univ. of Florida, Gainesville.

Yepez, P.G., W.E. Kunkle, and D.B. Bates. 1992. “Effect of delaying storage on the dry matter recovery and quality of bermudagrass ensiled in round bales wrapped with plastic.” *J. Anim. Sci.* 70:191 (Suppl. 1).

Table 1. Effects of grass and maturity (age in weeks) on forage quality.^a

Grass	TDN ^b			Voluntary intake ^c			Quality index ^d		
	4	6	8	4	6	8	4	6	8
Bahia	56	55	54	2.3	2.1	1.7	1.2	1.1	0.9
Bermuda	57	52	44	2.3	2.2	1.8	1.3	1.1	0.8
Star	60	53	49	2.4	2.5	2.1	1.4	1.3	1.0
Digit	60	58	57	2.5	2.7	2.2	1.5	1.5	1.2
Limpo	63	63	56	2.5	2.3	2.2	1.5	1.4	1.2

^aAdapted from Brown and Kalmbacher, pp 79-87, in 47th Florida Beef Cattle Shortcourse Proc., May 1988.

^bTotal Digestible Nutrients, percentage of dry matter.

^cIntake of dry matter expressed as percent of body weight.

^dVoluntary TDN intake relative to maintenance requirement: 1.0 = maintenance.

Table 2. Quality of coastal bermudagrass hay harvested at different maturities and seasons.^a

Item	Weeks of regrowth	Harvest date				
		06/14	07/12	08/09	09/06	10/04
TDN, % DM ^b	4	55	57	52	53	46
	6	52	51	47	49	48
	8	52	51	46	47	44
QI ^c	4	1.4	1.4	1.3	1.3	1.1
	6	1.3	1.4	1.0	1.2	1.2
	8	1.3	1.1	0.9	1.1	0.8
ADG, lb ^d	4	0.57	0.78	0.72	0.63	0.28
	6	0.34	0.48	-0.04	0.42	0.22
	8	0.16	0.07	-0.39	0.07	-0.39

^aAdapted from Nelson, et al., 1980, Louisiana Agr. Exp. Sta. Bul. 730.

^bTotal Digestible Nutrients, percentage of dry matter.

^cVoluntary TDN intake relative to maintenance requirement: 1.0 = maintenance.

^dDaily gains, in pounds; feeding trial conducted with steers from December through February for all hays.

Table 3. Effects of delaying time from baling to wrapping on bermudagrass round bale silage.^a

	Time from baling to wrapping, hr				S.E.
	1-4	19-22	27-31	48-52	
Year 1989 ^b					
Dry matter recovery, %	99.5	96.0	101.6	100.8	3.0
ADF-N ^c , % DM	0.17	0.15	0.13	0.14	0.02
IVOMD, % ^d	50.0	50.3	52.4	50.7	0.92
pH	5.81	6.02	5.99	5.95	0.07
Temperature, (days 9-14)					
- Peak, F°	88	93	95	104	1.4
- Average, F°	83	86	87	91	1.1
Year 1990 ^e					
Dry matter recovery, %	96.5	97.1	98.5	98.9	2.0
ADF-N ^c , % DM	0.08	0.08	0.08	0.10	0.01
IVOMD, % ^d	39.0	42.1	42.4	41.7	0.93
pH	5.40	5.47	5.46	5.77	0.11
Temperature, (days 3-8)					
- Peak, F°	118	133	136	150	1.8
- Average, F°	112	121	123	136	0.7

^aGarces-Yepez, et al., 1992.

^bBales made on October 31, 1989; bale weight averaged 1,035 lb; 37.3% dry matter.

^cAcid detergent fiber nitrogen as a percent of dry matter.^dIn-vitro organic matter digestibility.

^eBales made on September 6, 1990; bale weight averaged 1,275 lb; 38.0% dry matter.

Table 4. Depreciation and plastic costs for two round bale silage wrapping systems.^a

Bales/year ^b	Individually wrapped bales			In-line wrapped bales		
	500	1,000	2,000	500	1,000	2,000
Hay, tons ^c	180	360	720	180	360	720
Wrapper depreciation, \$/bale ^{de}	5.00	2.50	1.25	6.00	3.00	1.50
Stretch-wrap plastic, \$/bale	3.50	3.50	3.50	2.00	2.00	2.00
Total cost, \$/bale	8.50	6.00	4.75	8.00	5.00	3.50
Total cost, \$/ton dry forage	27.00	19.00	15.10	25.40	15.90	11.10

^aAdded costs above hay harvest, additional labor required in some situations.
^bBales 4 feet wide by 5 feet high, 1,400 lb, 45% dry matter, 630 lb dry matter/bale.
^cTons of hay with equivalent dry forage.
^dIndividual bale wrapper \$12,500 cost; depreciated over 5 years, \$2,500/year.
^eIn-line bale wrapper \$15,000 cost; depreciated over 5 years, \$3,000/year.

Table 5. Effects of wilting bermudagrass round bale silage on forage preservation and performance of growing cattle.^{ab}

	Wilting time, hours			Hay
	None	1-2	2-4	
Forage dry matter, %	26.8	36.2	45.9	87.4
Wet bale weight, lb	1,450	1,425	1,340	710
Dry bale weight, lb	390	515	620	620
Storage losses, %	12.5	11.5	8.8	3.0
Feeding losses, %	10.1	12.7	16.8	12.5
Dry matter intake, lb ^c	8.58	9.55	10.20	9.82
Daily gain, lb	-0.15	+0.02	+0.26	+0.28

^aBerthe et al., 1991; Research conducted over two years, bermudagrass cut after 5 to 6 weeks regrowth in Year 1 and 6 to 7 weeks regrowth in Year 2.
^bBales were 4 feet wide and 4.5 feet high made with New Holland 848 baler. High moisture forage wrapped with 4 layers of stretch-wrap plastic.
^cCattle weighed 550 lb at start of trial in Year 1; cattle weighed 520 lb at start of trial in Year 2.