

UF/IFAS Standardized Nutrient Recommendations for Vegetable Crop Production in Florida¹

Rao Mylavarapu, George Hochmuth, and Guodong Liu²

Introduction

This publication presents the fertilization recommendations for vegetable crops based on soil tests performed by the IFAS Extension Soil Testing Laboratory (ESTL). It contains the basic information from which ESTL soil test reports and fertilization recommendations are generated. Additional information on nutrient recommendations is presented in the Vegetable Production Handbook of Florida, 2017–2018. Similarly, UF/IFAS Standardized Nutrient Recommendations for Agronomic Crops can be found in SL129 (<http://edis.ifas.ufl.edu/ss163>) (Mylavarapu 2015).

Soil Testing

Soil testing is a scientific tool for effective nutrient management that provides an estimate or an index of the available nutrient-supplying capacity of the soil. This index can then be used to develop recommendations for nutrient applications based on plant need and the contribution of nutrients already in the soil to the crop nutrient requirement during the growing season. This best management practice (BMP) helps farmers achieve profitable crop yields while protecting the environment from excessive fertilization and nutrient losses. Successful use of soil testing requires that: (1) you send soil samples that adequately represent your field or management unit to the lab, (2) the laboratory uses calibrated soil test methods for predicting fertility

requirements, and (3) the fertilizer recommendations you get are based on measured crop responses.

The ESTL extracts phosphorus (P), potassium (K), Mg, Ca, copper (Cu), manganese (Mn), and zinc (Zn) from soil samples with the Mehlich-3 extractant and bases fertilizer recommendations for those nutrients on the test results. The use of the Mehlich-3 extractant enhances the extraction of micronutrients and allows for a broader range of applicability (most normal soil pH ranges) when compared to the Mehlich-1 extractant. Other advantages of the Mehlich-3 extractant include improved extraction of P in soils with high iron and aluminum (Al) accumulations by facilitating the dissociation of P from these compounds through the addition of fluoride, as well as the extraction of exchangeable cations by the addition of ammonium nitrate.

Nitrogen (N) fertilization is not based on soil tests but on crop need documented in research literature. Liming recommendations are based on the Adams-Evans lime requirement test (a calibration equation developed for Florida soils) and the target pH for the crop for which the recommendation is being made.

Soil Sampling

Soil testing depends on soil samples that are representative of the soil for the field in question. Samples should be collected from a relevant depth at specific locations to obtain a

1. This document is CIR1152, one of a series of the Soil and Water Sciences Department, UF/IFAS Extension. Original publication date March 2000. Revised October 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Rao Mylavarapu, professor, Soil and Water Sciences Department; George Hochmuth, professor emeritus, Soil and Water Sciences Department; and Guodong Liu, assistant professor, Horticultural Sciences Department; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

total number of samples that represent the conditions of the field.

Samples should be collected from the top six inches of the soil, because this is the part of the soil that is typically tilled and contains many of the nutrient-absorbing roots. The number of samples taken depends on the number of management units within the field. Management units are areas that will receive different agricultural practices (different crops or different planting dates) or have different soil types. These differences potentially contribute to different fertilization needs and management approaches. Once management units have been determined based on variability in the field, each management unit should be sampled separately. Samples should be collected from each management unit using a random approach to sample location selection in order to obtain a total of 20 samples. Each sample should be collected with a soil sampling probe and retained in a plastic bucket to composite all of the samples. Once sampling of the management unit is complete and the soils in the bucket are thoroughly mixed together, a sample volume of about one-half pint is obtained from the bucket and placed in the paper bag provided for soil-testing submissions to the lab. Additional information on management units and soil sampling schemes can be found in the EDIS document SL190, UF/IFAS Nutrient Management Series: Soil Sampling Strategies for Precision Agriculture (<http://edis.ifas.ufl.edu/ss402>) (Mylavarapu and Lee 2014).

Water Management

Nutrients can leach in Florida's sandy soils due to heavy rainfall or excessive irrigation. The fertilization recommendations presented in this circular were developed from research and on-farm experience with optimum water management. Irrigation requirements are primarily determined by crop water requirements, the characteristics of the irrigation system, management practices, and the physical and certain chemical characteristics of the soil in the irrigated area. Irrigation water quality criteria should also be considered in relation to fertilization, including salinity of the irrigation water, sodium adsorption ratio - i.e., the ratio of sodium to calcium (Ca) and magnesium (Mg) (Reeve et al. 1954; Oster and Sposito 1988), pH, alkalinity, organic contaminants, and heavy metals. A well-managed irrigation program will use water that does not curtail the effects of fertilization and will keep water and nutrients in the root zone, where both inputs will benefit yield and vegetable quality while minimizing chances for negative environmental impact. For more information on water management, see AE260, Principles and Practices of

Irrigation Management for Vegetables (<http://edis.ifas.ufl.edu/cv107>).

Managing Soil pH

In the southeastern United States, most native mineral soils are naturally acidic and require management for increasing pH (via liming) and nutrients to maintain optimal soil fertility. Liming reduces soil acidity, thereby changing numerous soil parameters related to fertilization in order to improve the yield or quality of a crop. Additionally, many of the soils found in Florida have historically been managed extensively for pH and may not require immediate additional liming. However, there are regions in south Florida that have calcareous soils with as much as 90% free calcium carbonate on the soil surface, thereby limiting nutrient availability. In these soils, liming should be absolutely avoided and appropriate methods to lower the pH should be considered and implemented instead. Other factors that can affect soil pH, including the soils, crops produced, ecosystem, local BMPs implemented, and economics should be understood before soils are managed for pH.

While calibrated lime requirement tests are part of standard soil tests in this region, the affordability and availability of agricultural lime and amendments has led to a tendency to over-lime soils. Soil pH should not be lowered unless there is evidence that plant growth is being adversely affected by pH. When lime is applied, adequate time for the liming material to react with the soil and raise pH should be allowed for before additional management actions are taken. If liming is required, the level of dissolved calcium carbonate in irrigation water drawn from groundwater should be considered when determining lime application rates.

Benefits of liming with dolomitic limestone, when necessary, include: reduction of Al toxicity in mineral soils, additional Mg and Ca as nutrients, and increased availability of other nutrients such as P. Conversely, overliming can cause low-P stress, Mn deficiency, other micronutrient deficiencies, and plant physiological disorders. In areas with naturally occurring carbonates that contribute to high soil pH, application of elemental sulfur (or ammonium sulfate when soil pH is lower than 7.25) and micronutrients in a band is recommended in order to most practically avoid the adverse effects of high pH. Application of a nitrogen fertilizer that provides sufficient hydronium ions in the root zone during chemical or biological reactions in soil will lower pH gradually with long-term use. In some cases, when the cause of high soil pH is natural, there may not be a cost-effective mechanism for lowering the pH and crops with a tolerance for higher pH ranges should be grown. For

more information on pH management, see EDIS documents Liming of Agronomic Crops (SS-AGR-153) (<http://edis.ifas.ufl.edu/aa128>), Lowering Soil pH to Optimize Nutrient Management and Crop Production (SL437) (<http://edis.ifas.ufl.edu/ss651>), and Agricultural Soils of Florida (SL441) (<http://edis.ifas.ufl.edu/ss655>).

Major Fertilization Factors

Depending on the crop and soils, natural fertility may not provide adequate levels of all required nutrients for desired plant growth. Fertilizers are used to provide additional nutrients in order to achieve economical crop production. In order to attain adequate nutrients for crop production while minimizing the risk of loss of nutrients to the environment, attention must be given to the four major soil fertilization factors: *right source*, *right rate*, *right placement*, and *right timing*. These factors, known as the 4Rs, should be evaluated when reviewing soil testing results to develop a personalized, integrated approach to nutrient management that makes efficient use of fertilizer investment for crop production and for environmental protection. The rate of fertilizer is a part of the overall nutrient management program. The recommended rates have been determined to provide adequate nutrient amounts even under highest yield potentials. Rate recommendations may change depending on new research. The 4Rs are fully interdependent and linked in the cropping system—they must each be considered and work together in order to sustain the economy, society, and the environment. Considerations for selecting the right source may include the ease of application of a nutrient, the cost per unit of the nutrient, and the efficiency of the nutrient. Practicing the right rate of nutrient application is intrinsically linked to soil testing. Soil testing can determine the amount of fertilizer needed for the crop production that meets plant requirements and protects against nutrient losses. The right timing of nutrients takes into consideration the growth pattern of the specific crop and changes in nutrient demand during the growing season in order to meet the needs of growth while also minimizing the chance of leaching of nutrients. The right placement of nutrients involves determining where the plant will have the best access to the nutrients—primarily in the root zone or just ahead of the advancing root system—and using a nutrient placement approach (such as banding and broadcasting) to meet those needs. The right placement should also consider the nutrient being applied. For example, P can become unavailable in some soils when broadcast-applied and can accumulate over time in unavailable forms, and ammoniacal N can significantly volatilize when left on the surface of a soil with a pH higher than 7.3. Examples of how the 4Rs influence fertilization include

consideration of controlled release or organic fertilizers (right source), using soil testing results to determine the amount of fertilizer needed (right rate), application by banding or broadcasting depending on the type of crop and the development or spread of the root system (right placement), and anticipating changes in plant-specific growth and nutrient demand so that application can be timed to meet the needs of growth (right timing). For more information on the 4Rs, please refer to the EDIS document The Four Rs of Fertilizer Management (SL411) (<http://edis.ifas.ufl.edu/ss624>).

Soil Test Rating Level

The current UF/IFAS interpretation of the Mehlich-3 extractant is presented in Table 1. This soil test has been extensively researched to correlate the soil text index (the concentration of nutrients extracted from the soil sample) with crop yield. This index tells us the relative level of a nutrient that will likely contribute to the crop nutrient requirement during the growing season. The response curve of these data are used to partition the range of soil text index results into corresponding low, medium, and high indexes. Through the process of calibration, crop nutrient requirements are determined for soil test values that fall within the indexes. This interpretation scale has been verified by field research on growers' fields throughout Florida for vegetable production.

Interpretations of results for Mehlich-3 extractable micronutrients have been developed from experience and field testing with vegetables (Table 2). Because responses to micronutrients are commodity-specific, these interpretations should be used as guides only. Zinc, Cu, and Mn can build up with time since they are quite immobile in the soil. The decision to add micronutrients should include an accounting for all sources, such as fungicides and micronutrient content in irrigation water.

Soil Test Recommendations

Soil test reports from the ESTL are computer-generated from soil test data and crop information. Reports contain the results of the tests (soil pH and ppm extractable P, K, Mg, Ca, and Cu, Mn, and Zn, if requested), a rating of the P, K, and Mg (high to low), and a fertilization recommendation. The recommendation is composed of two parts: (1) the rates of N, P₂O₅, and K₂O fertilizer to apply and (2) footnotes that give important information about fertilization management, such as application timing, special crop requirements, etc. Soil testing should be performed

annually in most cases, with future sampling frequency based on several successive years of soil testing results.

Table 3 contains crop descriptions, target pH, and N, P₂O₅, and K₂O recommendations for each of the three soil test rating levels, for which footnotes will be printed for each of the crop reports and the references upon which the recommendations are based.

Recommended fertilizer rates have been determined from research using typical standard bed spacing (Table 5). These fertilizer rates are expressed on a “per acre” basis, which can be converted to pounds per 100 linear bed feet. For planting patterns other than the typical bed spacing, refer to Table 6 for the equivalent fertilizer application rate given in pounds of nutrient per 100 linear bed feet. Using Table 5 and Table 6 will ensure correct fertilizer application rate using variable planting patterns. For more information on the linear bed foot system, please see SL409 (<http://edis.ifas.ufl.edu/ss622>).

References

Dukes, M.D., L. Zotarelli, G.D. Liu, and E. H. Somonne. 2015. *Principles and Practices of Irrigation Management for Vegetables*. AE260. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ae260>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Eggplant in Florida*. CV228. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv228>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Pepper in Florida*. CV230. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv230>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Cucumber in Florida*. CV226. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv226>

Hochmuth, G., and E. Hanlon. 2014. *A Summary of N, P, and K Research with Potato in Florida*. CV233. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv233>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Snap Bean in Florida*. CV234.

Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv234>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Sweet Corn in Florida*. CV235. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv235>

Hochmuth, G., and E. Hanlon. 2016. *A Summary of N, P, and K Research with Watermelon in Florida*. CV232. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv232>

Hochmuth, G., and E. Hanlon. 2014. *A Summary of N, P, and K Research with Tomato in Florida*. CV236. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv236>

Hochmuth, G. J., and E. A. Hanlon. 2016. *Calculating recommended fertilizer rates for vegetables grown in raised-bed, mulched cultural systems*. SS516. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss516>

Hochmuth, G., R. Mylavarapu, and E. Hanlon. 2014. *Developing a Soil Test Extractant: The Correlation and Calibration Processes*. SS622. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss622>

Hochmuth, G., R. Mylavarapu, and E. Hanlon. 2014. *Fertilizer Recommendation Philosophies*. SS623. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss623>

Hochmuth, G., R. Mylavarapu, and E. Hanlon. 2014. *Soil Testing for Plant-Available Nutrients- What Is it and Why Do We Use It?* SS621. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss621>

Hochmuth, G., R. Mylavarapu, and E. Hanlon. 2014. *The Four Rs of Fertilizer Management*. SS624. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss624>

Liu, G., K. Morgan, Y. Li, L. Zotarelli, J. DeValerio, and Q. Wang. 2015. *What is 4R Nutrient Stewardship?* HS1264. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/hs1264>

Mylavarapu, R., G. Hochmuth, C. Mackowiak, A. Wright, and M. Silveira. 2016. *Lowering Soil pH to Optimize Nutrient Management and Crop Production*. SS651. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss651>

Mylavarapu, R., W.G. Harris, and G. Hochmuth. 2016. *Agricultural Soils of Florida*. SS655. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss655>

Mylavarapu, R., G. Hochmuth, C. Mackowiak, A. Wright, and M. Silveira. 2016. *Lowering Soil pH to Optimize Nutrient Management and Crop Production*. SS651. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss651>

Mylavarapu, R., T. Obreza, K. Morgan, G. Hochmuth, V. Nair, and A. Wright. 2014. *Extraction of Soil Nutrients Using Mehlich-3 Reagent for Acid-Mineral Soils of Florida*. SS620. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss620>

Mylavarapu, R., and D.L. Wonsuk. 2014. *UF/IFAS Nutrient Management Series: Soil Sampling Strategies for Precision Agriculture*. SS402. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss402>

Mylavarapu, R., D. Wright, and G. Kidder. 2015. *UF/IFAS Standardized Fertilization Recommendations for Agronomic Crops*. SS163. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/ss163>

Oster, J. D., and G. Sposito. 1980. "The Gapon coefficient and the exchangeable sodium percentage-sodium adsorption ratio relation". *Soil Science Society of America Journal*. 44(2): 258. doi:10.2136/sssaj1980.03615995004400020011x

Reeve, R.C., C.A. Bower, R.H. Brooks, and F.B. Gschwend. 1954. "A comparison of the effects of exchangeable sodium and potassium upon the physical condition of soils". *Soil Science Society of America Journal*. 18 (2): 130. doi:10.2136/sssaj1954.03615995001800020004x.

Wright, D.L., C. Mackowiak, and E.B. Whitty. 2014. *Liming of Agronomic Crops*. AA128. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/aa128>

Table 1. Mehlich-3 soil test interpretations used for vegetable crops on mineral soils.

| Element | Mehlich-3, mg/kg ¹ (ppm) | | |
|---------|-------------------------------------|--------|------|
| | Low | Medium | High |
| P | ≤25 | 26–45 | >45 |
| K | ≤35 | 36–60 | >60 |
| Mg | ≤20 | 21–40 | >40 |

Source: Mylavarapu, Obreza, Morgan, Hochmuth, Nair, and Wright (2014)

Table 2. Interpretations of Mehlich-3 soil test for micronutrients.

| Interpretations | Soil pH (mineral soils only) | | |
|---|--------------------------------|-----------|-----------|
| | 5.5–5.9 | 6.0–6.4 | 6.5–7.0 |
| | Test level (parts per million) | | |
| Test level below which there may be a crop response to applied copper. | 0.7–1.0 | 1.0–1.3 | 1.3 |
| Test level above which copper toxicity may occur. | 3.5–5.0 | 5.0–8.0 | 8.0 |
| Test level below which there may be a crop response to applied manganese. | 10.3–12.7 | 12.7–15.1 | 15.1–17.6 |
| Test level below which there may be a crop response to applied zinc. | 1.7 | 1.7–2.2 | 2.2–4.2 |

From “Notes in Soil Science” No. 9 1983, and Mylavarapu et al. 2002.

Table 3. Target pH and recommended N, P₂O₅, and K₂O fertilizer rates for commercial vegetable production. Phosphorus and potassium rates are based on an interpretation of a Mehlich-3 soil test on a mineral soil.

| Crop Description | Target pH | Pounds/Acre/Cropping season | | | | | | | Footnotes | References |
|--|-----------|-----------------------------|-------------------------------|-----|----|------------------|-----|----|---------------------------------|-----------------------------|
| | | N (lb/Ac) | P ₂ O ₅ | | | K ₂ O | | | | |
| | | | L | MED | HI | L | MED | HI | | |
| BEAN Snap, Lima, Pole | 6.5 | 100 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 356 | HS853, HS1261, HS711, CV296 |
| BEET | 6.5 | 120 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 356 | HS711, CV296 |
| BROCCOLI Cauliflower, Brussels sprouts | 6.5 | 175 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 352 354 356 | HS711, CV296 |
| CABBAGE Collard, Chinese Cabbage | 6.5 | 175 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 352 354 356 | HS854, HS711, CV296 |
| CARROT | 6.5 | 175 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 356 | HS711, CV296 |
| CELERY | 6.5 | 200 | 150 | 100 | 0 | 150 | 100 | 0 | 250 251 350 354 356 | HS711, CV296 |
| CUCUMBER | 6.5 | 150 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 351 354 356 | HS855, HS711, CV296 |
| EGGPLANT | 6.5 | 200 | 130 | 100 | 0 | 130 | 100 | 0 | 250 251 350 351 352 353 354 356 | HS856, HS711, CV296 |
| LETTUCE Crisphead, Romaine, Endive, Escarole | 6.5 | 200 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 352 354 356 | HS711, CV296 |
| MUSKMELON | 6.5 | 150 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 354 356 | HS711, CV296 |
| MUSTARD Kale, Turnip | 6.5 | 120 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 356 | HS711, CV296 |
| OKRA | 6.5 | 120 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 356 | HS857, HS711, CV296 |

| Crop Description | Target pH | Pounds/Acre/Cropping season | | | | | | | Footnotes | References |
|--------------------------------|-----------|-----------------------------|-------------------------------|-----|----|------------------|-----|-----|------------------------------------|--|
| | | N (lb/Ac) | P ₂ O ₅ | | | K ₂ O | | | | |
| | | | L | MED | HI | L | MED | HI | | |
| ONION Bulb | 6.5 | 150 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 354 356 | HS711, CV296 |
| ONION Bunching, Leek | 6.5 | 120 | 100 | 100 | 0 | 100 | 100 | 0 | 250 251 350 356 | HS711, CV296 |
| PARSLEY | 6.5 | 120 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 356 | HS638, HS711, CV296 |
| PEA Southern, Snow, English | 6.5 | 60 | 80 | 60 | 0 | 80 | 60 | 0 | 250 251 350 356 | HS711, CV296 |
| PEPPER Bell, Specialty | 6.5 | 200 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 352 353 354 356 | HS859, HS711, CV296 |
| POTATO | 6.0 | 200 | 120 | 60 | 0 | 150 | 150 | 150 | 250 253 350 356 | HS945, HS711, CV296 |
| RADISH | 6.5 | 90 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 252 350 356 | HS711, CV296 |
| SPINACH | 6.5 | 90 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 356 | HS711, CV296 |
| SQUASH Summer, Winter, Pumpkin | 6.5 | 150 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 356 | HS861, HS711, CV296 |
| STRAWBERRY | 6.5 | 150 | 120 | 100 | 0 | 120 | 100 | 0 | 250 350 352 353 354 355 356 | HS956, HS1116, HS711, SL344, CIR1141, CV296 |
| SWEET POTATO | 6.5 | 60 | 100 | 80 | 0 | 100 | 80 | 0 | 250 251 350 356 | HS711, CV296 |
| TOMATO Slicing, Cherry, Plum | 6.5 | 200 | 120 | 100 | 0 | 125 | 100 | 0 | 250 251 350 351 352 353 354 356 | HS858, HS711, CV296 |
| WATERMELON | 6.0 | 150 | 120 | 100 | 0 | 120 | 100 | 0 | 250 251 350 351 352 353 354 356 | HS711, CV296 |

Table 4. Footnotes used with vegetable crops.

| |
|---|
| <p>250 Indicated fertilizer amounts, and the nutrients already in the soil, will satisfy the crop nutrient requirement for this cropping season. Fertilizer and water management are linked. Maximum fertilizer efficiency is achieved only with close attention to water management. Supply only enough irrigation water to satisfy crop requirements. Excess irrigation may result in leaching of N and K, creating possible plant deficiencies. Overfertilization has been shown to reduce vegetable quality. For subsurface irrigation, maintain a constant water table between 18 (at planting) and 24 inches (near harvest) below the top of the bed. Monitor water table depth and do not fluctuate, else N can be “scrubbed” from the root zone. On soils that have not been in vegetable production within the past 2 years, or where micronutrients are known to be deficient, apply 5 lb Mn, 3 lb Zn, 4 lb Fe, 3 lb Cu, and 1.5 lb B/A. Use soil testing to monitor micronutrient status every 2 years to avoid micronutrient toxicity, since some micronutrients can build up in the soil. When deciding about micronutrient applications, consider micronutrients added to the crop via fungicides. Up to 40 lb/acre Mg might be needed when soil test is medium or lower in Mg. Mg can be supplied in fertilizer or from dolomitic limestone, when liming is recommended. Calcium concentrations are typically adequate in most soils used continuously for vegetable production or where the Mehlich-3 Ca index is > 300 ppm. Calcium is added during liming activities and from calcium carbonate present in irrigation water drawn from aquifers in Florida. These sources should be considered in the determination of Ca fertilizer needs.</p> |
| <p>251 For unmulched crops, fertilizer should be applied in split applications to reduce leaching losses and lessen danger of fertilization burn. Broadcast all P₂O₅ and micronutrients, if any, and 25 to 30% of the N and K₂O in the bed at planting. Apply remaining N and K₂O in sidedress bands during the early part of the growing season. Additional, supplemental sidedress applications of 30 lb N/A and 20 lb K₂O/A should be applied only if rainfall/irrigation amounts exceed 3 inches within a 3-day period or exceed 4 inches within a 7-day period. Avoid mechanical damage to plants when applying fertilizers.</p> |
| <p>252 The amounts suggested are generally sufficient for 2 or 3 crops in succession.</p> |
| <p>253 Where scab-resistant cultivars are grown, a pH between 6.0 and 6.5 is optimum. Where scab-susceptible cultivars are grown, the pH should be below 5.2 or above 7.2. Band all phosphorus. Apply 50 to 70% of N and 50% of K₂O at emergence and the remaining N and K at 35 to 40 days after planting. Potatoes planted in cool soils might respond to up to 25 lb P₂O₅ applied as starter fertilizer in the furrow with the seed pieces.</p> |
| <p>350 Supply 25 to 50% of the N in the nitrate form if soils were treated with multi-purpose fumigants or if the soil temperature will stay below 60°F for up to one week following transplanting or germination.</p> |
| <p>351 For mulched crops and subsurface irrigation, incorporate 10 to 20% of the N and K₂O, plus all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O 2 to 3 inches deep in one or more bands about 6 to 10 inches from the plants. For drip irrigation, incorporate 20% to 40% of the N and K₂O and all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O periodically through drip tubes according to the rate or crop growth. Consult AE259, “Scheduling Tips For Drip Irrigation of Vegetables”, AE260, “Principles and Practices of Irrigation Management for Vegetables” and AE500, “How to Determine Run Time and Irrigation Cycles for Drip Irrigation: Tomato and Pepper Examples” for information on injection schedules. For management systems where both subsurface and drip irrigation are being used, apply no more than 20% of the N and K₂O, plus all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O periodically through drip tubes according to the rate of crop growth. For overhead irrigation, incorporate all of the N, P₂O₅, K₂O and micronutrients, if any, into the bed prior to installation of the plastic mulch.</p> |
| <p>352 Amounts suggested are for the first crop. Squash and cucumber following other crops on the same mulch may not need substantial additional fertilizer. If fertilizer is needed for the second crop, apply fertilizer using a liquid-injection wheel or via drip irrigation. Apply no more than 30 to 40 lb/acre N and/or K₂O in any single injection wheel application.</p> |
| <p>353 From 25 to 30% of the N may be supplied from slow-release N sources, such as sulfur-coated urea, polymer-coated fertilizers, or isobutylidene-diurea (IBDU).</p> |
| <p>354 Transplants may benefit from application of a dilute, soluble starter fertilizer, especially at cool soil temperatures. Starter solution rates of N and P₂O₅ need not exceed 10 to 15 lb/acre each.</p> |
| <p>355 For overhead irrigation, broadcast all the P₂O₅ and micronutrients, if any, and 25% of the N and K₂O into the bed. Band remaining N and K₂O in center of bed 3 inches deep. For subsurface irrigation, incorporate 10 to 20% of the N and K₂O, plus all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O 2 to 3 inches deep in one or more bands about 6 to 10 inches from the plants. For drip irrigation, incorporate 20% of the N and K₂O and all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O periodically through drip tubes according to the rate of crop growth; see AE354, “Automatic Irrigation Based on Soil Moisture for Vegetable Crops”, AE500, “How to Determine Run Time and Irrigation Cycles for Drip Irrigation: Tomato and Pepper Examples”, AE260, “Principles and Practices of Irrigation Management for Vegetables” and Circular 1141, “Fertilization of Strawberries in Florida.” For management systems where both subsurface and drip irrigation are being used, apply no more than 20% of the N and K₂O, plus all of the P₂O₅ and micronutrients, if any, into the bed. Apply the remainder of the N and K₂O periodically through drip tubes according to the rate of crop growth.</p> |
| <p>356 UF/IFAS fertilization and liming recommendations are advisory in nature and emphasize efficient fertilizer use and environmentally sound nutrient management without losses of yield or crop quality. It is generally assumed the nutrients will be supplied from purchased commercial fertilizer and the expected crop yields and quality will be typical of economically viable production. Growers should consider UF/IFAS recommendations in the context of their entire management strategy, such as return on investment in fertilizer and the benefits of applying manure or biosolids (sewage sludge) to their land. There is insufficient research available to support the use of UF/IFAS soil test results for environmental nutrient management purposes. Such use is discouraged until correlation is proven.</p> |

Table 5. Typical bed (row) spacings for vegetables.

| Crop | Bed (row) spacing | Number of rows (per bed) |
|--|-------------------|--------------------------|
| Bean, snap, lima | 30 inches | 1 |
| Broccoli, cauliflower, Brussels sprout | 6 ft (mulched) | 2 |
| Cabbage, collard, Chinese cabbage, kale | 6 ft (mulched) | 2 |
| Carrot | 4 ft | 2-3 |
| Celery | 4 ft | 2 |
| Cucumber | 6 ft (mulched) | 2 |
| Eggplant | 6 ft (mulched) | 1 |
| Lettuce, crisphead, romaine, endive, escarole | 4 ft | 2 |
| Muskmelon | 5 ft | 1 |
| Okra | 6 ft (mulched) | 2 |
| Onion | 6 ft | 4 |
| Pea, southern | 30 inches | 1 |
| Pepper, bell, specialty | 6 ft (mulched) | 2 |
| Potato | 42 inches | 1 |
| Squash, summer | 6 ft (mulched) | 2 |
| Strawberry | 4 ft (mulched) | 2 |
| Sweet corn | 36 inches | 1 |
| Sweet potato | 42 inches | 1 |
| Tomato, slicing, cherry, plum | 6 ft (mulched) | 1 |
| Watermelon | 8 ft | 1 |
| For the following crops, see footnote¹ | | |
| Mustard | | |
| Turnip | | |
| Parsley | | |
| Pea, snow, English | | |
| Radish | | |
| Spinach | | |

¹These crops are generally produced on wide (40 to 48-inch) beds on 6-ft centers with 4 to 6 multiple rows. Some of the crops are also sown in broadcast-fashion on the bed.
Source: Hochmuth and Hanlon, SL409, 2016.

Table 6. Conversion of fertilizer rates in lb/A to lb/100 linear bed ft (LBF)¹.

| Bed spacing (ft) | Recommended fertilizer (N, P ₂ O ₅ , or K ₂ O) | | | | | | | | | | | | |
|--|---|------|------|------|------|------|------|------|------|------|------|------|------|
| | ----- lb/A ----- | | | | | | | | | | | | |
| | 20 | 25 | 40 | 50 | 60 | 75 | 80 | 100 | 120 | 140 | 160 | 180 | 200 |
| Pounds of fertilizer (N, P ₂ O ₅ , or K ₂ O) to apply per 100 LBF | | | | | | | | | | | | | |
| 3 | 0.14 | 0.17 | 0.28 | 0.35 | 0.41 | 0.52 | 0.55 | 0.69 | 0.83 | 0.96 | 1.10 | 1.24 | 1.38 |
| 4 | 0.18 | 0.23 | 0.37 | 0.46 | 0.55 | 0.69 | 0.73 | 0.92 | 1.10 | 1.29 | 1.47 | 1.65 | 1.84 |
| 5 | 0.23 | 0.29 | 0.46 | 0.57 | 0.69 | 0.86 | 0.92 | 1.15 | 1.38 | 1.61 | 1.84 | 2.07 | 2.30 |
| 6 | 0.28 | 0.34 | 0.55 | 0.69 | 0.83 | 1.03 | 1.10 | 1.38 | 1.65 | 1.93 | 2.20 | 2.48 | 2.77 |
| 8 | 0.37 | 0.46 | 0.73 | 0.92 | 1.10 | 1.38 | 1.47 | 1.84 | 2.20 | 2.57 | 2.94 | 3.31 | 3.67 |

¹This table is used correctly by (1) determining the typical bed spacing from Table 5 for the crop; (2) locating the column containing the recommended fertilizer rate in pounds per acre; and (3) reading down the column until reaching the row containing the typical bed spacing. The resulting number in pounds per 100 LBF should be used even in situations where the farmer's bed spacing differs from the typical bed spacing. Use of the table will involve doubling the rate, for example where the column for 100 pounds per acre was used in the calculation of pounds per 100 LBF for a recommended rate of 200 pounds per acre.
Source: Hochmuth and Hanlon, SL409, 2016.