

# Chapter 24.

## Tropical Root Crop Production in Florida

M.L. Lamberts and S.M. Olson

### INTRODUCTION

There are four main tropical root crops produced in Florida. These are: cassava, tannia, taro and boniato. The following table lists the botanical families, scientific names and both English and Spanish common names for these crops.

English Common Name	Family	Scientific name	Spanish Common Name(s)
Cassava	Euphorbiaceae	<i>Manihot esculenta</i> Crantz	Yuca
Taro, eddoe, dasheen	Araceae (Aroid)	<i>Colocasia esculenta</i> (L.) Schott	Malanga isleña or malanga
Tannia or tanier	Araceae (Aroid)	<i>Xanthosoma sagittifolium</i> (L.) Schott	Malanga or yautía
Tropical Sweetpotato	Convolvulaceae	<i>Ipomoea batatas</i> (L.) Lam.	Boniato or batatas or camote

### BRIEF DESCRIPTIONS OF TROPICAL ROOT CROPS

*Cassava* - a short-lived perennial tropical shrub growing 3 to 12 ft. tall (Fig. 24-1). Adventitious roots arise from stem cuttings. They vary in shape from long and slender to globose and enlarge during the starch storage process. Stems are woody and variously branched. Simple leaves, generally dark green, have palmate lobes.

*Taro* - produces an enlarged edible corm. Its leaves are peltate and when viewed from the abaxial surface resemble a heart with a heavily pigmented spot at the point where the petiole is attached to its adaxial surface (Fig. 24-2).

*Tannia* - produces an enlarged corm that may or may not be edible. Underground club-shaped cormels are produced during later growth stages (Fig. 24-3). Leaves have long petioles and broad sagittate lamina (Fig. 24-4).

*Tropical sweetpotato* - resembles other sweetpotatoes grown in the United States, but the skin ranges in color from light pink to a deep wine red and the flesh is white and starchier than yellow- or orange-fleshed cultivars.

### VARIETIES

*Cassava* - varieties are often separated based of their cyanogenic glucoside (HCN) content into either low HCN, a.k.a. "sweet," or high HCN, a.k.a. "bitter," types. The term "bitter" comes from a bitter flavor that is commonly believed to accompany the HCN. No named varieties are currently known in southern Florida. 'Senorita' was locally popular in the 1980s and attempts were made to introduce the CIAT variety 'Mantiqueira' at that same time because it was able to produce acceptable yields even with high levels of rootknot nematode. The range of local genotypes covers a few unnamed clones that have been imported from various Caribbean basin countries. Federal regulations prohibit further importation of cuttings or botanical seed.

*Taro* - 'Malanga Isleña,' produces one large white-fleshed central corm; a few unnamed Polynesian types are grown for the Asian market.

*Tannia* - 'South Dade White,' producing white-fleshed cormels; 'Malanga Amarilla,' producing a yellow-fleshed edible corm; and 'Vinola,' producing purple-fleshed cormels.

*Tropical sweetpotato* - 'Picadito' is the main variety grown in Miami-Dade County. It has deeply lobed leaves and wine colored skin (Fig. 24-5).

### PLANTING AND SEEDING

*Cassava* - In Miami-Dade County, theoretically it can be planted year round (Table 1). Normally, the planting is done in the early spring due to slow early growth and

**Table 1.** Planting information for cassava.

Planting dates	
Miami-Dade Florida	Year-round (mainly February to April)
Planting information	
Distance between rows (in)	48
Distance between plants (in)	24
Planting depth (in)	3 - 4
Propagules needed per acre	5,445
Days to harvest from planting	270 - 360
Plant populations (acre)	5,445

**Table 2.** Planting information for tannia and taro.

Planting dates	
Miami-Dade Florida	Year-round (mainly February to October)
Planting information	
Distance between rows (in)	48
Distance between plants (in)	24
Planting depth (in)	3 - 4
Propagules needed per acre	5,445
Days to maturity from transplant	270 - 360
Plant populations (acre)	5,445

the possibility of a winter frost. Stem cuttings 4–5” long (Fig. 24-6) are selected from the lower mature sections of healthy plants (Fig. 24-7) and planted horizontally, 3-6” below the soil surface in shallow Rockdale soils. Cuttings are planted by hand in furrows spaced 48” apart with an in row spacing of 24”. Since unrooted cuttings are sensitive to water loss, the field should be irrigated soon after planting.

*Taro & Tannia* - In Miami-Dade County, both crops can be planted year round. Normally, the colder winter months are avoided due to slow growth and susceptibility to frost. For tannia, the top portion of the mother corm is used for planting., For taro, the smaller unmarketable side shoots are used. Both are planted by hand in furrows spaced 48” apart with an in row spacing of 24”.

*Tropical sweetpotato* – In Miami-Dade County, this crop can be planted year round, though planting is sometimes delayed if extremely cold weather has been forecast. As with orange- or yellow-fleshed varieties, vine cuttings are used for planting.

## FERTILIZER & LIME

*Cassava* – Apply all  $P_2O_5$ , micronutrients, and 25 to 50% of N and  $K_2O$  in a band along the planted row one to two months after planting. Sidedress the remaining N and  $K_2O$  by banding them on the side of the bed four months after planting as a layby. Specific fertilizer recommendations for Rockdale soils are not available due to lack of a reliable and readily available soil analysis procedure for this soil type. In general, cassava has a medium requirement for N,  $P_2O_5$  and  $K_2O$ . Since the soils in Miami-Dade have a high pH, liming is not needed. There are no remediation materials for the high pH condition, though chelated forms of micronutrients such as Fe, Mg, Mn and Zn can be applied to treat deficiencies.

*Taro & Tannia* – Apply all  $P_2O_5$ , micronutrients, and 25 to 50% of N and  $K_2O$  in a band along the planted row one to two months after planting. Sidedress the remaining N and  $K_2O$  by banding them on the side of the bed four months after planting as a layby. Specific fertilizer recom-

mendations Rockdale soils are not available due to lack of a reliable and readily available soil analysis procedure for this soil type. In general, taro and tannia have medium to high requirements for N, medium to low requirements for  $P_2O_5$  and medium requirements for  $K_2O$ .

*Tropical sweetpotato* – see Chapter 22, Sweetpotato Production in Florida.

## IRRIGATION

*Cassava* - has relatively low requirements for irrigation. But, during active growth stages it must be irrigated when extended dry spells occur. Water requirements and subsequent irrigation requirements are reduced during the last few weeks of growth. This usually coincides with the dry season when cassava is most likely to be affected by mites, a situation which can be helped by using overhead irrigation since it increases humidity in the field.

*Taro & Tannia* - grow best when the soil is maintained moist, but not wet, at all times. Proper water management is essential for optimum corm and cormel sizing. Water requirements and subsequent irrigation requirements may be reduced during the last few weeks of growth.

*Tropical sweetpotato* – see Chapter 22, Sweetpotato Production in Florida.

## CULTURAL PRACTICES

### Soil Preparation

*Cassava* - grows best when the soil is turned 2 to 3 months before planting. Plowing early helps rot plant debris and reduce some nematode and disease problems. Soils in Miami-Dade County (except for marl soils) should be scarified or “rock plowed” prior to planting to improve drainage and increase available soil depth. For cassava, marl soils will always produce the best-shaped and best-looking storage roots. However, these soils are prone to flooding, making them less desirable for cassava production. Plants grown on Rockdale soils must be irrigated during periods of dry weather to avoid reduced yields. Wet weather for extended periods can cause leaching of N and K, requiring the addition of more fertilizer. More frequent applications of smaller amounts of fertilizer per application are suggested for Rockdale soils. In general, fields that have not produced a crop of cassava in the last 2 to 3 years are preferred. Avoid fields that have very high nematode populations or use a resistant variety if one is legally available.

*Taro & Tannier* - grow best when the soil is turned 2 to 3 months before planting. Plowing early helps rot plant debris and reduce some nematode and disease problems.

Soils in Miami-Dade County (except for marl soils) should be scarified or “rock plowed” prior to planting to improve drainage and increase available soil depth. For tannia, marl soils will always produce the best-shaped and best-looking cormels. These soils are preferred if supplemental irrigation cannot be supplied. Plants grown on Rockdale soils must be irrigated during periods of dry weather to avoid reduced yields. Wet weather for extended periods can cause leaching of N and K, requiring the addition of more fertilizer. More frequent applications of fertilizer using smaller amounts each time are suggested for Rockdale soils. In general, fields that have not produced a crop of tannia or taro in the last 2 to 3 years are preferred. Avoid fields that have very high nematode populations.

*Tropical sweetpotato* - see Chapter 22, Sweetpotato Production in Florida.

### Bedding

*Cassava, Taro & Tannia* – Plants are established in rows on flat land. During the cultivation process, ridges are formed down the rows of plants. Ridges provide a place for storage organ formation, improve drainage and facilitate harvesting.

*Tropical sweetpotato* – see Chapter 22, Sweetpotato Production in Florida.

### Cover Crops

*Taro & Tannier* – Cover crops are not recommended immediately after tannia and taro production. Cultivation is needed after harvest to prevent feral plant establishment.

*Tropical sweetpotato* – see Chapter 22, Sweetpotato Production in Florida.

### Disease Management

*Cassava* – There are several virus diseases of cassava. However, none have been reported to occur in the United States. Systemic diseases are also common in cassava. Planting material should pass through a phase of tissue culture and thermo therapy. Cassava bacterial blight (see PP-40: edis.ifas.ufl.edu/VH053) and *Cercospora* leaf spot are known disease problems in the United States. When pathogens are present, yields can be reduced by 50% if healthy, pathogen-free or disease resistant cuttings are not used. This emphasizes the need to select clean, healthy planting material. Since cassava is vegetatively propagated, virus and other systemic diseases can be carried from one

**Table 3.** Fungicides approved for disease management on cassava, taro and tannia.

Chemical Name	Rate/acre	Minimum Days to Harvest	Comments
Acrobat 50 WP	6.4 oz	30	Suppression of Phytophthora rot and leaf blight, not for cassava
Actinovate AG	12 oz	0	Biological material
Azoxystrobin (several brands)	See label	14	See label for maximum use rate and restrictions
Apron XL LS	0.64 oz/cwt.	0	Apply to seed pieces
Evito 480 SC	3.8 oz	7	See label for maximum use rate and restrictions
Headline	12 oz	3	See label for maximum use rate and restrictions
Maxim 4 FS	0.16 oz/cwt.	0	Apply to seed pieces
Metalaxyl (several brands)	4 pts	0	apply preplant or shortly after planting
Oxidate	1/100	0	
Potassium phosphite (several brands)	See label	0	
Presidio	4 oz	7	See label for maximum use rate and restrictions
Reason 500 SC	8.2 oz	14	See label for maximum use rate and restrictions
Revus	8 oz	14	See label for maximum use rate and restrictions
Ridomil Gold EC	See label		apply preplant or shortly after planting
Scala SC	7 oz	7	
Serenade ASO, MAX	See label	0	
Sonata	2-4 qts	0	
Switch 62.5 WG	14 oz	7	See label for maximum use rate and restrictions
Trilogy	2 gal	0	

planting to another in the planting material. Effective disease control for cassava is based on prevention. Most of the important diseases are caused by pathogens that can be easily spread by wind, rain and workers or are capable of spreading systemically through the plant. It is usually not possible to restore the health of an affected plant once the disease can be detected. A crop rotation of at least 3 years is an important means of controlling diseases. Chemicals approved for management of cassava diseases are shown in Table 3.

*Taro & Tannia* – Dasheen mosaic virus (Fig. 24-8; (see PP-40: [edis.ifas.ufl.edu/VH053](http://edis.ifas.ufl.edu/VH053))) and systemic diseases are common in tannia and taro. Planting material should pass through a phase of tissue culture and thermo therapy. Bacterial leaf blight (see PP-40: [edis.ifas.ufl.edu/VH053](http://edis.ifas.ufl.edu/VH053))

and root rot are two other maladies affecting tannia. Taro has few known disease problems in the United States. Losses due to these diseases have not been documented, but they can be significant. Genetic resistance to dasheen mosaic virus and root rot diseases is not well documented. Effective disease control for tannia and taro are based on prevention. Most of the important diseases are caused by root pathogens or are capable of spreading systemically through the plant. It is generally not possible to restore the health of an affected plant once the disease can be detected. Since taro and tannia are vegetatively propagated, virus and other systemic diseases can be carried from one planting to another in the propagules. A crop rotation of at least 3 years is an important means of controlling diseases. Chemicals approved for management of tannia and taro diseases are listed in Table 3.

**Table 4.** Selected insecticides for insect management on cassava, taro and tannia.

Chemical Name	Rate/acre	Minimum Days to Harvest	Comments	Notes
Actara	1.5 oz	14	leafhoppers	
Assail 30 SG, 70 WP	see label	7	aphids, leafhoppers, flea beetles	
Avaunt	6 oz	7	various caterpillars	
Azadirachtin (many brands)	see label	0	whiteflies, hornworms	
Baythroid XL	2.8 oz	0	various caterpillars, flea beetles	
Bifenthrin (various brands)	See label	See label	cucumber beetles, wireworms, flea beetles, white grubs	may be applied to soil prior to planting
BT ( <i>bacillus thuringiensis</i> )	see label	0	various caterpillars	
Deltamethrin (various brands)	See label	3	various caterpillars, flea beetles, green peach aphid	
Fulfill	2.75 oz	14	aphids, whiteflies	
Imidacloprid (many brands)	see label	125	aphids, leafhoppers, whiteflies	soil application at planting
Imidacloprid (many brands)	see label	7	aphids, leafhoppers, whiteflies	foliar applied
Intrepid 2F	6-16 oz	14	armyworms, cutworms, loopers, webworms	
Oberon 2SC	8-16 oz	7	whiteflies two spotted spider mites	
Oil, insecticidal	1-2 gal/100 gal water	1	leafminers, mites, whiteflies	
Renounce 20WP	1-3.5 oz	0	cutworms, loopers, flea beetles, leafhoppers	
Soap, insecticidal	1-2 gal/100 gal water	0	aphids	
Spinosad (various brands)	See label	7	Various caterpillars, dipteran leaf-miners, thrips	See label for detailed use instructions

*Tropical sweetpotato* – see Chapter 22, Sweetpotato Production in Florida.

### **Insect Management**

*Cassava* – Whitefly, spider mites, shoot fly and tomato hornworm are occasionally observed. Since there is no documentation of economic loss due to these pests, pesticides (Table 4.) are not normally used. Although mites can cause leaf drop, they are normally not present in significant numbers until the winter months when the plants nearing harvest. Keeping fallowed fields free of feral plants is recommended.

*Taro & Tannia* – Diaprepes root weevil is the main pest of tannia. Larval feeding causes physical damage to the cormels, causing them to be placed in a lower grade. Whitefly, thrips, and salt marsh caterpillar are occasionally observed. However, there is no documentation of economic loss due to these pests. Resistance is not known to any of these pests. Keeping fallowed fields free of feral plants is recommended. Table 4 lists chemicals labeled for taro and tannia.

*Tropical sweetpotato* — see Chapter 22, Sweetpotato Production in Florida.

### **Nematode Management**

*Cassava* – can be damaged by root-knot nematodes. It may cause stunting and yield loss; root-knot nematodes in the storage roots may cause cracking or internal dark lesions that severely reduce the value of the product. The only step to minimize nematode injury to cassava is to include crop rotation unless nematode-tolerant varieties can be introduced legally.

*Taro & Tannia* – Taro is frequently damaged by root-knot nematodes. Nematodes may cause stunting and yield loss; root-knot nematodes in the corms may cause cracking or internal dark lesions that severely reduce the value of the product. Several steps to minimize nematode injury to taro include crop rotation and use of nematode-free propagules. Using a field immediately after a crop that has legally been treated for nematodes may reduce injury.

*Tropical sweetpotato* — see Chapter 22, Sweetpotato Production in Florida.

### **EPA Crop Grouping and labels**

The Environmental Protection Agency establishes tolerances for pesticide residues in raw agricultural commodities. The crop grouping scheme allows tolerance to be established on a large number of commodities from research on certain specific crops in that group. The tropical root crops fall under the “Tuberous and Corm vegetable” subgroup of the “Root and Tuber Vegetable” group.

This means that if a label states that the pesticide is labeled for root and tuber crops, or for tuberous and corm vegetables, it may legally be applied to the tropical root vegetables including: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, cassava, chayote root, chufa, dasheen (taro), ginger, leren, sweet potato, tanier, taro, tumeric, yam bean, and true yam.

Check the labels of the pesticides to see if the crop group is listed.

### **Weed Management**

The following herbicides are labeled under the “tuberous and corm vegetable” crop group.

Clomozone (Command 3 ME) Apply preemergence or preplant at 2-3.3 pints/A. Trials with sweet potatoes and true yams has shown good tolerance and weed control. Command has not been tried on other tropical root crops in Florida. Use on a trial basis only.

Sethoxydim (Poast). Controls emerged grasses. Apply at 1 to 2.5 pints/A. Add a crop oil concentrate (coc) at 1% v/v. There is a 30 day phi.

Glyphosate (Roundup Original Max, Roundup UltraMax II, Roundup UltraMax TR, Roundup UltraDry, Roundup WeatherMax) may be applied as a directed shielded spray to row middles at 0.5 to 1.0 lb ai/A. Direct contact with the crop will cause significant yield reduction. (Note to Steve: Dow products have not been included since these crops are not listed on the labels I checked)

Carfentrazone (Aim) May be applied as a directed hooded spray in row middles at 0.33 to 1 fluid ounce/A. Controls a wide range of broadleaf weeds. Does not control grass weeds. Add a non-ionic surfactant or a COC at recommended rates.

*Cassava* – is long-season crop. Control of weeds during production can be difficult. Early season competition from weeds is extremely detrimental to crop yield, so a major emphasis on weed control should be made during this period. Growers must plan a total program that integrates mechanical and cultural methods of weed control with the use of herbicides. Cultivation is an effective way to manage weeds early in the season. Hilling blades can uproot many annual weeds that have emerged since the last cultivation.

*Taro & Tannia* – are long-season crops. Control of weeds during the extended production period can be difficult. Early season competition of weeds is extremely detrimental to crop yield, so a major emphasis on weed control should be made during this period. Growers must plan a

total program that integrates mechanical and cultural methods of weed control with the use of herbicides. Cultivation is an effective way to manage weeds early in the season. Hilling blades can uproot many annual weeds that have emerged since the last cultivation.

*Tropical sweetpotato* — see Chapter 22, Sweetpotato Production in Florida.

### Harvesting / Packing

*Cassava* – Unlike most vegetable crops, cassava does not have a definite stage where it is classified as mature since plants will continue to grow as long as there are green leaves. The crop should be harvested nine to 12 months after planting when it has produced the highest percentage of edible storage roots of the desired size (Fig. 24-9). The crop should be harvested before or soon after killing frosts. Critical temperatures for root damage in the field are not known. Care should be taken to avoid skinning roots during the harvesting process (Fig. 24-10). Once harvested, the roots are very perishable. Waxing of the roots is a common practice to improve shelf life.

*Taro & Tannia* – Unlike most vegetable crops, taro (Fig. 24-11) and tannia (Fig. 24-12) do not have a definite stage where they are classified as mature since plants will continue to grow as long as there are green leaves. The crop should be harvested when it has produced the highest percentage of the edible portion (cormels or corms) of the desired size. Tannia and taro should be harvested before killing frosts. Critical temperatures for corm and cormel damage in the field are not known.

*Tropical sweetpotato* — see Chapter 22, Sweetpotato Production in Florida.

### Crop Rotation

*Cassava, Taro & Tannia* – There are usually not as many root-knot nematodes where the preceding crop was a grass or small grain. Most vegetable crops are among the worst crops to precede cassava, from the standpoint of

building up hazardous nematode populations. Sweet corn may be the best rotation crop of the vegetables. Fields that have been planted to okra as a preceding crop should be avoided since rootknot nematode populations tend to build up in okra fields. Do not plant any of these crops in the same field in successive years.

*Tropical sweetpotato* — see Chapter 22, Sweetpotato Production in Florida.

### Flood Tolerance

*Cassava* – plants may tolerate a few days of mild flooding. However, storage root quality is significantly reduced and planting material may be lost if flooding is severe.

*Taro & Tannia* – Taro (malanga isleña) is more flood tolerant than tannia. The variety grown in Miami-Dade does not tolerate continuous flooding. Tannia plants will tolerate a few days of flooding. However, cormel quality is significantly reduced, due to wart like growths that form on the cormels.

### Other Information

*Cassava* – HCN: Fresh roots and leaves are toxic due to the presence of free and bound HCN. The total HCN content varies considerably with variety, environment and plant age. Levels in the peel, peeled root and leaves ranged from 5 to 77, 1 to 40 and 0.3 to 29 mg/100 g (fresh wt.), respectively. Cooking the leaves or roots and changing the cooking water are methods for reducing HCN concentrations. Roots should always be peeled prior to cooling. Toxicity from cassava may develop when considerable quantities are consumed over a period of time. This is particularly true if the prepared cassava has high HCN concentrations and the diet is poorly balanced nutritionally.

*Taro & Tannia* – Nematode-free planting material: Do not take nematodes or other soil borne problems to the field by planting contaminated plants. If plants must be propagated from suspect soil, use cleaned cuttings to avoid carrying potential problems into the field.