

Coffee Senna, *Senna occidentalis* L.¹

David W. Hall, Vernon V. Vandiver, and Jason A. Ferrell²

Classification

Common Name: Coffee Senna

Scientific Name: *Senna occidentalis* (L.) Link

Family: Leguminosae (Fabaceae), Bean Family

Seedling

The cotyledons are smooth, round, about 1 cm long, and usually less than 1 cm wide with 3 distinct veins in the upper surface (Figure 1). The stems have visible hairs just above and below the cotyledons.

Mature Plant

Coffee Senna is a smooth annual that can be 2 m tall (Figure 2). The leaves are compound. The leaflets are in 4-6 pairs and have a sharp leaf apex. These leaflets are 2-9 cm long and 2-3 cm wide with a distinct gland 3-5 mm from the base of the stalk. Flowering occurs in the leaf axils. The sepals are green and 6-9 mm long. The petals are yellow and 1-2 cm long. The 6-7 stamens are of two different lengths. The seed pods are dark brown, 8 to 12 cm long, 7-10 mm wide and curve slightly upward. The



Figure 1. Seedling, Coffee Senna (*Senna occidentalis* L.)

seeds are dull brown, 4-5 mm long and flattened on both ends.

1. This document is an excerpt from Weeds in Florida, SP 37, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: May 1991. Revised: December 2004. Reviewed: February 2009. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>.
2. David W. Hall, former extension botanist, Herbarium, Florida Museum of Natural History; Vernon V. Vandiver, associate professor emeritus, Agronomy Department; Jason A. Ferrell, assistant professor, Agronomy Department; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer, Interim Dean



Figure 2. Mature plant, Coffee Senna (*Senna occidentalis* L.)

History

Senna is an ancient Arabic name for these plants. The Latin word *occidentalis* means western, and refers to the origin.

Habitat

S. occidentalis is widespread in warm areas of the world except for Australasia.

Biology

On two different soil types growth was greater the higher the pH, 4.7-6.3. The seeds are known to be weakly toxic to various stock animals. Animals normally avoid ingesting these seeds. Increased germination is obtained by seed scarification.

Control

Peanuts

There are effectively no preemergence herbicides that provide highly effective coffee senna control. Pursuit, applied preemergence, will deliver approximately 70% control, but all other herbicides will provide 50% control or less.

The most effective herbicide combination is paraquat plus Basagran or Storm applied at-cracking. These combinations will commonly control greater than 90% of all coffee senna present at time of application; no soil-residual activity will be observed. Conversely, paraquat applied alone will not

satisfactorily control coffee senna. Postemergence applications of Cadre or Basagran are effective against coffee senna. Although Cadre and Basagran will generally not provide 100% control, 80 to 90% control is commonly observed.

Cotton

Staple has been shown to provide good control of coffee senna when applied preemergence. All other preemergence herbicides are considered ineffective. Although sicklepod and coffee senna are closely related species, Cotoran does not possess adequate levels of coffee senna activity, as is commonly observed with sicklepod. Postemergence applications of glyphosate and/or Staple provide greater than 90% control of coffee senna. All postemergence directed herbicides, except Cobra and MSMA, provide greater than 90% coffee senna control.