

# University of Florida Potato Variety Trials Spotlight: 'Peter Wilcox'<sup>1</sup>

Rodrick Z. Mwatuwa, Christian T. Christensen, Pam Solano, Kathleen G. Haynes, and Lincoln Zotarelli<sup>2</sup>

## General Comments

'Peter Wilcox' is a fresh market potato variety selected from progeny of a cross between B0810-1 and B0918-5 and tested under the pedigree B1816-5 by K.G. Haynes. It was jointly released by the United States Department of Agriculture, the North Carolina Agricultural Research Service, the Agricultural Experiment Stations of New Jersey, Pennsylvania, Florida, and New York, and the Maine Agricultural and Forest Experiment Station in 2007. 'Peter Wilcox' demonstrates good tuber characteristics and high yields. Yields are slightly lower than the commercial standard 'Red LaSoda' (RLAS). Tuber production and quality results provided in this spotlight are from Florida Potato Variety Trials conducted at the UF/IFAS Hastings Agricultural Extension Center between 2001 and 2016.

## General Characteristics

'Peter Wilcox' has a semi-erect plant habit with intermediate foliage. Tubers have a purple and slightly netted skin with a medium yellow flesh (Figure 1), according to Florida rating codes for potato tuber characteristics (Table 1). The tubers have a fair to good appearance with round to oblong shape and intermediate eye depth with apical distribution (Tables 1 and 3). The tubers have a low to medium specific gravity of 1.065 (Table 2) with a long dormancy (time required for sprout emergence). 'Peter Wilcox' has high yield potential under Florida production conditions with 219

cwt/acre marketable yield and 74% of the tubers produced in tuber size distribution classes A1 and A3 (Table 2).



Figure 1. Typical tuber and internal flesh color of 'Peter Wilcox' potato variety.

Credits: Lincoln Zotarelli, UF/IFAS

## Diseases

'Peter Wilcox' demonstrates no incidence of hollow heart, brown rot, corky ring spot, or internal heat necrosis (Table 3). It is resistant to powdery scab (*Spongospora subterranea*) and susceptible to late blight (*Phytophthora infestans*), early

1. This document is HS1295, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date May 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Rodrick Z. Mwatuwa, research assistant; Christian T. Christensen, postdoctoral research associate, Pam Solano; biological scientist; Kathleen G. Haynes, Genetic Improvement of Fruits and Vegetables Laboratory, USDA-ARS, Beltsville, MD; and Lincoln Zotarelli, assistant professor; Horticultural Sciences Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

blight (*Alternaria solani*), common scab (*Streptomyces scabies*), potato virus Y, and potato virus S. The cultivar is highly susceptible to Verticillium wilt (*Verticillium albo-atrum* and *Verticillium dahliae*). The standard UF/IFAS Extension-recommended disease and weed control program described under “Potato Production” (Chapter 13 of the *Vegetable Production Handbook for Florida* <http://edis.ifas.ufl.edu/cv131>) should be followed.

## Season Length and Growth

‘Peter Wilcox’ is a medium maturing cultivar under Florida growing conditions. Season length was 91 days on average from planting to harvest. This depended on weather conditions during the growing season. The plants should be harvested two to three weeks after vine kill to improve tuber maturation and skin set. Potatoes with proper skin set maintain better skin color, lose less weight in storage, and are more resistant to bruising and soft rot. For more information about vine killing on potatoes, see *Potato Vine Killing or Desiccation* described in Zotarelli et al. (2011). Late in the season, tuber size should be checked regularly in order to harvest tubers with desirable marketable size. Soil moisture should be managed late in the season to avoid high soil moisture conditions that cause enlarged lenticels and delayed skin set.

## Fertilization

UF/IFAS trial plots are normally fertilized with 200 to 230 lb/A of N. The first application of 100 lb/A of N (granular) is typically incorporated in the bed prior to planting, followed by one or two side dress fertilizer applications at emergence and/or at tuber initiation. Phosphorus and potassium applications follow the UF/IFAS guidelines described in Liu et al. (2016) and normally range between 45 to 100 lb/A of P<sub>2</sub>O<sub>5</sub> and 170 to 235 lb/A of K<sub>2</sub>O.

## Planting

A seed piece of 2.5 to 3 oz is recommended for planting. This variety should be planted with 40 inches between rows and 8 inches between plants, at 3 to 4 inches deep. A seed rate of 2,000 to 3,000 lb/acre of seed is expected.

## Other Information

For additional information on cultivation and weed and disease management see the “Potato Production” chapter of the *Vegetable Production Handbook* available at <http://edis.ifas.ufl.edu/cv131>.

## References

- Haynes, K. G., G. Craig Yench, M. E. Clough, M. R. Henninger, X. S. Qu, B. J. Christ, et al. 2015. “Peter Wilcox: a New Purple-Skin, Yellow-Flesh Fresh Market Potato Cultivar with Moderate Resistance to Powdery Scab.” *American Journal of Potato Research* 92(5): 573–581.
- Hutchinson, C. M., J. M. White, D. M., Gergela, P. A. Solano, K. G. Haynes, R. Wenrich, and C. S. Lippi. 2003. “Performance of chip processing potato varieties in north-eastern Florida.” *HortTechnology*, 13(4), 706–711.
- Liu, G., E.H. Simonne, K.T. Morgan, G.J. Hochmuth, M. Ozores-Hampton, and S. Agehara. 2016. “Fertilizer management for vegetable production in Florida.” In: *Vegetable Production Handbook of Florida 2016–17*. J.S. Freeman et al. (eds). Farm Media Journal. p.3–10.
- Sisson, J.A. and G.A. Porter. 2002. “Performance evaluations of potato clones and varieties in the northeastern states-1999.” *Maine Agr. For. Expt. Sta., Misc. Publ.* 751.
- Zotarelli, L., J. P. Dittmar, P. D. Roberts, P. Stansly, H. A. Smith, and S. E. Webb. 2016. *Chapter 13. Potato Production. Vegetable Production Handbook for Florida, 2015–2016 Edition*. HS733. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv131>
- Zotarelli, L., S. Sargent, P. Dittmar, M. Makani. 2011. *Potato vine Killing or Desiccation*. HS181. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/hs181>

Table 1. Florida rating codes for potato vine maturity and tuber characteristics.

Tuber Characteristics <sup>1</sup>							
Rating Code	Vine Maturity	Internal Flesh Color	Skin Color	Skin Texture	Tuber Shape	Eye Depth	Overall Tuber Appearance
1	dead	white	purple	partial russet	round	very deep	very poor
2	+–	cream	red	heavy russet	mostly russet	--	--
3	yellow and dying	light yellow	pink	moderate russet	round to oblong	deep	poor
4	+–	medium yellow	dark brown	light russet	mostly oblong	--	--
5	moderately senesced	dark yellow	brown	netted	oblong	intermediate	fair
6	+–	pink	tan	slightly netted	oblong to long	--	--
7	starting to senesce	red	buff	moderately smooth	mostly long	shallow	good
8	+–	blue	white	smooth	long	--	--
9	green and vigorous	purple	cream	very smooth	cylindrical	very shallow	excellent

<sup>1</sup> Adapted from Hutchinson, C.M. et al. (2003) and Sison, J.A. and G.A. Porter (2002)

Table 2. Summary of production statistics and specific gravity of 'Peter Wilcox', a purple-skinned fresh market potato variety grown at the UF/IFAS Hastings Agricultural Extension Center, Hastings, FL from 2001 to 2016.

Year	Total Yield (cwt/ac)	Marketable yield <sup>1</sup> (cwt/ac)	% of STD	Standard	Size Class (Distribution by class %) <sup>2</sup>						Range %		Specific Gravity
					C	B	A1	A2	A3	A4	A1 to A3	Culls	
2001	313	259	93	'Red Lasoda'	0	14*	77	8	1	0	86	4	1.069
2002	339	296	87	'Red Lasoda'	0	10*	72	17	1	0	90	3	1.064
2003	461	400	95	'Red Lasoda'	5	5	51	33	6	0	90	4	1.062
2004	330	248	76	'Red Lasoda'	12	12	70	6	0	0	75	1	1.076
2005	272	197	66	'Red Lasoda'	1	23	75	1	0	0	76	5	1.072
2006	333	273	68	'Red Lasoda'	2	15	84	0	0	0	84	3	1.070
2007	354	278	80	'Red Lasoda'	2	19	76	3	0	0	79	0	1.067
2008	256	150	78	'Red Lasoda'	4	35	59	2	0	0	61	3	1.071
2009	287	207	61	'Red Lasoda'	2	13	65	17	3	0	85	20	1.062
2010	286	110	47	'Red Lasoda'	10	50	39	0	0	0	39	3	1.060
2011	351	245	109	'Red Lasoda'	5	21	74	0	0	0	74	10	1.062
2012	311	214	124	'Red Lasoda'	3	17	68	12	0	0	80	15	1.060
2013	229	165	154	'Red Lasoda'	4	17	73	6	0	0	79	10	1.055
2014	339	208	93	'Red Lasoda'	4	32	63	1	0	0	64	4	1.062
2015	109	61	43	'Red Lasoda'	5	35	60	0	0	0	60	7	1.060
2016	282	189	208	'Red Lasoda'	5	26	68	1	0	0	69	3	1.069
<b>Average</b>	<b>303</b>	<b>219</b>	<b>93</b>		<b>4</b>	<b>22</b>	<b>67</b>	<b>7</b>	<b>1</b>	<b>0</b>	<b>74</b>	<b>6</b>	<b>1.065</b>

<sup>1</sup> Marketable yield: Sum of size classes A1 to A3.

<sup>2</sup> Size classes: C = 0.5 to 1.5 inches, B = 1.5 to 1 7/8 inches, A1 = 1 7/8 to 2.5 inches, A2 = 2.5 to 3.25 inches, A3 = 3.25 to 4 inches, A4 >4 inches; Size distribution by class: Class (wt)/(Total Yield [wt] – culls [wt])

\* classification = < 1 7/8 inches (C and B included in this classification)

Table 3. Vine maturity, tuber characteristics, and internal tuber defects of 'Peter Wilcox', a purple-skinned fresh market potato variety grown at the UF/IFAS Hastings Agricultural Extension Center, Hastings, FL from 2001 to 2016.

Year	Vine Maturity	Tuber Characteristics <sup>1</sup>						Internal Defects <sup>2</sup>			
		Internal Flesh color	Skin Color	Skin Texture	Tuber Shape	Eye Depth	Overall Appearance	HH	BR	CRS	IHN
2001	3	4	1	6	3	7	6	0	0	0	0
2002	4	4	1	7	3	7	6	0	0	0	0
2003	8	5	1	5	3	6	6	3	0	0	0
2004	6	4	1	6	3	6	6	1	0	0	0
2005	6	5	1	5	4	5	6	0	0	0	0
2006	8	4	2	6	3	7	6	0	0	0	0
2007	6	4	1	7	4	5	6	0	0	0	0
2008	7	5	1	7	3	6	6	0	0	0	0
2009	6	4	1	6	4	4	6	0	0	0	0
2010	7	4	1	6	4	7	6	0	0	0	0
2011	7	4	1	6	4	3	6	0	0	0	0
2012	9	5	1	6	4	4	6	0	1	0	0
2013	7	3	1	5	2	4	5	0	0	1	0
2014	4	5	1	7	3	4	8	0	0	0	0
2015	5	na <sup>3</sup>	na	na	na	na	6	0	0	0	0
2016	6	3	1	7	3	7	7	0	0	3	1
<b>Average</b>	<b>6</b>	<b>4</b>	<b>1</b>	<b>6</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>

<sup>1</sup> See rating system outlined in Florida Rating Code Table (Table 1).

<sup>2</sup> Percent tuber defects. HH = hollow heart, BR = brown rot, CRS = corky ring spot, IHN = internal heat necrosis.

<sup>3</sup> na = data not available