

# University of Florida Potato Variety Trials Spotlight: ‘Yukon Gold’<sup>1</sup>

Rodrick Z. Mwatuwa, Christian T. Christensen, Pam Solano, and Lincoln Zotarelli<sup>2</sup>

## General Comments

‘Yukon Gold’ is a fresh market potato variety selected from the progeny of a cross between W5279-4 and ‘Norgleam’ and tested under the pedigree G6666-4Y. It was released jointly by Agriculture Canada and Ontario Agricultural College, University of Guelph, Canada in 1981 (Johnston and Rowberry 1981). ‘Yukon Gold’ demonstrates high yield and good tuber characteristics compared to the commercial standard ‘La Chipper’. Tuber production and quality results provided in this spotlight are from Florida Potato Variety Trials conducted at the UF/IFAS Hastings Agricultural Extension Center between 1999 and 2014.

## General Characteristics

‘Yukon Gold’ tubers have a buff colored, slightly netted skin with a medium yellow flesh color (Figure 1). According to Florida’s rating codes for potato tuber characteristics (Table 1), the tubers are round to oblong oval shaped and have intermediate pink eyes that are not well distributed, unlike other buff-skinned and yellow-fleshed varieties. ‘Yukon Gold’ has high yield potential with a low to medium specific gravity of 1.071 (Tables 2 and 3) under Florida production conditions. The variety shows reduced dormancy (time required for sprout emergence). Marketable yield was 236 cwt/acre on average, approximately 5% less than ‘La Chipper’. Approximately 86% of the tubers were between the A1 and A3 tuber size classification (Table 2).



Figure 1. Typical tuber and internal flesh color of ‘Yukon Gold’ potato variety.

Credits: Lincoln Zotarelli, UF/IFAS

## Diseases

‘Yukon Gold’ is resistant to mild mosaic, moderately resistant to leafroll virus, and susceptible to virus Y, common scab (*Streptomyces scabies*) (Johnston and Rowberry 1981), PVY, early blight (*Alternaria solani*), dry rot (*Fusarium* spp.), silver scurf (*Helminthosporium solani*), black scurf (*Rhizoctonia solani*), and soft rot (*Pectobacterium* spp.) in storage (University of Nebraska-Lincoln). In all trials, ‘Yukon Gold’ showed a slight susceptibility of 1% to

1. This document is HS1296, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date May 2016. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Rodrick Z. Mwatuwa, research assistant; Christian T. Christensen, postdoctoral research associate; Pam Solano, biological scientist; and Lincoln Zotarelli, assistant professor; Horticultural Sciences Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county’s UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

internal heat necrosis, and less than 0.5% to corky ring spot and hollow heart under Florida conditions (Table 3). The standard UF/IFAS Extension-recommended disease and weed control program described under “Potato Production” (Chapter 13 of the *Vegetable Production Handbook for Florida* <http://edis.ifas.ufl.edu/cv131>) should be followed.

## Season Length and Growth

‘Yukon Gold’ is an early to medium maturing variety under Florida growing conditions. Season length was 92 days on average from planting to harvest. This depended on weather conditions during the growing season. The plants should be harvested two weeks after vine kill to improve tuber maturation and skin set. Potatoes with proper skin set maintain better skin color, lose less weight in storage, and are more resistant to bruising and soft rot. For more information about vine killing on potatoes, see *Potato Vine Killing or Desiccation* (Zotarelli et al. 2011). Late in the season, tuber size should be checked regularly to harvest tubers with desirable marketable size. Soil moisture should be managed late in the season to avoid high soil moisture conditions that cause enlarged lenticels and delayed skin set.

## Fertilization

UF/IFAS trial plots are normally fertilized with 200 to 230 lb/A of N. The first application of 100 lb/A of N (granular) is typically incorporated in the bed prior to planting, followed by one or two side dress fertilizer applications at emergence and/or at tuber initiation. Phosphorus and potassium applications follow the UF/IFAS guidelines described in Liu et al. (2016) and normally range between 45 to 100 lb/A of P<sub>2</sub>O<sub>5</sub> and 170 to 235 lb/A of K<sub>2</sub>O.

## Planting

A seed piece of 2.5 to 3 oz is recommended for planting. Crop should be planted with 40 inches between rows and 8 inches between plants, at 3 to 4 inches deep. A seed rate of 2,000 to 3,000 lb/acre of seed is expected.

## Other Information

For additional information on cultivation and weed and disease management see the Potato Production chapter of the *Vegetable Production Handbook* available at <http://edis.ifas.ufl.edu/cv131>.

## References

- Hutchinson, C. M., J. M. White, D. M., Gergela, P. A. Solano, K. G. Haynes, R. Wenrich, and C. S. Lippi. 2003. “Performance of chip processing potato varieties in north-eastern Florida.” *HortTechnology*, 13(4), 706–711.
- Johnston, G. R. and R. G. Rowberry. 1981. “Yukon Gold’: a new yellow-fleshed, medium-early, high quality table and french-fry cultivar.” *American Potato Journal*, 58(5), 241–244.
- Liu, G., E.H. Simonne, K.T. Morgan, G.J. Hochmuth, M. Ozores-Hampton, and S. Agehara. 2016. “Fertilizer management for vegetable production in Florida.” In: *Vegetable Production Handbook of Florida 2016—17*. J.S. Freeman et al. (eds). Farm Media Journal. p. 3–10.
- Nebraska Institute of Agriculture and Natural Resources. 2017. *Yukon gold characteristics*. Crop Watch [http://crop-watch.unl.edu/potato/yukongold\\_characteristics](http://crop-watch.unl.edu/potato/yukongold_characteristics). Accessed on 5 December 2016.
- Sisson, J.A. and G.A. Porter. 2002. “Performance evaluations of potato clones and varieties in the northeastern states-1999.” Maine Agr. For. Expt. Sta., Misc. Publ. 751.
- The Potato Association of America. 2017. ‘Yukon Gold’ (*Solanum tuberosum*). <http://potatoassociation.org/industry/varieties/white-varieties/yukon-gold-solanum-tuberosum>. Accessed on 5 December 2016.
- Zotarelli, L., S. Sargent, P. Dittmar, M. Makani. 2011. *Potato vine Killing or Desiccation*. HS181. Gainesville: University of Florida Institute of Food and Agricultural Sciences. Available at: <http://edis.ifas.ufl.edu/hs181>
- Zotarelli, L., J. P. Dittmar, P. D. Roberts, P. Stansly, H. A. Smith, and S. E. Webb. 2016. *Chapter 13. Potato Production. Vegetable Production Handbook for Florida, 2015–2016 Edition*. HS733. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cv131>

Table 1. Florida rating codes for potato vine maturity and tuber characteristics.

Tuber Characteristics <sup>1</sup>							
Rating Code	Vine Maturity	Internal Flesh Color	Skin Color	Skin Texture	Tuber Shape	Eye Depth	Overall Tuber Appearance
1	dead	white	purple	partial russet	round	very deep	very poor
2	+-	cream	red	heavy russet	mostly round	--	--
3	yellow and dying	light yellow	pink	moderate russet	round to oblong	deep	poor
4	+-	medium yellow	dark brown	light russet	mostly oblong	--	--
5	moderately senesced	dark yellow	brown	netted	oblong	intermediate	fair
6	+-	pink	tan	slightly netted	oblong to long	--	--
7	starting to senesce	red	buff	moderately smooth	mostly long	shallow	good
8	+-	blue	white	smooth	long	--	--
9	green and vigorous	purple	cream	very smooth	cylindrical	very shallow	excellent

<sup>1</sup> Adapted from Hutchinson, C.M. et al. (2003) and Sison, J.A. and G.A. Porter (2002)

Table 2. Summary of production statistics and specific gravity of 'Yukon Gold' potato variety grown at the UF/IFAS Hastings Agricultural Extension Center, Hastings, FL from 1999 to 2014.

Year	Total Yield (cwt/ac)	Marketable yield (cwt/ac)	% of Standard	Standard	Size Class (Distribution by class %)						Range %		Specific Gravity
					C	B	A1	A2	A3	A4	A1 to A3	Culls	
1999	422	392	114%	LAC	*	7*	73	20	0	0	93	5	1.070
2001	325	277	91%	LAC	*	2*	37	42	19	1	97	12	1.078
2002	298	258	88%	LAC	*	4*	40	49	8	0	96	9	1.074
2003	391	303	101%	LAC	*	14*	42	31	13	0	86	11	1.076
2004	311	217	95%	LAC	16	11	57	16	2	0	74	6	1.076
2005	226	180	72%	LAC	2	15	64	18	1	0	83	4	1.073
2006	342	275	89%	LAC	1	8	71	20	0	0	91	11	1.069
2007	325	286	100%	LAC	1	7	65	20	6	0	92	4	1.073
2008	280	213	111%	LAC	2	16	67	13	3	0	82	7	1.082
2009	361	268	105%	LAC	1	5	48	23	23	0	93	21	1.070
2010	326	250	131%	LAC	2	13	72	10	3	0	85	10	1.067
2011	348	248	88%	LAC	3	13	56	15	13	0	83	14	1.068
2012	275	179	74%	LAC	2	9	48	28	12	0	89	27	1.066
2013	129	45	49%	LAC	7	17	68	8	0	0	76	43	1.061
2014	271	145	100%	LAC	4	26	61	7	2	0	70	24	1.063
<b>Average</b>	<b>309</b>	<b>236</b>	<b>94%</b>		<b>4</b>	<b>13</b>	<b>58</b>	<b>21</b>	<b>7</b>	<b>0</b>	<b>86</b>	<b>14</b>	<b>1.072</b>

<sup>1</sup> Marketable yield: Sum of size classes A1 to A3.

<sup>2</sup> Size classes: C = 0.5 to 1.5 inches, B = 1.5 to 1 7/8 inches, A1 = 1 7/8 to 2.5 inches, A2 = 2.5 to 3.25 inches, A3 = 3.25 to 4 inches, A4 >4 inches; Size distribution by class: Class (wt)/(Total Yield [wt] – culls [wt])

<sup>3</sup> n.a. = not available

\* classification = <1 7/8 inches (C and B included in this classification)

Table 3. Yield, vine maturity, tuber characteristics, and internal tuber defects of 'Yukon Gold' potato variety grown at the UF/IFAS Hastings Agricultural Extension Center, Hastings, FL from 1999 to 2014.

Year	Vine Maturity (vine kill)	Tuber Characteristics <sup>1</sup>						Internal Defects <sup>2</sup>			
		Internal Flesh color	Skin Color	Skin Texture	Tuber Shape	Eye Depth	Overall Appearance	HH	BR	CRS	IHN
1999	*	*	7	5	3	4	4	2	0	0	1
2001	1	3	8	7	3	6	6	0	0	5	0
2002	2	3	7	7	3	7	6	0	1	0	5
2003	5	4	7	7	3	6	6	0	0	0	1
2004	5	4	7	7	3	6	6	3	0	0	0
2005	4	4	7	6	3	5	5	0	0	0	0
2006	7	2	4	4	2	3	6	1	0	0	0
2007	5	4	7	6	4	6	6	0	0	0	0
2008	7	3	7	6	2	8	5	0	0	0	3
2009	4	4	7	6	3	5	*	0	0	0	0
2010	7	4	7	7	3	5	6	0	0	0	0
2011	7	5	7	6	4	6	5	0	0	0	3
2012	9	4	7	6	3	4	5	0	0	0	0
2013	5	5	7	7	3	5	5	0	0	0	0
2014	3	5	7	6	3	6	5	0	0	0	0
<b>Average</b>	<b>5</b>	<b>4</b>	<b>7</b>	<b>6</b>	<b>3</b>	<b>5</b>	<b>5</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>

<sup>1</sup> See rating system outlined in Florida Rating Code Table (Table 1).

<sup>2</sup> Percent tuber defects. HH = hollow heart, BR = brown rot, CRS = corky ring spot, IHN = internal heat necrosis.

\* not available