

Cottony Cushion Scale, *Icerya purchasi* Maskell¹

A. B. Hamon and T. R. Fasulo²

Introduction

The cottony cushion scale (Figure 1) was described by Maskell (1878) from specimens sent to him by Dr. Purchas from Auckland, New Zealand. The host was kangaroo acacia and the insect was named for Dr. Purchas. At that time only one other species was known in the genus *Icerya* (Maskell 1878). This scale is apparently native to Australia and made its way to California on acacia plants around 1868 or 1869, and in about ten years was causing damage to citrus groves in southern California (Ebeling 1959). The measures taken to control the cottony cushion scale in California were to change the methods of insect control and research first in California and later in the whole United States. These new control methods led to the use of biological control and legislative quarantine (Ebeling 1959).

The following account of the introduction of this scale insect into Florida is largely taken from Gossard (1901). The ladybug, *Rodolia cardinalis* (Mulsant), was introduced into California in 1888 for the biological control of this scale (DeBach 1973). In 1893, the owners of a nursery in Keene, Florida (Pinellas County) sent an inquiry to someone in



Figure 1. Cottony cushion scales, *Icerya purchasi*, on twig. Credits: James Castner, University of Florida

California about the possibility of the ladybug being used to control other scale insects in Florida. Apparently interpreting this as a request for the ladybug the party sent a shipment of ladybugs and included some cottony cushion scales as food for the ladybugs. The nursery owners who either didn't see the scales or assumed they would be of no consequence, left the packing container near a citrus tree which eventually became infested. Florida now had the dreaded cottony cushion scale with which to contend. The originally infested Florida trees were destroyed, but infested trees appeared again in late 1898. However, this was presumably from a new

1. This document is EENY-034, one of a series of the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date June 1998. Reviewed March 2011. This document is also available as a Featured Creature at <http://creatures.ifas.ufl.edu/>. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. A. B. Hamon and T. R. Fasulo, Entomology and Nematology Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean

introduction from a tramp peddler of nursery stock in about 1895 (Gossard 1901).

Distribution

The cottony cushion scale is now widespread throughout the world wherever citrus is grown (Ebeling 1959). In Florida, this scale has been reported from most counties.

Description

The cottony cushion scale can be distinguished easily from other scale insects in Florida. It is the only species of *Icerya* present in Florida. The mature females (actually hemaphrodites) have bright orange-red, yellow, or brown bodies (Ebeling 1959). The body is partially or entirely covered with yellowish or white wax (Figure 2). The most conspicuous feature is the large fluted egg sac, which will frequently be 2 to 2.5 times longer than the body. The egg sac contains about 1000 red eggs (Gossard 1901). Depending on the temperature eggs hatch in a few days to two months. The newly hatched nymphs are bright red with dark antennae and thin brown legs. The antennae are six-segmented. This is the primary dispersal stage, and can be wind blown to new locations, crawl to nearby plants, or possibly hitchhike on other animals. After three molts the adult begins to deposit eggs and secrete the conspicuous egg sac. As the egg sac is formed the scale abdomen becomes more tilted until the scale appears to be standing on its head.



Figure 2. Adult female cottony cushion scale, *Icerya purchasi* Maskell. Credits: James Castner, University of Florida

Males are rare. They are winged with dark red body and dark colored antennae. Dark whorls of setae extend from each antennal segment, except the first (Ebeling 1959). It is interesting that the female is always a hemaphrodite with both testes and ovaries. If self-fertilization occurs only hemaphrodites are produced; however, when a hermaphrodite mates with a male more males and hemaphrodites are produced (Ebeling 1959).



Figure 3. Adult vedalia beetles, *Rodolia cardinalis* (Mulsant), feeding on cottony cushion scale, *Icerya purchasi* Maskell. Credits: James Castner, University of Florida

Host Plants

Cottony cushion scale is most frequently collected on *Citrus* and *Pittosporum* in Florida. However, numerous records on other host plants are in the Division of Plant Industry insect files.

Economic Importance

The cottony cushion scale can severely damage trees, resets, and nursery stock. Decreased tree vitality, fruit drop, and defoliation result from the feeding of this scale. Most damage occurs from the feeding of the early immature stages of the scale on the leaves, where they settle in rows along the midrib and veins, and on the smaller twigs. The older nymphs continue to feed, but migrate to the larger twigs, and finally, as adults, they settle on the larger branches and trunk. This scale is seldom found on the fruit. Added damage can result from the accumulation of sooty mold due to the honeydew excreted by the scale.

Management

Citrus Pest Management Guide for Scale Insects

Selected References

Debach, P. 1973. Biological control of insect pests and weeds, Chapman and Hall, London. 844 p.

Ebeling, W. 1959. Subtropical fruit pests. Univ. of California Press, Los Angeles. 436 p.

Fasulo, T.R., and R.F. Brooks. (January 2001). Scale Pests of Florida Citrus. UF/IFAS ENY814. <http://edis.ifas.ufl.edu/CH059> (13 September 2001).

Gossard, H.A. 1901. The cottony cushion scale. Florida Agr. Exp. Sta. Bull. 56:309-356.

Maskell, W.M. 1878. On some Coccidae in New Zealand, Trans. Proc. New Zealand Inst. 11:220-223.

Morrison, H. 1928. A classification of the higher groups and genera of the coccid family Margarodidae. USDA Tech. Bull. 52:203-211.