

Bed Bug, *Cimex lectularius*, Linnaeus (Insecta: Hemiptera: Cimicidae)¹

Shawn E. Brooks²

Introduction

Sometimes referred to as "red coats," "chinchies," or "mahogany flats" (USDA 1976), bed bugs, *Cimex lectularius* Linnaeus, are blood feeding parasites of humans, chickens, bats and occasionally domesticated animals (Usinger 1966). Bed bugs are suspected carriers of leprosy, oriental sore, Q-fever, and brucellosis (Krueger 2000) but have never been implicated in the spread of disease to humans (Dolling 1991). After the development and use of modern insecticides, such as DDT, bed bug infestations have virtually disappeared. However, since 1995, pest management professionals have noticed an increase in bed bug related complaints (Krueger 2000).

Distribution

Human dwellings, birds nests, and bat caves make the most suitable habitats for bed bugs since they offer warmth, areas to hide, and most importantly hosts on which to feed (Dolling 1991). Bed bugs are not evenly distributed throughout the environment but are instead concentrated in harborages (Usinger 1966). Within human dwellings,



Figure 1. Adult bed bug, *Cimex lectularius* Linnaeus, feeding. Credits: Joseph Smith, University of Florida

harborages include cracks and crevices in walls, furniture, behind wallpaper and wood paneling, or under carpeting (Krueger 2000). Bed bugs are usually only active during night but will feed during the day when hungry (Usinger 1966). Bed bugs can be transported on clothing, in traveler's luggage, or in bedding and furniture (USDA 1976) but lack appendages to enable them to cling to hair, fur, or feathers, so are rarely found on hosts (Dolling 1991).

1. This document is EENY-140 (IN297), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: June 2000. Revised: December 2005. Reviewed: March 2008. This document is also available on Featured Creatures Website at <http://creatures.ifas.ufl.edu>. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>.

2. Shawn E. Brooks, Entomology and Nematology Department, University of Florida, Institute of Food and Agricultural Sciences, Gainesville, FL 32611.

Description

The adult bed bug is a broadly flattened, ovoid, insect with greatly reduced wings (Schuh and Slater 1995). The reduced fore wings, or hemelytra, are broader than they are long, with a somewhat rectangular appearance. The sides of the pronotum are covered with short, stiff hairs (Furman and Catts 1970). Before feeding, bed bugs are usually brown in color and range from 6 to 9.5 mm in length. After feeding, the body is often swollen and red in color (USDA 1976).



Figure 2. Adult bed bug, *Cimex lectularius* Linnaeus. Credits: David Almquist, University of Florida



Figure 3. Lateral view of an adult bed bug, *Cimex lectularius* Linnaeus. Credits: Joseph Smith, University of Florida

The two bed bugs most important to man are the common bed bug, *Cimex lectularius*, and the tropical bed bug, *Cimex hemipterus*. These two species of bed bugs can easily be distinguished by looking at the prothorax, the first segment of the thorax. The prothorax of the common bed bug is more expanded laterally and the extreme margins are more flattened than that of the tropical bed bug (Smith 1973).



Figure 4. Nymph of the bed bug, *Cimex lectularius* Linnaeus. Credits: Joseph Smith, University of Florida

Life Cycle

Because of their confined living spaces, copulation among male and female bed bugs is difficult. The female possesses a secondary copulatory aperture, Ribaga's organ or paragenital sinus, on the fourth abdominal sternum where spermatozoa from the male are injected. The spermatozoa then migrate to the ovaries by passing through the haemocoel, or body cavity (Dolling 1991). The female bed bug lays approximately 200 eggs during her life span at a rate of one to 12 eggs per day (Krueger 2000). The eggs are laid on rough surfaces and coated with a transparent cement to adhere them to the substrate (Usinger 1966). Within six to 17 days bed bug nymphs, almost devoid of color, emerge from the eggs. After five molts, which takes approximately ten weeks, the nymphs reach maturity (USDA 1976).

Survey and Management

Bed bugs are most active at night, they are extremely shy and wary so their infestations are not easily located (Snetsinger 1997). However, when bed bugs are numerous, a foul odor from oily secretions can easily be detected (USDA 1976). Other recognizable signs of a bed bug infestation include excrement left around points of entry and exit to their hiding places (Dolling 1991) and reddish brown spots on mattresses and furniture (Frishman 2000). Good sanitation is the first step to controlling the spread of bed bugs. However, upscale hotels and private homes

have recently noted infestations, suggesting that good sanitation is not enough to stop a bed bug infestation (Krueger 2000).

If bed bugs are located in bedding material or mattresses, control should focus on mechanical methods of control, such as vacuuming, caulking and removing or sealing loose wallpaper, to minimize the use of pesticides (Frishman 2000). The effectiveness of using steam cleaners or hot water to clean mattresses is questionable. Heat is readily absorbed by the mattress and does no harm to the bed bug in fact the moisture may produce favorable conditions for house dust mites. Pillows should be removed and dry-cleaned or replaced (Snetsinger 1997). For severe infestations, however, pesticides may be used. Care should be taken not to soak mattresses and upholstery with pesticides. Allow bedding and furniture to dry thoroughly before using.

For more information, see the Insect Management Guide for Bed Bugs (<http://edis.ifas.ufl.edu/IG083>).

Selected References

Dolling WR. 1991. *The Hemiptera*. Oxford University Press, New York, New York.

Fasulo TR. 2002. Bloodsucking Insects. UF/IFAS SW 156.

Fasulo TR, Kern W, Koehler PG, Short DE. 2005. *Pests In and Around the Home*. Version 2.0. UF/IFAS CD-ROM. SW 126.

Frishman A. 2000. Bed Bug basics and control measures. *Pest Control* 68:24.

Furman DP, Catts E. 1970. *Manual of Medical Entomology*, 3rd ed. National Press Books, Palo Alto, California.

Ghauri MSK. 1973. Hemiptera (bugs), pp. 373-393. *In* K.G.V. Smith [ed], *Insects and other Arthropods of Medical Importance*. British Museum, London, England. [USDA] U.S. Department of Agriculture. 1976. *How to control bed bugs*. USDA. Washington D.C.

Koehler PG, Hertz J. 2005. Bed bugs and blood-sucking conenose. EDIS. "<http://edis.ifas.ufl.edu/IG083> (9 December 2005).

Krueger L. 2000. Don't get bitten by the resurgence of bed bugs. *Pest Control* 68:58-64.

Potter, MF 2004. *Bed Bugs*. University of Kentucky Entomology FactSheets. <http://www.uky.edu/Agriculture/Entomology/entfacts/struct/ef636.htm> (9 December 2005).

Snetsinger R. 1997. Bed bugs & other bugs, pp. 393-425. *In* A. Mallis and S.A. Hedges [eds.], *Handbook of Pest Control*, 8th ed. Franzak & Foster Co., Cleveland, Ohio.

Schuh R, Slater JA 1995. *True Bugs of the World (Hemiptera : Heteroptera) Classification and Natural History*. Cornell University Press, Ithica, New York.

Usinger RL. 1966. *Monograph of Cimicidae (Hemiptera - Heteroptera)*. Entomological Society of America, College Park, Maryland.