

# Pantropical Huntsman Spider, *Heteropoda venatoria* (Linnaeus) (Arachnida: Araneae: Sparassidae)<sup>1</sup>

G. B. Edwards, Jr.<sup>2</sup>

## Introduction

The pantropical huntsman spider, *Heteropoda venatoria* (L.), sometimes called the giant crab spider or the banana spider (due to its occasional appearance in marketed bananas), is a cosmopolitan species introduced into and now occurring in the United States, in subtropical areas of Florida, Texas, and California, and in coastal areas of Georgia and South Carolina. It is presumed to have been introduced from Asia, where many of its closest relatives live (Gertsch 1948). It is sometimes mistaken for a large **brown recluse**, *Loxosceles reclusa* Gertsch and Mulaik, a poisonous venomous spider in the family Loxoscelidae-Sicariidae, but it is neither related nor is it dangerous. Some authors placed this spider in the family Heteropodidae, due to the uncertainty of the name Sparassidae (Platnick and Levi 1973), but the family name Sparassidae was stabilized by Jäger (1999).

## Abbreviated Synonymy

*Aranea venatoria* Linnaeus (1767)

*Aranea regia* Fabricius (1793)

*Heteropoda venatoria*, Latreille (1804)

*Thomisus leucosia* Walckenaer (1805)

*Micrommata setulosa* Perty (1833)



Figure 1. Adult male huntsman spider, *Heteropoda venatoria* (Linnaeus).

Credits: Marie Knight

## Description

*Heteropoda venatoria* is a large brown spider with a flattened body structure and very little dorsal pattern. Adult specimens have a body length of 2.2 to 2.8 cm (about 1 inch), and have a leg span of 7 to 12 cm (3 to 5 inches).

1. This document is EENY-160, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date October 2000. Revised June 2009. Reviewed April 2018. Visit the EDIS website at <http://edis.ifas.ufl.edu>. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. G. B. Edwards, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Florida State Collectoin of Arthropods; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Adult females have a larger body size, especially the abdomen, than males. Adult males have longer legs than females, and the long male palpi have the terminal segment enlarged and the ventral sclerites exposed, as in most true spiders. Both sexes have a yellow to cream clypeus and a wide marginal band encircling the rest of the carapace, tan in females and cream in males. In addition, males have a dark, longitudinal stripe on the abdomen and a light-bordered pale area behind the eyes. The legs of both sexes have distinct black spots from each of which arises an erectile macroseta. Otherwise, the spider is not conspicuously hairy.



Figure 2. Female brown recluse spider, *Loxosceles reclusa* Gertsch and Mulaik, dorsal view for comparison with dorsal view of male huntsman spider, *Heteropoda venatoria* (Linnaeus).  
Credits: Jeffrey Lotz, DPI



Figure 3. Frontal view of adult male huntsman spider, *Heteropoda venatoria* (Linnaeus), showing the flattened body structure.  
Credits: Marie Knight



Figure 4. Closeup of adult male huntsman spider, *Heteropoda venatoria* (Linnaeus). See the dark, longitudinal stripe on the abdomen and a light-bordered pale area behind the eyes.  
Credits: Marie Knight

## Habits and Habitat

Females of *H. venatoria* make flattened, disc-like eggsacs about 1.5 cm in diameter which contain over 200 eggs. The eggsac is carried under the body, its size and shape probably causing the female to remain relatively immotile. All stages of development of juveniles and adults appear to occur simultaneously throughout the year.

This and similar species are highly valued in tropical countries because they capture and feed on cockroaches and other domestic insect pests. As with other vagrant spiders, huntsman spiders do not use webs to capture prey. Their great speed and strong chelicerae (jaws) are used to capture the insects on which they feed. Venom is also injected into the prey from glands extending from the chelicerae into the cephalothorax.

The flattened body enables this large spider to fit into surprisingly small cracks and crevices. This ability, along with its adaptability to human habitations, helps explain its frequent occurrence in houses, barns, sheds, under boards on the ground, and in other sheltered areas. Being cold-sensitive, these spiders cannot exist outdoors in areas with freezing winter temperatures; occasionally they occur in greenhouses and other heated buildings in temperate climates. On the other hand, in southern Florida where frost is rare, these spiders have become “wild” (i.e., no longer requiring man-made structures). In the Homestead area, *H. venatoria* is now established in the “wild” and is common in avocado groves (Whitcomb unpublished).

They can be easily collected at night by using a headlight (Wallace 1937) as their eyes reflect light, appearing as blue spots on the trunks of trees and on the ground, much like wolf spiders.

## Survey and Detection

In Florida, *H. venatoria* may be distinguished from other large, cursorial spiders by its flattened brown body and the black spots around the macrosetae on the legs. In warm weather, it may be found in and about human habitations; in cold weather it will be found indoors, under furniture or cabinets, behind wall hangings, and in closets and garages. It is not a dangerous spider, but a locally painful bite, sometimes with noticeable swelling, can be delivered to any human who carelessly handles a huntsman spider or accidentally traps it, for example, in a shoe.

## Selected References

Breene RG. (2003). Common Names of Arachnids. *The American Arachnological Society Information on Arachnids*. [http://www.americanarachnology.org/assets/pdfs/arachnid\\_common\\_names2003.pdf](http://www.americanarachnology.org/assets/pdfs/arachnid_common_names2003.pdf) (April 2018).

Florida Nature. (2004). *Heteropoda venatoria*, Huntsman Spider. *FloridaNature.org*. [http://flnature.org/species.asp?species=Heteropoda\\_venatoria](http://flnature.org/species.asp?species=Heteropoda_venatoria) (22 June 2009).

Gertsch WJ. 1948. *American Spiders*. New York, NY: D. Van Nostrand Co., Inc. 284 pp.

Jäger, P. 1999. "Sparassidae – the valid scientific name for the huntsman spiders (Arachnida: Araneae)." *Arachnologische Mitteilungen* 17: 1–10.

Platnick NI, Levi HW. 1974. "On names of spiders." *British Arachnology Society Bulletin* 2: 166–167.

Wallace HK. 1937. "The use of a headlight in collecting nocturnal spiders." *Entomological News* 48: 160–161.