

## South American Cucurbit Fruit Fly, *Anastrepha grandis* (Macquart) (Insecta: Diptera: Tephritidae)<sup>1</sup>

---

H. V. Weems, Jr.<sup>2</sup>

### Introduction

The South American cucurbit fruit fly, *Anastrepha grandis* (Macquart), which exists in several South American countries, attacks watermelon and other fruits of the family Cucurbitaceae. Once a pest of minor to moderate importance generally, in recent years it has become a rather important pest. The pest status differs in each country and has sometimes changed within a country. This species is potentially of economic importance in Florida and southern Texas should it ever be introduced there. It has been intercepted in the United States in pumpkin from Argentina and Brazil, and in banana debris from Panama.

### Synonymy

*Acrotoxa grandis* (Macquart)

*Anastrepha schineri* Hendel

*Anastrepha latifasciata* Hering

*Tephritis grandis* Macquart

*Trypeta grandis* (Macquart)

### Distribution

Argentina, Bolivia, Brazil (except north), Colombia, Ecuador, Panama (NAPPO 2009), Paraguay, Peru and Venezuela.

### Identification

Rather large, yellow-brown, with yellow and dark-brown markings. Mesonotum 3.3 to 4.0 mm long, yellow-brown, with humerus, median stripe widening to include acrostichal bristles but not reaching scutellum, lateral stripes from just before transverse suture to side of scutellum, stripe below notopleuron, metapleuron, and scutellum except extreme base yellow; a sublateral stripe from level of humeral bristle to scutellum, broken at transverse suture, a band along scutoscutellar suture, intensified medially, and a spot on pteropleuron dark brown; metanotum blackened laterally. Macrochaetae dark brown; pile yellowish brown. No sternopleural bristle. Wing 9.0 to 10.5 mm long, the bands yellow brown, rather diffuse; costal and S bands broadly connected, and no distinct hyaline spot anterior to

---

1. This document is EENY-205 (IN362) (originally published as DPI Entomology Circular 334), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: March 2001. Revised November 2009. This document is also available on Featured Creatures Web site at <http://entomology.ifas.ufl.edu/creatures>.

2. H. V. Weems, Jr. (retired), Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL.

vein  $R_{4+5}$ ; distal arm of V band absent, the proximal arm not joining S band.



**Figure 1.** Adult female South American cucurbit fruit fly, *Anastrepha grandis* (Macquart). Credits: Division of Plant Industry




**Figure 2.** Adult male South American cucurbit fruit fly, *Anastrepha grandis* (Macquart). Credits: Division of Plant Industry


Female terminalia: Ovipositor sheath 5.8 to 6.2 mm long, tapering posteriorly to apical third, which is distinctly depressed and broadened; in profile the sheath is distinctly concave dorsally on median half and concave ventrally on apical third. Rasper well developed, of slender, curved hooks in five or six rows. Ovipositor slightly longer than length of ovipositor sheath, being somewhat curved dorsoventrally to permit fitting into sheath; tip long and slender, without serrations; extreme base slightly widened (Stone 1942).

Steyskal (1977) distinguished adult *A. grandis* from the other four species of the *Grandis* group as follows:

1. Vein  $R_3$  somewhat undulant; metanotum yellow . . . . *A. bezzii* Lima and *A. balloui* Stone



**Figure 3.** Rasters at tip of ovipositor sheath. Credits: Division of Plant Industry



**Figure 4.** Ovipositor Credits: Division of Plant Industry

1'. Vein  $R_3$ , not undulant; metanotum marked with black . . . . 2

2. Mesonotum not striped with black . . . . *A. atrigona* Hendel

2'. Mesoscutum with black stripes . . . . 3

3. Mesoscutum and pleura largely black, wing with pattern darkened . . . . *A. shannoni* Stone

3'. Mesoscutum and pleura with little black, wing pattern very little darkened . . . . *A. grandis* (Macquart)

## Hosts

- *Citrullus lanatus* (Thunberg) Matsumura & Nakai - watermelon
- *Cucumis sativus* Linnaeus - cucumber
- *Cucurbita maxima* Duchesne - autumn and winter squash, pumpkin
- *Cucurbita pepo* Linnaeus - summer and autumn pumpkin and squash, gourd, marrow
- *Cucurbita moschata* (Duchesne ex Lam.) Duchesne ex Poiret - "amyama," "pumpkin"
- *Lagenaria siceraria* (Molina) Standl. - "camasa"

- *Cucumis melo* L. - "honeydew melon"

It has also been known to attack common guava, *Psidium guajava* (Norrbom 2004) The record of rearing from oranges (Greene 1934) is erroneous. Immature fruits of cucurbits apparently are preferred, but mature or nearly mature fruits of some varieties are attacked occasionally.

## Survey and Detection

Adults are easily recognized by their very large size (wing length 10 mm), long aculeus (a sharp spine jutting from beneath the margin of the eighth sternite) and rather diffuse wing markings which are not of the typical *Anastrepha* type.

Larvae can be collected from infested fruit and are very difficult to identify to species except when reared to adults. For best larval preservation, kill in boiling water, place in 50% isopropyl alcohol for two days, then in 75% isopropyl alcohol. Adults may be collected on stickyboard and in baited traps.

## Selected References

- Carroll LE, Norrbom AL, Dallwitz MJ, Thompson FC. (December 2006). *Anastrepha grandis* (Macquart). *Pest Fruit Flies of the World – Larvae*.  
[http://delta-intkey.com/ffl/www/ana\\_gran.htm](http://delta-intkey.com/ffl/www/ana_gran.htm) (30 October 2009).
- Enkerlin D, Laura-Garcia R, Fidel-Lopez M. 1989. pp. 83-90. In Robinson AS, Hooper G (editors), *Fruit Flies, Their Biology, Natural Enemies and Control*. 3A: I-xii, 1-372.
- Foote RH. 1967. Family Tephritidae. In Vanzolini, M. (editor), *A Catalogue of the Diptera of the Americas South of the United States*. 57: 1-91. Dept. Zool., Sec. Agric., São Paulo, Brazil.
- Greene CT. 1934. A revision of the genus *Anastrepha* based on a study of the wings and on the length of the ovipositor sheath (Diptera: Trypetidae). *Proceedings of the Entomological Society of Washington* 36: 127-179.
- Macquart J. 1846. Diptères exotiques nouveaux ou peu connus. (1845) (1844): 133-364, illus. *Mém. Soc. Roy. Sci., Agric., Arts Lille*. (published separately as *Supplément I*, p. 5-238.
- NAAPO. (July 2009). Outbreak of *Anastrepha grandis* (South American cucurbit fruit fly) in Panama. *Phytosanitary Alert System*.  
<http://www.pestalert.org/viewNewsAlert.cfm?naid=76> (30 October 2009).
- Norrbom AL. (31 October 2001). *Anastrepha grandis* (Macquart). *The Diptera site*.  
<http://www.sel.barc.usda.gov/diptera/tephriti/Anastrep/grandis.htm> (14 January 2002).
- Norrbom AL. (2004). Host plant database for *Anastrepha* and *Toxotrypana* (Diptera: Tephritidae: Toxotrypanini). *Diptera Data Dissemination Disk 2* (CD-ROM). North American Dipterist's Society. Washington, D.C.
- Steyskal GC. 1977. Pictorial key to species of the genus *Anastrepha* (Diptera: Tephritidae). (Special publication) *Entomological Society of Washington* 35 pp.
- Stone A. 1942. *The Fruitflies of the Genus Anastrepha*. U.S. Department of Agriculture Miscellaneous Publication No. 439, Washington, DC. 112 pp.
- White IM, Elson-Harris MM. 1994. *Fruit Flies of Economic Significance: Their Identification and Bionomics*. CAB International. Oxon, UK. 601 pp.