

Red admiral *Vanessa atalanta rubria* (Fruhstorfer) (Insecta: Lepidoptera: Nymphalidae: Nymphalinae) ¹

Donald W. Hall and Jerry F. Butler²

Introduction

The red admiral is a distinctive, medium sized, brightly colored black and orange butterfly that is common throughout much of the United States.

Distribution

The red admiral occurs in Europe, northern Africa, Eurasia, and in North America from northern Canada to Central America. It is also found in Hawaii, New Zealand, and some of the Caribbean Islands (Opler et al. 1984, Scott 1986). Adults are found throughout Florida, but it is not known whether or not there are breeding populations in the Florida Keys (Minno and Emmel 1993).

Description

Adult: The wing spread is 1.75 to 2.50 inches (Daniels 2003). The upper side of the wings is black. The front wings have a broad, orange median band with white spots nearer the apex. The hind wings have a marginal orange band. Both fore and hind wings have a series of curved white dashes along the margins. The underside of the front wings are dark

with white spots apically and a pinkish to orange median band. The undersides of the hind wings are marbled.



Figure 1. Adult red admiral, *Vanessa atalanta rubria* (Fruhstorfer), dorsal view. Credits: D.W. Hall, University of Florida

Egg: The eggs are light green and are surrounded with a series of white vertical ridges.

Larva: The larvae are up to 1.4 inches in length (Minno et al. 2005). They are highly variable in color. The head is black with short spines and hairs. The body varies from pale brown with a yellowish stripe on the sides to black with small white spots and a broken white stripe on the sides. Each segment has

-
1. This document is EENY-446 (IN810), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: April 2009. This document is also available on Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
 2. Donald W. Hall, professor and Jerry F. Butler, professor, Entomology and Nematology Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean



Figure 2. Adult red admiral, *Vanessa atalanta rubria* (Fruhstorfer), ventral view. Credits: J.F. Butler, University of Florida

a transverse row of long, branching dark or pale spines (scoli) with reddish bases.



Figure 3. Larva of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: J.F. Butler, University of Florida



Figure 4. Pupa of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: J.F. Butler, University of Florida

Life Cycle and Biology

Eggs are laid singly on the leaves of the host plants. The larvae make leaf nests by silking together several of the smaller leaves of *Parietaria* species or by silking together the sides of the larger leaves of nettles to form a tube. Each caterpillar makes a number of shelters during its lifetime (Wagner 2005). Older larvae chew partially through the petioles of the folded leaf nests causing them to droop (Scott 1986).

Red admirals are most common in moist open areas near woods (Glassberg et al. 2000) where their host plants are commonly found. Males defend territories and fly out to meet females for mating. Red admirals prefer fermenting tree sap, fruit, and fresh dung to flower nectar (Allen 1997, Minno and Minno 1999, Opler and Krizek 1984). However, they also



Figure 5. Folded leaf nest of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer), on false nettle. Credits: D.W. Hall, University of Florida

drink nectar from a variety of flowers (Cech and Tudor 2005, Tooker et al. 2002). They also drink from moist soil (Minno and Minno 1999).

There are two generations per year in the northern parts of the distribution and as many as four or more in the southern U.S. (Scott 1984). In the U.S., adults overwinter in the southern part of the range, and more northerly areas are re-colonized each year (Opler et al. 1984, Scott 1986) with the migrating butterflies sometimes forming mass migrations. Walker (2001) reported spring migrations northward in Florida but no significant southward migration during the fall. There is some disagreement concerning whether or not pupae overwinter. Cech and Tudor (2005) and Wagner (2005) state that pupae may overwinter in some parts of the range, but Scott (1986) is doubtful that pupae overwinter.

Hosts

Larvae feed on herbs in the nettle family (Urticaceae) – most commonly plants in the genera *Urtica*, *Boehmeria*, *Laportea*, and *Parietaria*. In Florida, the most common hosts are: false nettle (*Boehmeria cylindrica* [L.]Sw.), Florida pellitory (*Parietaria floridana* Nutt.), heartleaf nettle (*Urtica chamaedryoides* Pursh) and the naturalized exotic stinging nettle (*Urtica dioica* L.) and burning nettle (*Urtica urens* L.). Canadian wood nettle (*Laportea canadensis* [L.]Wedd.) is commonly utilized farther north but this species is rare in Florida and restricted to the Panhandle. Plant names are from Wunderlin and Hansen (2003) and Wunderlin and Hansen (2008).



Figure 6. False nettle, *Boehmeria cylindrical* [L.]Sw., a host of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: D.W. Hall, University of Florida



Figure 7. Florida pellitory, *Parietaria floridana* Nutt., a host of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: D.W. Hall, University of Florida



Figure 8. Florida pellitory, *Parietaria floridana* Nutt., a host of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: D.W. Hall, University of Florida



Figure 9. Canadian wood nettle, *Laportea canadensis* [L.]Wedd., a host of the red admiral, *Vanessa atalanta rubria* (Fruhstorfer). Credits: D.W. Hall, University of Florida

Selected References

Allen TJ. 1997. The Butterflies of West Virginia and their Caterpillars. University of Pittsburgh Press. Pittsburgh, Pennsylvania. 400 pp.

Cech R, Tudor G. 2005. Butterflies of the East Coast: An Observer's Guide. Princeton University Press. Princeton, New Jersey. 345 pp.

Daniels JC. 2000. Butterflies 1: Butterflies of the Southeast. UF/IFAS. Card Set. SP 273

Daniels JC. 2003. Butterflies of Florida: Field Guide. Adventure Publications, Inc. Cambridge, Minnesota. 256 pp.

Glassberg J, Minno MC, Calhoun JV. 2000. Butterflies through Binoculars: Florida. Oxford University Press. New York, New York. 256 pp.

Minno MC, Butler JF, Hall DW. 2005. Florida Butterfly Caterpillars and their Host Plants. University Press of Florida. Gainesville, Florida. 341 pp.

Minno MC, Emmel TC. 1993. Butterflies of the Florida Keys. Scientific Publishers. Gainesville, Florida. 168 pp.

Minno MC, Minno M. 1999. Florida Butterfly Gardening: A Complete Guide to Attracting, Identifying, and Enjoying Butterflies of the Lower South. University Press of Florida. Gainesville, Florida. 210 pp.

Opler PA, Krizek GO. 1984. Butterflies East of the Great Plains. The Johns Hopkins University Press. Baltimore, Maryland. 294 pp.

Scott JA. 1986. The Butterflies of North America: A Natural History and Field Guide. Stanford University Press. Stanford, California. 583 pp.

Tooker JF, Reigel PF, Hanks LM. 2002. Nectar sources of day-flying Lepidoptera of central Illinois. Annals of the Entomological Society of America. 95: 84-96.

Wagner DL. 2005. Caterpillars of Eastern North America. Princeton University Press. Princeton, New Jersey. 512 pp.

Walker TJ 2001. Butterfly migrations in Florida: seasonal patterns and long term changes. Environmental Entomology 30: 1052-1060.

Wunderlin RP, Hansen BF. 2003. Guide to the Vascular Plants of Florida. 2nd ed. University Press of Florida. Gainesville, Florida. 787 pp.

Wunderlin RP, Hansen BF. (2008). Atlas of Florida Vascular Plants. *Institute for Systematic Botany*. <http://www.plantatlas.usf.edu/> (1 April 2009).