

# Bean, Jack —*Canavalia ensiformis* (L.) D.C.Bean, Sword —*Canavalia gladiata* (Jacq.) D.C.<sup>1</sup>

James M. Stephens<sup>2</sup>

The **jack bean** and the sword bean are very similar and are occasionally grown in home gardens around the state. Other names for them are Chickasaw lima bean, Brazilian broad bean, coffee bean, ensiform bean, horse bean, mole bean, go-ta-ki, overlook bean, Pearson bean, watanka, and raba de burro.

## DESCRIPTION

In both species, pods reach a length of 10 to 14 inches, and a width of 1 to 1½ inches. Seeds are large, ½ to ¾ inch



Figure 1. Jack bean.

long and nearly as broad. The two species differ mainly in the length of the seed hilum (scar). The hilum of sword bean (*C. gladiata*) is more than one-half the length of the seed, whereas that of the jack bean (*C. ensiformis*) is only about one-third as long as the seed. Most of the jack beans identified in Gainesville have been white with a black seed scar, while the sword beans have been red seeded.

## CULTURE

There probably are many varietal and environmental variations in the growth habit of the plants; however, it appears that the jack bean plant is bushy in comparison with the vining nature of the sword bean plant. One North Florida gardener noted that one sword bean plant covered his entire garden of 400 square feet.

## USE

Neither bean has commercial importance in this country. Both are reported to be edible in the young tender immature stage by slicing and boiling the tender pods, or peeling and using the seeds as broad beans. Also, reports have indicated the possibility of mild toxicity of beans when eaten in large quantities. Because of this, the boiling water should be drained off to remove any poisonous substances coming from the beans. Pharmaceutical companies have shown some interest in the beans as a possible source of the enzyme urease, and a few acres have been tried in Florida for that purpose.

1. This document is HS553, one of a series of the Horticultural Sciences Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date April 1994. Revised March 2009. Reviewed January 2012. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. James M. Stephens, professor, Horticultural Sciences Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.