

Cucumber, Armenian — *Cucumis melo* L. (Flexuosus group)¹

James M. Stephens²

Armenian cucumber is known by several other names such as snake cucumber, snake melon, and uri. It should not be confused with the snake gourd or club gourd, *Trichosanthes anguina*.



Figure 1. Armenian cucumber.

Description

Armenian cucumber is closely related to muskmelon, but is elongated like a cucumber. The slender fruit is usually about 3 feet long and 3 inches in diameter, almost always bent and twisted. It is dark green, marked with paler green longitudinal furrows, and is thicker at the blossom end. The fruit changes to yellow when ripe, at which time it has a strong muskmelon odor.

The annual vine is creeping, with slender roundish to angular stems covered with short hairs. The leaves are rounded, almost kidney-shaped with five angles (lobes). Both male and female flowers on the vine are small and pale yellow, with five rounded divisions. Seeds are more like those of a muskmelon than a cucumber.

Individual plants may be found bearing fruits that are long and twisted and at the same time other fruits that are broad and oval. Sometimes even the same fruit will be thin and snake-like near the stem end, but swollen at the other end similar to a melon.

Culture

In Florida, as in most parts of the country, the Armenian cucumber is grown as a curiosity and for pickling. Planted in late August in Gainesville, fruits suitable for use have been produced by late October.

It is a warm season crop, and should be planted from seed in the early spring or early fall in all areas of Florida except South Florida where it may be seeded October through February. Follow the cultural suggestions for cucumber and cantaloupe. Growers should be on the alert for powdery mildew and downy mildew, two common diseases of cucumbers and cantaloupes.

1. This document is HS589, one of a series of the Horticultural Sciences Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date May 1994. Revised March 2009. Reviewed February 2012. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. James M. Stephens, Professor, Horticultural Sciences Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.