

# IRAC's Insecticide Mode of Action Classification<sup>1</sup>

Frederick M. Fishel<sup>2</sup>

*This guide explains the rationale behind the Insecticide Resistance Action Committee's (IRAC) insecticide and acaricide mode of action classification and provides a listing of those insecticide common names with their groupings and primary modes of action for insecticides currently registered in Florida.*

## What is IRAC?

IRAC has groups formed in several countries, including the United States, Brazil, South Africa, Spain, India, and Australia. The group's purpose is to communicate and educate agricultural producers and crop protection professionals by providing resistance management information. Members of an IRAC group are generally professionals who are actively engaged in the insecticide and acaricide manufacturing industry. Some university researchers also participate.

## Resistance to Pesticides

Resistance refers to an inheritable change in the sensitivity of a pest population that is reflected by a repeated product failure to achieve the expected control level when used according to the label recommendation for that pest species. Resistance does not always occur, but it has been documented with insecticides as early as 1914. There are many known instances today where resistance is a problem. Not only has resistance occurred with insecticides, but it has also occurred with other pesticides, such as fungicides,

herbicides, and rodenticides. Complicating the understanding and management of resistance is the problem of knowing which type of resistance is present in a given pest population. Some pest populations are known to have cross-resistance. That is, they are not effectively controlled with pesticides having the same mode of action that generally targets the same site within the pest. For example, both the carbamate and organophosphate insecticides target acetylcholine esterase, although each group of insecticides is chemically different from one another. The greatest resistance concern arises when multiple-resistance is confirmed. Multiple-resistance is the situation of a pest population that is resistant to pesticides having different modes of action. It is the most difficult type of resistance to manage because the number of management options is reduced. For more information on resistance, see UF/IFAS EDIS Document ENY-624, *2014 Florida Citrus Pest Management Guide: Pesticide Resistance and Resistance Management*, <http://edis.ifas.ufl.edu/cg026>.

## IRAC's Classification Scheme

IRAC's insecticide classification scheme is based on mode of action. The goal of the scheme is to provide information to applicators of acaricides and insecticides so they can make sound decisions on selecting insecticides to prevent or manage resistance. Besides selecting products that have different modes of action, growers are also encouraged

1. This document is PI-83, one of a series of the Pesticide Information Office, UF/IFAS Extension. Original publication date October 2005. Revised December 2008, April 2011, April 2014, and March 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Frederick M. Fishel, professor, Agronomy Department, and director, Pesticide Information Office; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition. Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

to integrate other methods into insect and mite control programs. Table 1 contains those acaricides and insecticides registered for use in Florida, though it changes constantly. They are listed according to IRAC's classification scheme by their group and subgroup codes, primary target site of action, chemical sub-group or exemplifying active ingredient, and active ingredient, based on that appearing in version 8.2, 2017, prepared by the IRAC International MoA Working Group.

## Using the IRAC Classification Scheme with Product Labels

IRAC is currently encouraging manufacturers of pesticides to indicate the IRAC mode of action group number and description on their product labels; some registrants are now doing so, especially with newer products. Such information would be helpful in assisting pesticide applicators in the selection of acaricides and insecticides for use in resistance management strategies. See Figure 1 for an example of the manner that IRAC is encouraging registrants to list this information.




Figure 1.

## Additional Information

IRAC: <http://www.irc-online.org/groups/guide/>

Rogers, M.E., and M. M. Dewdney. 2016. *Florida Citrus Pest Management Guide: Pesticide Resistance and Resistance Management*. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/cg026>.

Table 1. IRAC's classification scheme for acaricides and insecticides registered for use in Florida.

Group	Subgroup	Primary Action Target Site	Chemical Subgroup or Exemplifying Active Ingredient	Active ingredients
1*	1A	Acetylcholine esterase inhibitors	Carbamates	Carbaryl Methiocarb Methomyl Oxamyl Propoxur Thiodicarb
	1B		Organophosphates	Acephate Chlorpyrifos Chlorpyrifos-methyl Coumaphos Diazinon Dichlorvos Dicrotophos Dimethoate Disulfoton Ethoprop Fosthiazate Malathion Methamidophos Methidathion Naled Oxydemeton-methyl Phorate Profenofos Temephos Terbufos Tetrachlorvinphos Trichlorfon
2	2A	GABA-gated chloride channel antagonists	Cyclodiene organochlorines	Endosulfan
	2B		Phenylpyrazoles (Fiproles)	Fipronil
3*	3A	Sodium channel modulators	Pyrethroids Pyrethrins	Allethrin and isomers Bifenthrin and isomers Cyfluthrin and isomers Cyhalothrin and isomers Cypermethrin and isomers Cyphenothrin Deltamethrin Esfenvalerate Etofenprox Fenpropathrin Fenvalerate Flumethrin Imiprothrin Permethrin Phenothrin Prallethrin Pyrethrins (pyrethrum) Resmethrin Tefluthrin Tetramethrin Talomethrin

Group	Subgroup	Primary Action Target Site	Chemical Subgroup or Exemplifying Active Ingredient	Active ingredients
4	4A	Nicotinic acetylcholine receptor agonists	Neonicotinoids	Acetamiprid Clothianidin Dinotefuran Imidacloprid Thiamethoxam
	4C		Sulfoximines	Sulfoxaflor
	4D		Butenolides	Flupryadifurone
5		Nicotinic acetylcholine receptor allosteric activators	Spinosyns	Spinetoram Spinosad
6		Chloride channel activators	Avermectins Milbemycins	Abamectin Emamectin benzoate Milbemectin
7	7A	Juvenile hormone mimics	Juvenile hormone analogues	Hydroprene Kinoprene Methoprene
	7B		Fenoxycarb	Fenoxycarb
	7C		Pyriproxyfen	Pyriproxyfen
8	8A	Miscellaneous non-specific (multi-site) inhibitors	Alkyl halides	Methyl bromide and other alkyl halides
	8B		Chloropicrin	Chloropicrin
	8C		Sulfuryl fluoride	Sulfuryl fluoride, cryolite
	8D		Borax	Borax, Boric acid, Disodium octaborate, Sodium borate, Sodium metaborate
9	9B	Modulators of chordotonal organs	Pymetrozine	Pymetrozine, Pyriproxyfen
	9C		Fonicamid	Fonicamid
10	10A	Mite growth inhibitors	Clofentezine Hexythiazox	Clofentezine Hexythiazox
	10B		Etoazole	Etoazole
11		Microbial disruptors of insect midgut membranes (includes transgenic crops expressing <i>B.t.</i> toxins)	<i>Bacillus thuringiensis</i> or <i>Bacillus sphaericus</i> and the insecticidal proteins they produce	<i>Bacillus thuringiensis</i> subsp. <i>israelensis</i> <i>Bacillus sphaericus</i> <i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i> <i>Bacillus thuringiensis</i> subsp. <i>aizawai</i> Bt crop proteins
12	12B	Inhibitors of mitochondrial ATP synthase	Organotin miticides	Fentutatin oxide
	12C		Propargite	Propargite
13		Uncouplers of oxidative phosphorylation via disruption of the proton gradient	Chlorfenapyr	Chlorfenapyr, Sulfluramid
15		Inhibitors of chitin biosynthesis, type 0, Lepidopteran	Benzoylureas	Diflubenzuron Hexaflumuron Lufenuron Novaluron Noviflumuron
16		Inhibitors of chitin biosynthesis, type 1	Buprofezin	Buprofezin
17		Molting disruptor, Dipteran	Cyromazine	Cyromazine

Group	Subgroup	Primary Action Target Site	Chemical Subgroup or Exemplifying Active Ingredient	Active ingredients
18		Ecdysone receptor agonists	Diacylhydrazines	Halofenozide Methoxyfenozide Tebufenozide
19		Octopamine receptor agonists	Amitraz	Amitraz
20*	20A	Mitochondrial complex III electron transport inhibitors	Hydramethylnon	Hydramethylnon
	20B		Acequinocyl	Acequinocyl
21*	21A	METI acaricides and insecticides	Fenazaquin	Fenazaquin Fenpyroximate Pyridaben Tolfenpyrad
	21B	Mitochondrial complex I electron transport inhibitors	Rotenone	Rotenone
22*	22A	Voltage-dependent sodium channel blockers	Indoxacarb	Indoxacarb
	22B		Metaflumizone	Metaflumizone
23		Inhibitors of acetyl CoA carboxylase	Tetronic and tetramic acid derivatives	Spirodiclofen Spiromesifen Spirotetramat
24	24A	Mitochondrial complex IV electron transport inhibitors	Phosphine	Aluminum phosphide Phosphine Zinc phosphide
25	25A	Mitochondrial complex II electron transport inhibitors	Beta-ketonitrile derivatives	Cyflumetofen
28		Ryanodine receptor modulators	Diamides	Chlorantraniliprole Cyantraniliprole Flubendiamide
29		Chordotonal organ modulators— undefined target site	Flonicamid	Flonicamid
UN		Compounds of unknown or uncertain mode of action®	Azadirachtin	Azadirachtin Bifenazate Dicofol Lime sulfur Pyridalyl Sulfur

\*Groups and Sub-groups: Although sharing the same primary target site, it is possible that not all members of a single mode of action class have been shown to be cross-resistant. Different resistance mechanisms that are not linked to the target site, such as enhanced metabolism, may be common for such a group of chemicals. In such cases, the mode of action grouping is further divided into sub-groups.

®A compound with an unknown or controversial mode of action or an unknown mode of toxicity will be held in category UN until evidence becomes available to enable that compound to be assigned to a more appropriate mode of action class.