

## Herbicides: How Toxic Are They?<sup>1</sup>

---

Fred Fishel, Jason Ferrell, Greg MacDonald, and Brent Sellers<sup>2</sup>

A herbicide is any agent used to bring about plant death. Although everything from salt to soap has been used for this purpose, herbicides are primarily synthetic chemicals manufactured for use in the agriculture, industrial, and ornamental and turf industries. For many years these products have been seen as toxins that poison plants and are equally harmful to the applicator. To compound this issue, organically produced food is becoming more popular because it is pesticide-free and is seen as being healthier. These factors have led to a prevalence of opinion that pesticides are bad for the environment and harmful to humans.

Although there have been pesticides that were toxic and dangerous to handle, most of these products are no longer used and have been replaced by newer chemistry. Pesticides now must go through rigorous testing by EPA before they can be sold. This has led to many herbicides that possess little or no mammalian toxicity and are less harmful than many everyday household products (Table 1). Surprisingly, household chemicals that many of us store under the kitchen sink pose more risk to the handler than herbicides.

A common way to document toxicity is by oral LD<sub>50</sub> values. LD<sub>50</sub> is the amount of chemical required to provide a “lethal dose” to 50% of the test population. LD<sub>50</sub> is measured in mg of chemical administered per kg of body weight. Therefore, an oral LD<sub>50</sub> of 500 means that 500 mg of chemical were needed to obtain lethality in a 1 kg subject (rabbit). The lower the LD<sub>50</sub> value, the less chemical that is required to reach lethality. A chemical with an LD<sub>50</sub> of 10 mg/kg is more acutely toxic than one with an LD<sub>50</sub> of 100 mg/kg.

Table 1 demonstrates that herbicides often have higher LD<sub>50</sub> values than many commonly used or consumed products. Why is this? Why are chemicals that are so effective on plant species not equally harmful to humans? The reason is two-fold. First, herbicides target highly specific biological or biochemical processes within plants, such as photosynthesis and production of branch-chain amino acids. However, mammals (humans included) do not photosynthesize or produce branch-chain amino acids. Therefore, herbicides that target photosynthesis or branch-chain amino acid production have no place to bind in our bodies and have very little impact.

- 
1. This document is PI-133, one of a series of the Pesticide Information Office, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date September 2006. Revised November 2007. Reviewed December 2009. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
  2. Fred Fishel, associate professor, Agronomy Department, and director, Pesticide Information Office; Jason Ferrell, assistant professor, Agronomy Department, University of Florida; Greg MacDonald, associate professor, Agronomy Department, University of Florida; Brent Sellers, assistant professor, Agronomy Department, Range Cattle Research and Education Center--Ona, FL; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

**Use herbicides safely. Read and follow directions on the manufacturer's label.**

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean

Secondly, since these herbicides do not bind in our bodies, they are often excreted in urine within 24 hours of the dose. This flushing of the herbicide does not allow concentrations to build up within the body to toxic levels. This in no way means that it is safe to intentionally ingest herbicides, but the fact is that our bodies are well equipped to safely dispose of accidental exposure to many common herbicides.

### What about Agent Orange?

Agent Orange was a herbicide product containing equal parts 2,4-D and 2,4,5-T. These herbicides were used extensively in the Vietnam War to defoliate the jungles in an attempt to expose troop movement. However, many veterans came forward after the war with illnesses ranging from a multitude of cancers to various respiratory diseases. It was speculated that Agent Orange was to blame for these disorders.

After numerous governmental inquiries, it has been learned that the herbicides 2,4-D and 2,4,5-T were not responsible for the human health effects of Agent Orange. Rather, Agent Orange was inadvertently contaminated with dioxin, a potent and known carcinogen. Dioxin was an unanticipated byproduct of the 2,4,5-T synthesis process. Although synthesis processes have been developed that allow 2,4,5-T to be produced without dioxin contamination, these procedures were not employed to fulfill this specific contract. Consequently, the production of 2,4,5-T has been totally banned by the US and numerous other countries. For a detailed discussion of this subject, see <http://www.nap.edu/catalog/11242.html#orgs>.

### Conclusion

It must be noted that some herbicides are harmful. Herbicides such as paraquat and endothal have "Danger" signal words on the label and must be handled with great care. Therefore, it is important that **all** herbicides be handled carefully and in a manner consistent with their labeling. Just because some herbicides are less toxic than table salt does not mean that any herbicides should be handled carelessly. But on the other hand, using a herbicide in accordance with the product label will not often result in personal injury or cause for alarm.

### Additional Information

Committee to Review the Health Effects in Vietnam Veterans of Exposure to Herbicides. 2004. Veterans and Agent Orange. National Academies Press. Washington, D.C. (<http://www.nap.edu/catalog/11242.html#orgs>). Visited September 2006.

Fishel, F.M. 2005. Evaluation of pesticides for carcinogenic potential. UF/IFAS EDIS Publication PI-37. <http://edis.ifas.ufl.edu/PI074> . Visited July 2005.

National Pesticide Information Center (1-800-858-7378 or <http://npic.orst.edu/> ). Visited July 2005.

Nesheim, O.N., F.M. Fishel, and M. Mossler. Toxicity of pesticides. UF/IFAS EDIS Publication PI-13. <http://edis.ifas.ufl.edu/PI008>. Visited August 2006.

Poison Information Center Network (1-800-222-1222 or <http://www.fpicn.org> ). Visited July 2005.

**Table 1.** Comparison of oral LD<sub>50</sub> values for commonly used herbicides and consumer goods.

<b>Herbicide</b>	<b>LD<sub>50</sub><sup>1</sup></b>	<b>Common consumer chemicals</b>	<b>LD<sub>50</sub></b>
Paraquat (Gramoxone)	~100	Nicotine	9
Triclopyr	630	Caffeine	192
2,4-D	666	Bleach	192
Pendimethalin (Prowl)	1050	Tylenol	338
Atrazine	3090	Household ammonia (10%)	350
Glyphosate (Roundup)	4900	Codeine	427
Imazaquin (Image)	>5000	Table salt	3000