

Exotic Diseases of Citrus

M.M. Dewdney, J.D. Yates,
M.E. Rogers, T.M. Spann

CITRUS VARIEGATED CHLOROSIS (CVC)



CVC (upper side of leaf)



CVC (underside of leaf)



CVC (small, hard fruit)

Healthy fruit

LEPROSIS



Leprosis (early symptoms)



Leprosis (advanced symptoms)



Leprosis on fruit




Leprosis on mature fruit




Leprosis bark scaling

SWEET ORANGE SCAB (SOS)



Sweet Orange Scab

CITRUS TRISTEZA



Citrus Tristeza Virus Stem-Pitting

PSEUDOCERCOSPORA FRUIT AND LEAF SPOT DISEASE



Pseudocercospora lesions on young flush



Pseudocercospora lesions on fully expanded fruit



Pseudocercospora lesions on mature fruit

For more information, contact the University of Florida / IFAS Citrus Research and Education Center 863-956-1151, www.crec.ifas.ufl.edu, or your local county citrus extension agent: at http://citrusagents.ifas.ufl.edu/citrus_agents_home_page/citrus_agents_home.html

1. This document is PP264, one of a series of the Department of Plant Pathology, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First published: April 2009. Revised July 2010.

2. Megan M. Dewdney, assistant professor, Department of Plant Pathology, Jamie D. Yates, coordinator for canker and greening extension education, Michael E. Rogers, assistant professor, Department of Entomology, Timothy M. Spann, assistant professor, Department of Horticulture, Citrus REC, Lake Alfred, Florida; Cooperative Extension Service, Institute of Food and Agricultural Sciences; University of Florida; Gainesville, FL 32611.

	CITRUS VARIEGATED CHLOROSIS (CVC)	LEPROSIS	PSEUDOCERCOSPORA FRUIT AND LEAF SPOT DISEASE	SWEET ORANGE SCAB (SOS)	CITRUS TRISTEZA VIRUS STEM-PITTING (CTV-SP)
Plant Tissue Location	Xylem	Cytoplasm or nucleus of infected cells	Fruit rind and leaves	Fruit rind	Phloem
Spread	Sharpshooters, graft-transmitted, possibly seed	Several species of <i>Brevipalpus</i> mites	Windborne conida, rain splash or movement of infected plant material	Requires moisture; presence of susceptible fruit; splash by raindrops	Several species of aphids, budding and grafting
Pathogen	<i>Xylella fastidiosa</i> (Bacterium)	<i>Citrus leprosis virus</i>	<i>Pseudocercospora angolensis</i>	<i>Elsinoë australis</i> (Fungus)	<i>Citrus tristeza virus</i>
Tree Symptoms	Reduced vigor and growth; tree not killed, but becomes unproductive; symptoms may appear in one branch or the whole canopy	Symptoms localized on fruit, leaves, or twigs	Premature defoliation and fruit drop	Unaffected	Scattered pits after removing bark from affected plants; depressed areas in the bark
Leaf Symptoms	Severe chlorosis, reddish to brown lesions on the lower side corresponding to yellow areas on the upper surface; may resemble zinc deficiency in early stages	Chlorotic at first then may become brown with or without a necrotic center; smooth to touch	Circular to irregularly shaped lesions that can coalesce with brown or grayish centers surrounded by a yellow halo	Unaffected	Chlorosis and dieback may occur
Stem/Twig Symptoms	Unaffected	Bark scaling, twig dieback	Unaffected	Unaffected	Twigs are brittle; bark is abnormally thick; trunk and branches may have a ropy appearance
Fruit Symptoms	Small and hard with high acid; fruit remains on the tree; may exhibit sunburn damage; normal color change	Premature fruit drop, flat or depressed lesions with yellow halo; often with brown centers	Lesions are circular to irregularly shaped; young fruit has nipple-like swellings with yellow halo; young fruit can be mummified; mature fruit lesions are dark brown to black and generally flat or sunken with a yellow halo; single or coalescent	Young fruit have corky wart-like pustules; tan to gray in color; mature fruit lesions become smoother	Small and misshapen
Flowering	Off-season bloom	Unaffected	Unaffected	Unaffected	Off-season bloom
Varieties Affected	Sweet orange	Mainly sweet orange, mandarins	All citrus varieties	Sweet orange and tangerines	Lime, grapefruit, sweet orange