

Wireworms in Florida Sugarcane ¹

D. G. Hall, R. H. Cherry, and R. A. Gilbert²

Wireworms, the larval stage of a click beetle, often cause severe damage to numerous crops in Florida. At least twelve species of wireworms have been found in southern Florida, but only the corn wireworm, *Melanotus communis*, is considered to cause significant economic damage to sugarcane. Since *Melanotus communis* is the important wireworm species, the rest of this document will pertain to this species.

Biology

Adult click beetles may be collected during all months of the year, but they are most numerous during May, June, and July and are scarce from September to March. The beetles are brown and about 5/8" long.

The larvae of click beetles are called wireworms because they resemble a jointed piece of stiff, bright, brownish-yellow wire. *Melanotus communis* wireworms are approximately 1-1/4" in length when fully grown (Figure 1). They are soil pests and attack the underground portions of the sugarcane plant.



Figure 1. Wireworm larva.

Wireworms are generally found under sugarcane stools rather than between stools in a row or between rows. Wireworm populations are lowest in the summer and increase rapidly in the fall staying fairly constant thereafter. However, *M. communis* are slightly smaller in the fall than the rest of the year.

Damage

Generally, wireworms are a pest of newly planted sugarcane and only rarely a pest in ratoon sugarcane. Wireworms feed on the buds and root

-
1. This document is ENY-665 (SC013), one of a series of the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First printed: October 1991. Revised: February 2008. This publication is also a part of the Florida Sugarcane Handbook, an electronic publication of the Agronomy Department. For more information you may contact the editor of the Sugarcane Handbook, R. A. Gilbert (ragilbert@gnv.ifas.ufl.edu). Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>.
 2. D. G. Hall, U.S.D.A., Ft. Pierce, FL; R. H. Cherry, professor, Everglades Research and Education Center; and R. A. Gilbert, associate professor, Everglades Research and Education Center, Florida Agricultural Experiment Stations, Institute of Food and Agricultural Sciences, University of Florida. The use of trade names in this publication is solely for the purpose of providing specific information. It is not a guarantee or warranty of the products named and does not signify that they are approved to the exclusion of others of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

primordia during germination of sugarcane seed pieces, and on shoots and roots after germination. Most of the injury to young shoots is near the point where the shoots join the seed piece or stubble. Wireworm injury can generally be identified as relatively large, ragged holes cut into seedpieces and buds, or into young shoots (Figure 2). The death of buds or young shoots leads to stand reductions. Wireworm injury has been reported as facilitating the entrance of the fungus that causes sugarcane red rot disease.




Figure 2. Wireworm damage.

Field studies have shown that significant stand reductions during early plant growth may occur at infestation levels of fewer than 4 wireworms per 5 feet of row. One recent study showed that one wireworm per 5 row feet could cause 6.2-7.8% stand reduction at twelve weeks after planting. Tillering may compensate for some of the early stand losses, but, at harvest, stalks from infested sugarcane fields tend to weigh less.

In a large plot study, the weight of cane harvested was reduced by 3.8% per wireworm per 5 feet of row. Stand reductions in plant cane caused by 8 to 12 wireworms per 5 feet of row carried over into the first ratoon crop and resulted in lower ratoon crop yield.

Control

Biological

Although there are a small number of insect parasites that attack wireworms, biological control of wireworms in south Florida by beneficial insects and diseases has generally been considered to be insignificant. Birds such as the cattle egret may sometimes be of value in reducing wireworm levels because they may consume wireworms that are exposed at the surface of the soil when a field is disked or cultivated.

Cultural Practices

Flooding for wireworm control can be effective but is a slow process and may not be practical. More studies are needed, but the current information suggests a minimum of six weeks of continuous flooding is needed during the summer to obtain wireworm control. Longer flooding durations are needed during colder months.

Flooding during late spring and summer will kill the wireworms and also prevent egg-laying by the adult click beetles. Fallow field flooding or growing rice as a rotation crop may eliminate the need to use a soil insecticide at sugarcane planting the following fall. Growing a ratoon rice crop may offer slightly increased control during years when click beetle activity extends into July and August.

Chemical

Soil insecticides are generally used in newly planted sugarcane for wireworm control. Insecticides are not used for wireworm control in ratoon sugarcane. No resistance to insecticides has been reported to occur in wireworms in Florida sugarcane.