

Managing Against the Development of Herbicide Resistant Weeds: Sugarcane ¹

B. A. Sellers, J. A. Ferrell, and G. E. MacDonald²

Producing a profitable sugarcane crop in Florida depends, in part, on effectively controlling weeds that can reduce yields. Because they provide an efficient and cost effective means of weed control, herbicides are a critical component of sugarcane weed management programs. However, in many cropping systems excessive use of a single herbicide, or group of herbicides with the same site of action, has resulted in the development of herbicide resistant weeds (for more information refer to EDIS document SS-AGR-243, *Herbicide Resistant Weeds*). When herbicide resistant weed populations appear, standard weed control treatments often become ineffective. As a result, alternative means of control must be used. In crops, such as sugarcane, where a limited number of herbicides are registered, the loss of a single effective herbicide can be very costly. Thus, it is critical to manage herbicides in order to prevent or delay the development of herbicide resistant weed populations.

In order to successfully manage herbicides against the development of herbicide-resistant weeds,

you must have a basic understanding of which herbicides have the same site of action. Table 1 lists herbicides by group number, site of action, chemical family, common name, and trade name.

When planning a herbicide program to manage against herbicide resistance, it is ideal to avoid using a single herbicide or herbicide group in consecutive years. However, because of the limited number of herbicides available and the perennial crop cycle of sugarcane Group 4 (2,4-D), Group 5 (atrazine, ametryn, metribuzin, and other triazines), and Group 18 (asulam) herbicides are typically used in every year of a sugarcane crop.

Worldwide over 60 weed species have developed resistance to the triazine herbicides. These biotypes include several members of the genera *Amaranthus*, *Chenopodium*, *Panicum*, and *Solanum*, which are commonly found in Florida sugarcane fields. Consequently, it is critical that other herbicide groups be utilized as part of an integrated weed

1. This document is SS-AGR-244, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date February 2006. Reviewed October 2008. This publication is also part of the Florida Sugarcane Handbook, an electronic publication of the Agronomy Department. For more information, you may contact the editor of the Sugarcane Handbook, R.A. Gilbert (ragilbert@ifas.ufl.edu). Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. B. A. Sellers, assistant professor, Agronomy Department, Range Cattle Research and Education Center--Ona, FL; J. A. Ferrell, assistant professor, Agronomy Department; G. E. MacDonald, associate professor, Agronomy Department; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. Use herbicides safely. Read and follow directions on the manufacturer's label.

control program to prevent the development of triazine resistant weed populations. Although there are no reported cases of resistance to asulam, there is always a chance that resistant populations could develop. Until recently, asulam (Group 18) was the only herbicide that could be used for postemergence control of grass weeds in sugarcane. However, the recent registration of Envoke (Group 2) provides an alternate site of action for postemergence grass weed control. For most grassy weeds, tank mixtures of asulam and Envoke are an effective resistance management strategy. Herbicide resistance is more likely to be a problem in fields successively planted to sugarcane. Rotational crops and fallow periods provide a valuable opportunity to control weeds using tillage, flooding, or herbicides with different sites of action.

Although it is likely that small populations of herbicide resistant weeds are already present in the EAA, herbicide resistance is currently not a significant problem. The continued use of integrated and properly managed weed control programs should ensure that resistance does not become a major issue in the future.

Table 1. Group number and site of action of herbicides commonly used in sugarcane and crops grown in rotation with sugarcane.

Group number and site of action	Chemical Family	Common Name	Trade Name(s)	Crop used in
Group 1 Acetyl CoA carboxylase (ACCase) inhibitors	aryloxyphenoxy-propanoates	fenoxaprop	Acclaim	sod
		fluzafop quizalofop	Fusilade DX Assure II	fallow, canal banks vegetables
Group 2 Acetolactate synthase (ALS) inhibitors	cyclohexanediones	clethodim	Select	vegetables
		sethoxydim	Poast, Poast plus	vegetables
		bensulfuron-methyl	Londax	rice
		chlorsulfuron halosulfuron-methyl	Corsair Semptra, Sandea	sod sugarcane
Group 3 Microtubule assembly inhibitors	sulfonyleureas	nicosulfuron	Accent	sweet corn
		trifloxysulfuron-sodium	Envoke	sugarcane
		bispyribac-sodium	Regiment	rice
		oryzalin	Snapshot, Surfian	sod
Group 4 Synthetic auxins	dinitroanilines	pendimethalin	Prowl 3.3, Prowl H2O, Pendimax	sugarcane
		prodiamine	Barricade	sod
		2,4-D	several	sugarcane, rice, sweet corn
Group 5 Photosystem II inhibitors	phenoxy acetic acids	dicamba	Banvel	sugarcane
		ametryn	Evik	sugarcane
		atrazine	Aatrex, Bicep II Magnum ²	sugarcane, sweet corn
Group 6 Photosystem II inhibitors (same site as group 5, but different binding characteristics)	benzothiadiazoles	hexazinone	K4 ¹	sugarcane
		metribuzin	Sencor, Lexone	sugarcane
		simazine	Princep, Simazine	sweet corn
		bentazon	Basagran	sweet corn, rice, vegetables

Table 1. Group number and site of action of herbicides commonly used in sugarcane and crops grown in rotation with sugarcane.

Group number and site of action	Chemical Family	Common Name	Trade Name(s)	Crop used in
Group 7 Photosystem II inhibitors (same site as group 5 and 6, but different binding characteristics)	Ureas	diuron	Karmex, K4 ¹	sugarcane
		linuron	Lorox	vegetables
Group 8 Lipid synthesis inhibition (not ACCase inhibition)	Amide	propanil	Stam M-4	rice
		napropamide	Devrinol	sod
Group 9 EPSP synthase inhibitors	Thiocarbamates	butylate	Sutan	sweet corn
		EPTC	Eradicane	sweet corn
Group 14 Protoporphyrinogen oxidase (PPO) inhibitors	no family name	thiobencarb	Bolero	rice
		glyphosate	Roundup, Touchdown, others	fallow
Group 15 unknown site of action	Aryl triazinone	carfentrazone	Aim	sugarcane, rice, sweet corn
	Diphenylethers	acifluorfen	Ultra Blazer	rice
		oxyfluorfen	Galligan, Goal	sweet corn
	N-phenylphthalimides	flumioxazin	Valor SX	sugarcane
Group 16 unknown site of action	Oxadiazole	oxadiazon	Ronstar	sod
	Chloroacetamides	metolachlor	Dual Magnum, Pennant	sweet corn, sod
Group 18 DHP (dihydropteroate synthase step) inhibitors		pronamide	Magnum	sod
		ethofumesate	Kerb	sod
Group 21 Cell wall synthesis inhibitor (site B)	Benzofuran	asulam	Prograss	sugarcane
	Carbamate	isoxaben	Gallery	sod
Group 22 Photosystem I electron diversion	Benzamide	paraquat	Gramoxone Extra	fallow
	Bipyridyliums	mesotrione	Callisto	sweet corn
Group 28 Hydroxyphenyl-pyruvate-dioxygenase inhibitors				