


UF/IFAS Nutrient Management Series: Computational Tools for Field Implementation of the Florida Phosphorus Index - Okeechobee Watershed Area ¹

G.W. Hurt, R.S. Mylavarapu and S.P. Boetger²



1. This is Circular 1339, one of a series of circulars from the Soil and Water Science Department, UF/IFAS Extension Originally published in July 2006. Revised August 2013. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. G.W. Hurt, National Leader for Hydric Soils, USDA-NRCS, R.S. Mylavarapu, Nutrient Management Specialist and Director of UF/IFAS ARL/ESTL, Soil and Water Science Department, University of Florida, Gainesville, FL 32611 and S.P. Boetger, Agronomist, NRCS, Gainesville, FL.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other extension publications, contact your county Cooperative Extension service. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, Dean.

CONTENTS

CONTENTS..... 2
PURPOSE..... 3
SCIENTIFIC SUPPORT 3
INTRODUCTION 3
COMPONENTS OF THE P INDEX..... 4
FIELD EVALUATION AND IMPLEMENTATION FOR OKEECHOBEE COUNTY..... 5
Phosphorus Transport Potential Due to Site & Transport Characteristics – Part A (Table 1)..... 5
Soil Erosion..... 5
C-Factors for Dual Cropping Systems..... 11
Soil Erosion Calculation Example 13
Runoff Potential 13
Artificial Drainage 15
Leaching Potential 16
Phosphorus Runoff and Leaching Potentials Ratings for Florida Soil Survey Map Units..... 17
Potential to Reach Water Body..... 18
Phosphorus Transport Potential Due to Phosphorus Source Management - Part B (Table 2) 19
Fertility Index Value:..... 19
Waste Water Application Volume:..... 19
RESULTING P INDEX..... 20
Conservation Planning Notes..... 20
REFERENCES 22
APPENDIX I. 24

PURPOSE

This Circular contains tables with numerical values for each of the various factors listed in the Florida P Index to be used for computation, with examples, during field implementation.

NOTE: This Circular was developed as a source of information and guidance for preparing nutrient management plans for agricultural farms in Florida, specifically to address Phosphorus management through manure/organic by-product applications. This material is therefore intended for any and all agricultural professionals with sufficient training and background in nutrient management to be certified as a Nutrient Management Specialist. This publication is NOT intended for those individuals seeking basic information regarding agricultural nutrients and their environmental impact.

Detailed information about the Florida Phosphorus Index (P-Index) including background, developmental process and considerations can be obtained from the NRCS Field Office Technical Guide (Florida Phosphorus Index Work Group. 2000) or by contacting any of the work group member listed below. The electronic Field Office Technical Guide (eFOTG) can be found at <http://efotg.nrcs.usda.gov/treemenuFS.aspx?Fips=12001&MenuName=menuFL.zip> (The Florida Phosphorus Index sheets are located in Section IV of the Table of Contents under C.Tools.) It is important that the reader has understood the concept and scope of the P index as described on the website before actual field evaluation and implementation. These fact sheets are available for each of the 67 counties in Florida as part of the Nutrient Management series, Circular 1263 and Circular 1273 through 1338, on the internet at: http://edis.ifas.ufl.edu/TOPIC_SERIES_Florida_Phosphorous_Index .

SCIENTIFIC SUPPORT

The following individuals are members of the Florida Phosphorus Index Work Group and were instrumental in the development of the P Index:

University of Florida, Institute of Food and Agricultural Science (UF/IFAS): D.A. Graetz, V.N. Nair, W.G. Harris, G. Kidder, K.L. Campbell, R.S. Mylavarapu, and R.D. Rhue.

Natural Resources Conservation Service (NRCS): S.P. Boetger, G.W. Hurt, W.G. Henderson, W.R. Reck, N. Watts, P.B. Deal and W.D. Tooke.

Florida Department of Agriculture and Consumer Service (FDACS): J.C. Love and D. Smith.

INTRODUCTION

The Phosphorus Index (P Index) is a site-specific, qualitative vulnerability assessment tool. This tool allows a conservation planner to determine the sites that are potentially most vulnerable to off-site movement of phosphorus. The P Index is used to determine whether application of manure/organic by-products should be based on either a nitrogen-based budget or a phosphorus-based budget. The P Index is NOT to be used in any area designated as phosphorus-limited by legislation (e.g. Everglades, Green Swamp, and Okeechobee Basin) to determine if a nitrogen-based nutrient budget can be used. These areas are to have phosphorus-based nutrient budgets regardless of the nutrient source or soil type. The P Index should, however, be used to implement conservation practices to reduce phosphorus movement in these areas.

The purpose of the P Index is to aid planners and others in the decision-making process involved in designing conservation plans related to land application of animal wastes. The P Index is not intended to be an evaluation tool to determine compliance of water quality standards by any regulatory agency. Any attempt to use the P Index as a regulatory tool would be grossly beyond the intent of the concept and philosophy of the P Index developers.

The P Index is a science-based decision-making tool that will support conservation planning and component planning of nutrient management. Concerns regarding P management of manure/organic by-product recycling can be effectively communicated to landowners if the P Index is consistently applied.

COMPONENTS OF THE P INDEX

The P Index assesses two major categories of characteristics: (1) those related to site and transport – Part A (Table 1); and (2) those related to phosphorus sources – Part B (Table 2). The P Index results are then obtained by multiplying the total for Part A by the total for Part B.

$$\text{P Index} = \text{Total for Part A (Site and Transport)} \times \text{Total for Part B (Source Management)}$$

Table 1. Phosphorus Index Worksheet – Part A

Part A: Transport Potential Due to Site and Transport Characteristics						
Site and Transport Characteristics	Phosphorus Transport Rating					Value
Soil Erosion	No Surface Outlet 0	<5T/A ^a 1	5-10 T/A 2	10-15 T/A 4	>15 T/A 8	
Runoff Potential	Very Low 0	Low 1	Medium 2	High 4	Very High 8	
Leaching Potential	Very Low 0	Low 1	Medium 2	High 4	Very High 8	
Potential To Reach Water Body	Very Low 0	Low 1	Medium 2	High 4		
Total for Part A: Site and Transport ^b						
^a T/A = tons per acre. ^b If the sum for Part A is 0 (zero), then change the sum to 1 (one).						

Table 2. Phosphorus Index Worksheet – Part B

Part B: Transport Potential Due to Phosphorus Source Management						
Phosphorus Source Management	Phosphorus Loss Rating					Value
Fertility Index Value	Soil Fertility Index x 0.025 (____ ppm P x 2 x 0.025) ^c					
P Application Source and Rate ^d	0.05 x (____ lbs P ₂ O ₅) for fertilizer, manure or compost 0.015 x (____ lbs P ₂ O ₅) for biosolids 0.10 x (____ lbs P ₂ O ₅) for waste water					
Application Method	No Surface Outlet Or Solids incorporated immediately or injected 0	Applied via irrigation Or Solids incorporated within 1 day of application 2	Solids incorporated within 5 days of application ^e 4	Solids not incorporated within 5 days of application 6		
Waste Water Application	0.20 x ____ acre inches/year					
Total for Part B: Phosphorus Source						
^c From soil test (Mehlich-3) results. ^d Initial evaluation should be N-based rates. ^e Solids include fertilizers, composts, biosolids, and manure and other animal wastes.						

The result of an analysis using the P Index gives the producer a vulnerability rating for each field or portion of a field analyzed (Table 3). This rating may be LOW, MEDIUM, HIGH, or VERY HIGH. As the vulnerability rating increases, so does the potential for phosphorus transport off-site, and for phosphorus to become associated with water quality impairment.

Table 3: Assessing the P Index Results

P Index for Site	Generalized Interpretation of P Index for Site
<75	LOW potential for P movement from the site. If current practices are maintained there is a low probability of an adverse impact to surface waters from P losses at this site. N-based nutrient management planning is satisfactory for this site. Soil P levels and P loss potential may increase in the future due to N-based nutrient management.
75-150	MEDIUM potential for P movement from this site. The chance for an adverse impact to surface waters exists. <i>Nitrogen-based nutrient management planning is satisfactory for this site when conservation measures are taken to lessen the probability of P loss.</i> Soil P levels and P loss potential may increase in the future due to N-based nutrient management.
151-225	HIGH potential for P movement from the site and for an adverse impact on surface waters to occur unless remedial action is taken. Soil and water conservation and P management practices are necessary (if practical) to reduce the risk of P movement and water quality degradation. If risk cannot be reduced then a P-based management budget based on soil test crop P requirements will be utilized.
>225	VERY HIGH potential for P movement from the site and for an adverse impact on surface waters. Remedial action is required to reduce the risk of P movement. All necessary soil and water conservation practices, plus a P-based management plan must be put in place to avoid the potential for water quality degradation. The P-based management plan will be based on soil test crop requirement to reduce P over a defined period (not to exceed 20 years).

FIELD EVALUATION AND IMPLEMENTATION FOR OKEECHOBEE WATERSHED AREA

Phosphorus Transport Potential Due to Site & Transport Characteristics – Part A (Table 1)

Phosphorus transport potential due to site and transport characteristics is as follows:

- Soil Erosion
- Runoff Potential
- Leaching Potential
- Potential to Reach Water Body

Soil Erosion

Soil erosion by water is defined as the loss of soil along a slope or unsheltered distance and is estimated from erosion prediction models. Soil erosion is not calculated for sites that have No Surface Outlet. For all other sites soil erosion by water is predicted using the Revised Universal Soil Loss Equation (RUSLE). RUSLE is used in this index to indicate an average annual long-term movement of soil, thus potential for sediment and attached P movement toward a water body. The RUSLE methodology presented here is a simplified version of that presented in Chapter 6, Florida Agronomy Field Handbook (Florida Ecological Sciences Staff. 1999, as revised) which is available from any NRCS office. Version 2 of the Revised Universal Soil Loss Equation (RUSLE 2) uses factors that represent the effects of climatic erosivity, soil erodibility, topography, cover-management and support practices to compute erosion. This Circular

provides values for calculating only RUSLE. However, those users that choose to use RUSLE2 may download the details from the following website:

http://fargo.nserl.purdue.edu/rusle2_dataweb/RUSLE2_Index.htm.

The average annual erosion expected on fields is computed by:

$$A = R * K * LS * C * P$$

Where:

A is the average soil loss. **A** is a computed value expressed in tons/acre/year.

R is the rainfall factor. For Charlotte County the R-factor is 475.
 For Glades County the R-factor is 475.
 For Hendry County the R-factor is 500.
 For Highlands County the R-factor is 475.
 For Lake County the R-factor is 450.
 For Martin County the R-factor is 550.
 For Okeechobee County the R-factor is 500.
 For Orange County the R-factor is 460.
 For Osceola County the R-factor is 475.
 For Palm Beach County the R-factor is 600.
 For Polk County the R-factor is 475.
 For St. Lucie County the R-factor is 525.

K is the soil erodibility factor. K-factor values are soil specific (see Appendix 1 for these values).

K-factors presented in Appendix 1 are values to be used in conjunction with the soil survey of the designated County if the surface texture of a field is the same as reported in the soil survey. The soil survey is available at the local NRCS field office. Since K-factors presented in the soil survey are only interpretations, they **should be confirmed by on-site investigations**. Where surface textures differ from those in the soil survey, the following K-factors should be used: muck = 0.2, mucky sand = 0.05, sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, and clay = 0.37.

LS is the topographic factor. Slope length (L) begins where runoff starts and ends where slope decreases and deposition begins, or it is the horizontal distance between terraces, or it includes the entire width of contoured or contour strip-cropped fields without terraces. L is expressed in feet and must be determined on-site. Average slope lengths in Okeechobee Watershed Area range from 40 to 120 feet. Slope (S) is the ratio of horizontal distance to vertical distance. S is expressed in percent and must be determined on-site.

Table 4, Table 5, and Table 6 contain common LS-factors for the Okeechobee Watershed Area. Additional LS-factors are available in Chapter 6, Florida Agronomy Field Handbook (Florida Ecological Sciences Staff, 1999).

Table 4. Values for topographic factor (LS) for rangeland and other land uses with cover.

Slope (%)	Horizontal slope length (ft.)						
	9	25	50	75	100	150	200
0.2	0.05	0.05	0.05	0.05	0.05	0.05	0.05
0.5	0.08	0.08	0.08	0.08	0.09	0.09	0.09
1.0	0.12	0.13	0.13	0.14	0.14	0.15	0.15
2.0	0.20	0.21	0.23	0.25	0.26	0.27	0.28
3.0	0.26	0.29	0.33	0.36	0.38	0.40	0.43
4.0	0.33	0.36	0.43	0.46	0.50	0.54	0.58
5.0	0.38	0.44	0.52	0.57	0.62	0.68	0.73
6.0	0.44	0.50	0.61	0.68	0.74	0.83	0.90
8.0	0.54	0.64	0.79	0.90	0.99	1.12	1.23
10.0	0.65	0.81	1.03	1.19	1.31	1.51	1.67
12.0	0.75	1.01	1.31	1.52	1.69	1.97	2.20
14.0	0.85	1.20	1.58	1.85	2.08	2.44	2.74
16.0	0.95	1.38	1.85	2.18	2.46	2.91	3.28
20.0	1.11	1.74	2.37	2.84	3.22	3.85	4.38

Table 5. Values for topographic factor (LS) for row-cropped agricultural and other land uses with little-to-moderate cover.

Slope (%)	Horizontal slope length (ft.)						
	9	25	50	75	100	150	200
0.2	0.05	0.05	0.05	0.05	0.05	0.05	0.05
0.5	0.07	0.08	0.08	0.08	0.09	0.09	0.09
1.0	0.11	0.12	0.13	0.14	0.14	0.15	0.16
2.0	0.17	0.19	0.22	0.25	0.27	0.29	0.31
3.0	0.22	0.25	0.32	0.36	0.39	0.44	0.48
4.0	0.26	0.31	0.40	0.47	0.52	0.60	0.67
5.0	0.30	0.37	0.49	0.58	0.65	0.76	0.85
6.0	0.34	0.43	0.58	0.69	0.78	0.93	1.05
8.0	0.42	0.53	0.74	0.91	1.04	1.26	1.45
10.0	0.50	0.67	0.97	1.19	1.38	1.71	1.98
12.0	0.58	0.84	1.23	1.53	1.79	2.23	2.61
14.0	0.65	1.00	1.48	1.86	2.19	2.76	3.25
16.0	0.72	1.15	1.73	2.20	2.60	3.30	3.90
20.0	0.85	1.45	2.22	2.85	3.40	4.36	5.21

Table 6. Values for topographic factor (LS) for freshly prepared construction and other highly disturbed soil conditions with little or no cover.

Slope (%)	Horizontal slope length (ft.)						
	9	25	50	75	100	150	200
0.2	0.05	0.05	0.05	0.05	0.05	0.05	0.05
0.5	0.07	0.07	0.08	0.08	0.09	0.09	0.10
1.0	0.09	0.10	0.13	0.14	0.15	0.17	0.18
2.0	0.13	0.16	0.21	0.25	0.28	0.33	0.37
3.0	0.17	0.21	0.30	0.36	0.41	0.50	0.57
4.0	0.20	0.26	0.38	0.47	0.55	0.68	0.79
5.0	0.23	0.31	0.46	0.58	0.68	0.86	1.02
6.0	0.26	0.36	0.54	0.69	0.82	1.05	1.25
8.0	0.32	0.45	0.70	0.91	1.10	1.43	1.72
10.0	0.37	0.57	0.91	1.20	1.46	1.92	2.34
12.0	0.45	0.71	1.15	1.54	1.88	2.51	3.07
14.0	0.45	0.85	1.40	1.87	2.31	3.09	3.81
16.0	0.56	0.98	1.64	2.21	2.73	3.68	4.56
20.0	0.67	1.24	2.10	2.86	3.57	4.85	6.04

C is the cover management factor. **C** is defined as the ratio of soil loss from an area with specified cover and management to soil loss from an identical area in tilled continuous fallow. **C**-factors for most crop-management systems have been computed and are listed in two tables. Table 7 contains **C**-factors for cultivated fields and pasture land and Table 8 contains **C**-factors for other agronomic land uses. The higher the number the higher the potential soil loss.

Table 7. C-Factor - Cover Management Factor (Cultivated Fields and Pasture Land).

Cover/Management	Remarks	C-Factor
Bahiagrass/Bermuda grass	Established with no grazing and no haying	0.006
Bahiagrass/Bermuda grass	From planting to 4 years, grazed	0.067
Bahiagrass/Bermuda grass	From planting to 4 years, hayed	0.057
Bahiagrass/Bermuda grass	From planting to 5 years, grazed	0.055
Bahiagrass/Bermuda grass	From planting to 6 years, grazed	0.047
Corn Conventional Tilled	Average yield 80 bushels/acre/year - 30 inch rows	0.348
Corn Conventional Tilled	Average yield 112 bushels/acre/year - 30 inch rows	0.253
Corn Conventional Tilled	Average yield 125 bushels/acre/year - 30 inch rows	0.229
Corn Conventional Tilled	Average yield 150 bushels/acre/year - 30 inch rows	0.198
Corn Conservation Tillage	Average yield 80 bushels/acre/year - 30 inch rows	0.282
Corn Conservation Tillage	Average yield 112 bushels/acre/year - 30 inch rows	0.187
Corn Conservation Tillage	Average yield 125 bushels/acre/year - 30 inch rows	0.180
Corn Conservation Tillage	Average yield 150 bushels/acre/year - 30 inch rows	0.136
Corn No Till	Average yield 80 bushels/acre/year - 30 inch rows	0.140
Corn No Till	Average yield 112 bushels/acre/year - 30 inch rows	0.082
Corn No Till	Average yield 125 bushels/acre/year - 30 inch rows	0.068
Corn No Till	Average yield 150 bushels/acre/year - 30 inch rows	0.051
Cotton Conventional Tilled	Average yield 500lbs/acre/year - 30 inch rows	0.375
Cotton Conventional Tilled	Average yield 500lbs/acre/year - 38 inch rows	0.436
Cotton Conventional Tilled	Average yield 750lbs/acre/year - 30 inch rows	0.310
Cotton Conventional Tilled	Average yield 750lbs/acre/year - 38 inch rows	0.381
Cotton Conventional Tilled	Average yield 1000lbs/acre/year - 30 inch rows	0.271
Cotton Conventional Tilled	Average yield 1000lbs/acre/year - 38 inch rows	0.298
Cotton Conservation Tillage Residue Not Removed	Planted in Rye Average yield 500lbs/acre/year - 30 inch rows	0.079
Cotton Conservation Tillage Residue Not Removed	Planted in Rye Average yield 500lbs/acre/year - 38 inch rows	0.094
Cotton Conservation Tillage Residue Not Removed	Planted in Rye Average yield 750lbs/acre/year - 30 inch rows	0.062
Cotton Conservation Tillage Residue Not Removed	Planted in Rye Average yield 750lbs/acre/year - 38 inch rows	0.081
Cotton Conservation Tillage Residue Not Removed	Planted in Rye Average yield 1000lbs/acre/year - 30 inch rows	0.050
Cotton No Till, Planted in last years cotton residue	Average yield 500lbs/acre/year - 30 inch rows	0.143

Table 7 (cont.). C-Factor - Cover Management Factor (Cultivated Fields and Pasture Land).

Cover/Management	Remarks	C-Factor
Cotton No Till, Planted in last years cotton residue	Average yield 500lbs/acre/year - 38 inch rows	0.177
Cotton No Till, Planted in last years cotton residue	Average yield 750lbs/acre/year - 30 inch rows	0.100
Cotton No Till, Planted in last years cotton residue	Average yield 750lbs/acre/year - 38 inch rows	0.137
Peanut Conventional Till Residue Not Removed	Average yield 2000lbs/acre/year - 36 inch rows	0.371
Peanut Conventional Till Residue Not Removed	Average yield 3000lbs/acre/year - 36 inch rows	0.281
Peanut Conventional Till Residue Not Removed	Average yield 4000lbs/acre/year - 36 inch rows	0.230
Peanut Conventional Till Residue Removed	Average yield 2000lbs/acre/year	0.534
Peanut Conventional Till Residue Removed	Average yield 3000lbs/acre/year	0.449
Peanut Conventional Till Residue Removed	Average yield 4000lbs/acre/year	0.436
Peanut Conservation Tillage Planted in Rye	Average yield 2000lbs/acre/year - Residue Removed	0.479
Peanut Conservation Tillage Planted in Rye	Average yield 3000lbs/acre/year - Residue Removed	0.362
Peanut Conservation Tillage Planted in Rye	Average yield 4000lbs/acre/year - Residue Removed	0.269
Peanut No Till, Residue Not Removed	Average yield 3000lbs/acre/year	0.084
Peanut No Till	Average yield 3000lbs/acre/year - Residue Removed	0.154
Peanut No Till Planted in Rye	Average yield 3000lbs/acre/year - Residue Removed	0.089
Ryegrass, grazed		0.273
Rye, grazed	2800lbs/acre Residue Remaining	0.113
Rye, not grazed	4200lbs/acre Residue Remaining	0.080
Soybeans	Average yield 35 bushels/acre/year	0.355
Watermelon		0.320
Watermelon	With good summer weed or grass cover	0.173
Watermelon	Followed by rye, not grazed	0.269
Weed/Grass, idle	With good summer weed/grass cover	0.079
Weed/Grass, idle	With good summer and winter weed/grass cover	0.035
Weed/Grass, idle	With good winter weed/grass cover	0.245

Table 8. C-Factor - Cover Management Factor for Groves/Orchards (citrus, blueberries, etc.) Rangeland, Disturbed Forest Land, and Long-Term Hay Land, and Idle Land.

Vegetation Canopy Type			Percentage Surface Contact of Ground Cover					
Type and Height of Canopy	Canopy Cover ^a	Type ^b	0	20	40	60	80	>95
No appreciable canopy		G	.450	.200	.100	.013	.013	.003
		W	.450	.240	.150	.090	.043	.011
Tall weeds/short brush ^c	25	G	.360	.170	.090	.038	.012	.003
		W	.360	.200	.130	.082	.041	.011
	50	G	.260	.130	.070	.035	.012	.003
		W	.260	.160	.110	.075	.039	.011
	75	G	.170	.100	.060	.031	.011	.003
		W	.170	.120	.090	.067	.038	.011
Brush or bushes ^d	25	G	.400	.180	.090	.040	.013	.003
		W	.400	.220	.140	.085	.042	.011
	50	G	.340	.160	.085	.038	.012	.003
		W	.340	.190	.130	.081	.041	.011
	75	G	.280	.140	.080	.036	.012	.003
		W	.280	.170	.120	.077	.040	.011
Trees ^e	25	G	.420	.190	.100	.041	.013	.003
		W	.420	.230	.140	.087	.042	.011
	50	G	.390	.180	.090	.040	.013	.003
		W	.390	.210	.140	.085	.042	.011
	75	G	.360	.170	.090	.039	.012	.003
		W	.360	.200	.130	.083	.041	.011

^a Percent of total surface area hidden from view by canopy.
^b G = Surface cover is grass, grasslike plants, and/or decaying litter at least 2 inches thick.
W = Surface cover is broadleaf herbaceous plants and/or decaying litter less than 2 inches thick.
^c Average height that water drops from canopy in autumn is less than 3 feet.
^d Average height that water drops from canopy in autumn is 3 to 12 feet.
^e Average height that water drops from canopy in autumn is more than 12 feet.

C-Factors for Dual Cropping Systems

The C-factor for dual cropping systems is determined by averaging the individual C-Factors. For example, bahiagrass (6 years, grazed) followed by ryegrass, grazed would have a C-Factor calculated as follows:

C-Factor for bahiagrass, 6 years grazed 0.047 (from Table 7) plus C-Factor for ryegrass, grazed 0.273 (from Table 7) divided by 2 equals a C-Factor of 0.160 .

P is the support practice factor. **P** is the ratio of soil loss with a conservation support practice (contour cropping, contour strip cropping, or terracing) to soil loss with straight-row farming up and down the slope. P-factors for these conservation support practices have been computed and are listed in Table 9, Table 10, and Table 11).

The methodology provided herein to calculate P-Factor is a simplified version. A more thorough methodology is explained in Chapter 6, Florida Agronomy Field Handbook, NRCS.

Table 9. P-Factors for Up and Down Hill Cropping and Contour Cropping.

Land Slope Percent	Up and Down Hill Farming P-Factor	Contour Farming P-Factor
1.1 to 2	1.0	0.60
2.1 to 7	1.0	0.50
7.1 to 12	1.0	0.60
12.1 to 18	1.0	0.80
18.1 to 24	1.0	0.90

Table 10. P-Factors for Contour Strip Cropping.

Land Slope Percent	P-Factor ^a	P-Factor ^b	P-Factor ^c	Contour Strip Width (feet) ^d	Maximum Slope Length (feet) ^e
1.0 to 2.5	0.30	0.45	0.60	130	800
2.6 to 5.5	0.25	0.38	0.50	100	600
5.6 to 8.5	0.25	0.38	0.50	100	400
8.6 to 12.5	0.30	0.45	0.60	80	240
12.6 to 16.5	0.35	0.52	0.70	80	160
16.5 to 20.5	0.40	0.60	0.80	60	120
21.5 to 25	0.45	0.68	0.90	50	100

^a For 4-year rotation of row crop, small grain with grass seeding, and 2 years of grass. A second row crop can replace the small grain if grass is established following harvest.

^b For 4-year rotation of 2 years of row crops, 1 year of winter grain with grass seeding, and 1 year of grass.

^c For alternative strips of row crop and small grain.

^d Adjust strip width limits, generally downward, to accommodate widths of equipment.

^e Length limits may be increased by 10 percent if residue cover after crop planting will regularly exceed 50 percent.

Table 11. P-Factors for Terraces.

Horizontal Interval (feet)	Closed Outlet ^a	Open Outlets with Percent Channel Grade Indicated ^b		
		P-Factor	P-Factor for 0.1-0.3	P-Factor for 0.4-0.7
<110	0.50	0.60	0.70	1.0
111-140	0.60	0.70	0.80	1.0
141-180	0.70	0.80	0.90	1.0
181-225	0.80	0.80	0.90	1.0
226-300	0.90	0.90	1.0	1.0
>300	1.0	1.0	1.0	1.0

^a P-Factors for closed outlet terraces also apply to terraces with underground outlets and to level terraces with open outlets.

^b The channel grade is measured on the 300 feet of terrace or the 1/3 of total terrace length closest to the outlet, whichever is less.

Possible phosphorus transport rating values for soil erosion are (Part A - Table 1):

0 for fields with no surface outlet (such as for karst areas in the Suwannee River watershed).

1 for fields with a calculated soil loss (A) of less than 5 tons/acre/year.

2 for fields with a calculated soil loss (A) of between 5 and 10 tons/acre/year.

4 for fields with a calculated soil loss (A) of between 10 and 15 tons/acre/year.

8 for fields with a calculated soil loss (A) of more than 15 tons/acre/year.

Soil Erosion Calculation Example

Situation: An area in the southeastern portion of Okeechobee county has the following conditions:

Soil: From soil survey the soil is map unit 11 (Immokalee fine sand). The soil was verified on-site as being Immokalee fine sand, on a 0.5 percent slope with a slope length of 100 feet.

Crop: The field is a Bermuda grass pasture planted every 4 years and grazed.

$$A = R * K * LS * C * P$$

R = 500 (for all of Okeechobee County)

K = 0.10 (from Appendix I - Table 13. Okeechobee County)

L S= 0.09 (from Table 4)

C = 0.067 (from Table 7)

P = 1.0 (from Table 9; field is not contour cropped, contour strip cropped, or terraced)

A = 500 * 0.10 * 0.09 * 0.067 * 1.0

A = 0.3 tons/acre/year

The resulting Soil Erosion value assigned to the Phosphorus Transport Rating - Part A (Table 1) would be 1 (<5 T/A). This is the most common result obtained in Okeechobee County.

Runoff Potential

Usage of the following runoff potential criteria is based on a minimum of 10 observations (soil borings) per spray field/application area unless the number of borings identifies the site as a problem area or a uniform area. At least one observation is to be made in each of the landforms present. Examples of landforms are flats, flatwoods, depressions, terraces, rises, knolls, hills, hillsides, sideslopes, toeslopes, footslopes, etc. If there is no surface outlet for the field in consideration, the rating is Very Low (**0**) for Runoff Potential.

The NRCS Hydrologic Soil Groups, slope, and the presence or absence of artificial drainage are used to evaluate runoff potentials.

Runoff Potential Rating Criteria - Part A (see Table 1)

Very Low (0):

Soils in Hydrologic Soil Group A with $\geq 75\%$ ground cover **and slopes of 8% or less.**

or:

any Hydrologic Soil Group with no surface outlet.

Low (1):

Soils in Hydrologic Soil Groups A with $< 75\%$ ground cover with surface outlet and A/D (with effective drainage depth of greater than 48") **and slopes of 8% or less** (Effective drainage is water control that is designed and maintained according to NRCS standards that will perform the desired water control.)

Medium (2):

Soils in Hydrologic Group A and A/D (with effective drainage depth of 37" to 48") **and slopes of more than 8%.**

or:

Soils in Hydrologic Groups B and A/D or B/D (with effective drainage depth of 37" to 48") **and slopes of 5% or less.**

High (4):

Soils in Hydrologic Group B and B/D (with effective drainage depth of 20" to 36") **and slopes of more than 5% up to and including 8%.**

or:

Soils in Hydrologic Groups C and A/D, B/D or C/D (with effective drainage depth of 20" to 36") **and slopes of 5% or less.**

Very High (8):

Soils in Hydrologic Group B and B/D (with effective drainage depth of 37" to 48") **and slopes of more than 8%.**

or:

Soils in Hydrologic Groups C and C/D (with effective drainage depth of 20" to 36") **and slopes of more than 5%.**

or:

Soils in Hydrologic Groups D and A/D, B/D, and C/D in undrained condition.

Runoff Potentials are presented in Appendix I for each County based on the above criteria and the definitions of the four hydrologic soil groups below. These are potentials to be used in conjunction with the soil surveys of the Okeechobee Watershed Area. Potentials presented are interpretations and are not factual data. As with all interpretations, **runoff potentials should be confirmed by on-site investigations. Slope and hydrologic group should be determined on-site.**

Group A: Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well-drained to excessively-rained sands or gravelly sands. These soils have a high rate of water transmission.

Group B: Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well-drained or well-drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C: Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D: Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink/swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

Artificial Drainage

Presence of artificial drainage can change the runoff potential of a soil. Drained Runoff Potentials in Table 13 in Appendix I have been assigned to those soils deemed drainable by NRCS. Drained Runoff Potentials presented are based on NRCS “Technical Release No. 55-Urban Hydrology for Small Watersheds, Amendment FL3” (Table 12).

Table 12. Reclassification of Runoff Potential and Hydrologic Group Based on Drainage.

Effective Drainage Depth (Inches) ^a	Drained Runoff Potential	Drained Hydrologic Group
Less than 20	Very High	D
20-36	High	C
37-48	Medium	B
Greater than 48	Low	A

^a Effective drainage is defined as having good surface drainage with a designed subsurface drainage system properly installed and maintained with a water removal rate of at least 0.5 inches/day. **Rarely have agricultural fields in Okeechobee County been effectively drained to a depth of more than 24 inches.**

Drained Runoff Potentials in Appendix I - Table 13 for each county are based on the maximum effective drainage depth expected for each soil. Actual effective drainage may be less than the maximum. For example, Immokalee (Table 13-Okeechobee County -map unit 11) has a drained runoff potential of Medium. This rating is based on a maximum effective drainage depth of 37 to 48 inches. If field conditions indicate a site had been effectively drained to a depth of only 24 inches, then the on-site runoff potential would be High (Table 12) and the resulting Phosphorus Transport Rating – Part A value for runoff would be 4 (Table 1).

Leaching Potential

Usage of the following leaching potential criteria is based on a minimum of 5 observations (e.g. soil borings) per 40 acres of application area unless the number of borings identify the site as a problem area or a uniform area. Ground penetrating radar (GPR) should be used for the assessment of all Karst areas. At least one observation is to be made in each landform present.

Presence or absence of a loamy/clayey layer and thicknesses of sandy layers, and presence or absence of coated sand are used to evaluate leaching potentials.

Leaching Potential Rating Criteria – Part A (see Table 1)

Very Low (0):

At least 80 percent of observations have a loamy or clayey layer at least 25 cm (10 inches) thick starting within 50 cm (20 inches). Typically, these soils are Typic Paleudults.

Low (1):

At least 80 percent of observations have a loamy or clayey layer at least 25 cm (10 inches) thick starting within 200 cm (80 inches). Typically, these soils are Arenic and Grossarenic Paleudults.

Medium (2):

At least 80 percent of observations have a loamy or clayey layer at least 25 cm (10 inches) thick starting at a depth below 200 cm (80 inches) but above seasonal high saturation **and** sand grains in the E and Bw horizons have coatings (chroma ≥ 3) to a depth of at least 100 cm (40 inches); or at least 80 percent of observations have no loamy or clayey layer at least 25 cm (10") thick, but have a layer at least 200 cm (80") thick with coated sand grains (chroma equal to or greater than 3). The entire 200 cm (80") layer must be above seasonal high saturation.

High (4):

At least 20 percent of observations have no loamy or clayey layer,(or the loamy or clayey layer is less than 25 cm (10 inches) thick) **and** the combined thickness of layers with coated sand grains (chroma ≥ 3 in the E, Bw, and C horizons and any chroma in the Bh horizons) is more than 50 cm (20 inches) and less than 200 cm (80 inches).

Very High (8):

At least 20 percent of observations have no loamy or clayey layer (or the layer is less than 25 cm (10 inches) thick) **and** the combined thickness of layers with coated sand grains (chroma ≥ 3 in the E, Bw, and C horizons and any chroma in the Bh horizons) is equal to or less than 50 cm (20 inches).

Leaching Potentials are presented in Appendix 1 - Table 13 based on the above criteria. These are potentials to be used in conjunction with the soil surveys of the Okeechobee Watershed Area. Potentials presented are interpretations, and are not factual data. As with all interpretations, **leaching potentials should be confirmed by on-site investigations.**

The rating of Medium Leaching Potential may be unique to Florida. This rating is based on deeper observation of soils that would normally be rated as having a High or Very High Leaching Potential. The rating of Medium Leaching Potential is given to soils with a significant loamy/clayey layer below the normal (2m or 80 inches) soil classification depth. Use of Ground Penetrating Radar (GPR) and/or geological investigations is needed to rate a site as having a Medium Leaching Potential and the depth to the loamy/clayey layer must be **above** the seasonal high saturation (water table).

Sinkholes occur where calcareous limestone below the land surface has been naturally dissolved by circulating ground water. A sinkhole forms when soil or weakened rock falls into underlying cavernous limestone. The sinkhole depth to width ratio tends to relate to soil slope stability-- typically the width is 5 times the depth. Okeechobee Watershed Area does not have areas considered to be high risk for sinkhole development. However, if sinkhole development is suspect, the GPR will be used to determine the leaching potential.

Phosphorus Runoff and Leaching Potentials Ratings for Florida Soil Survey Map Units

The runoff and leaching potentials (Appendix I - Table 13) were created by comparing estimated soil properties found in the County's soil survey with the above criteria. The potentials presented herein are interpretations, and not factual data. As with all interpretations based on information in a published soil survey or other sources of estimated soil properties, **phosphorus runoff and leaching potentials should be confirmed by on-site investigations.** However, a soil survey is an excellent place to initiate off-site investigation before making on-site determinations. For information on how to use a soil survey, see Circular 959 Soil Ratings for Crop Production and Water Quality Protection (Brown, et. al. 1991). However, note that phosphorus runoff and leaching potentials were derived from criteria that are different from the criteria used to derive the pesticide runoff and leaching potentials.

Potential to Reach Water Body

This parameter is used to address the potential for runoff to reach a water body. If there is no direct discharge from the edge of a field, the potential to affect a water body is considered to be “very low.” If the P concentration of the runoff can be attenuated by flow through a wetland, buffer strip or overland treatment area, the potential is considered “low.” If there is ditch drainage or direct discharge to a water body, the index value is increased to “medium.” When there is potential for direct discharge to a lake, sinkhole, or natural stream the potential for water quality degradation by P is enhanced and the index rating is increased to “high.”

Potential to Reach Water Body Rating Criteria (see Table 1)

Very Low (0):

No direct discharge from the edge of the field.

Low (1):

Discharge through wetlands, buffer area (refer to table below for buffer width), storm water detention, or overland treatment.

Medium (2):

No buffer, ditch drainage to or direct discharge to a water body.

High (4):

Direct discharges to a lake, sinkhole, or natural stream.

Non-Application Buffer Widths ¹		
Object, Site	Situation	Base Buffer Width from Object, Site (ft.)
Well, potable	Located up-slope of application site	150
Well, potable	Located down-slope of application site provided conditions warrant application	300
Waterbody, Stream ^{2/} , sinkhole or wetland	Good vegetation ^{3/} . Add 2 feet for each 1% slope for slopes up to 8%.	50 (+)
Waterbody, Stream ^{2/} , sinkhole or wetland	Poor vegetative cover or Predominant slope > 8% ^{3/}	100
Public Road – roadside ditch	Irrigated wastewater or solids applied with spreader	30

1/ Research has shown that forested or forest/grass buffers are more effective at removing phosphorus. Grass buffers are more effective at removing nitrogen. Every effort should be made to reduce phosphorus inputs at their sources. If phosphorus is managed responsibly on-site, buffers can store significant amounts of the excess; but if phosphorus is uncontrolled buffers can quickly become saturated and over whelmed. Even with their limits, buffers still perform a valuable service by displacing phosphorus-producing activities away from streams and regulating the flow of phosphorus. Taken in part from “A Review Of The Scientific Literature On Riparian Buffer Width, Extent And Vegetation”, Institute of Ecology, University of Georgia.

2/ Waterbody includes pond, lake, or open sinkhole. Open sinks include paleo sinks without a confining layer within 80 inches of the surface. Stream includes both perennial and intermittent streams and canals.

3/ Good vegetation refers to a well-managed, dense stand that is not overgrazed.

Phosphorus Transport Potential Due to Phosphorus Source Management - Part B (Table 2)

Phosphorus transport potential due to phosphorus source management is as follows:

- Fertility Index Value
- P Application Source and Rate
- Application Method
- Waste Water Application

Criteria

Fertility Index Value:

Existing soil P levels are included in the P Index and identified as the “fertility index”. The “fertility index” is defined as Mehlich-3 extractable P, of a 0-15 cm (0-6 inches) depth soil sample, in ppm (parts per million) multiplied by 2 to convert to pounds per acre. The 0.025 multiplication factor was selected to provide a value range similar to those used for other parameters in the P Index.

Obtain soil samples by taking 15 to 20 small cores (for areas up to 40 acres) at random over the entire area to a depth of about 6 inches. Place the 15 to 20 plugs in a container, mix them thoroughly, and send approximately one pint of the mixed sample to the UF/IFAS Extension Soil Testing Laboratory (ESTL) or other qualified laboratory for analysis.

P Application Source and Rate:

The multiplication factors for the application of P vary based on the source (fertilizer, manure, compost, biosolids, or waste water). Fertilizer, manure, and compost have the multiplier 0.05. For biosolids the multiplier is lower (0.015) because of evidence that the Fe and Al content of biosolids will decrease the P availability in biosolids-amended soils. In contrast, P in water from municipal and lagoon effluents is mostly in a soluble form and therefore the multiplier is higher (0.10).

Application Method:

The application method is not a consideration for sites that have No Surface Outlet or where solids are incorporated immediately after application or injected (value 0). For all other sites, effluent applied via irrigation are typically applied frequently (weekly, bi-weekly) and in small amounts or where solids are incorporated within one day of application; therefore, the potential for P loss is low (value 2). In contrast, solids (fertilizers, compost, biosolids, manures) surface-applied and not incorporated would have a higher potential for loss, particularly through surface runoff (value 6). Incorporated solids within 5 days of application have a medium potential for loss (value 4).

Waste Water Application Volume:

Excessive volumes of water may exacerbate movement of P via downward or lateral leaching, depending on the landscape. The 0.20 multiplication factor was selected to provide a value range similar to those used for other parameters in the P Index.

RESULTING P INDEX

The P Index is obtained by multiplying the site and transport characteristics totals – Part A (Table 1) by the phosphorus source totals – Part B (Table 2). The results are interpreted according to guidelines in Table 3.

On sites with a LOW or MEDIUM vulnerability rating, it is possible to use a nitrogen-based budget to determine application rates. On sites with a HIGH or VERY HIGH vulnerability rating, it is necessary to use a phosphorus-based budget to determine application rates.

Assessing the P Index Results

The numerical result of the P Index has no absolute value, but is immediately translated into a qualitative rating (LOW, MEDIUM, HIGH, or VERY HIGH). For each qualitative rating a description is given for the level of concern that each specifically assessed field has for P loss potential (Table 3). Some general guidance is given for each qualitative level as to the intensity and type of remedial action or mitigation that would be necessary to reduce P loss risk.

Conservation Planning Notes

Since output from the P Index includes information that is specific to each of the site and transport characteristics – Part A (Table 1) and phosphorus source management – Part B (Table 2), the conservation planner can identify which characteristics/management have the greatest influence in determining the final vulnerability rating and may be targeted for remedial action. Table 14 may be used to record notes to explain, clarify, and/or define site characteristics and source management used to evaluate a site. Each factor can be revisited and planning changes made, thereby changing the resulting P Index. For example, terraces can be installed, thereby lowering soil erosion and the final P Index. Similarly, the P Index can be lowered by reducing the planned P application rate.

Table 14. Conservation Planning Notes.

Client Name:	County:	Date:
Planner:	Field(s):	Crop:
Site and Transport Characteristics	Remarks	
Soil Erosion		
Runoff Potential		
Leaching Potential		
Potential to Reach Water Body		
Phosphorus Source Management		
Fertility Index Value		
P Application Source and Rate		
P Application Method		
Waste Water Application		

GLOSSARY (as used in the P Index the following definitions apply)

No Surface Outlet – The combination of slope and permeability of the application site that will not discharge surface flow from that site in a 2 year – 24 hour rainfall event.

(This level of evaluating runoff is not intended to require calculation for the rainfall events but is intended to evaluate those sites that do not have external surface flows during most years. Where these sites occur, additional comments may need to be recorded on the back of form FL-CPA-41)

Compost – animal wastes and plant debris that has gone through the composting process.

Biosolids – Residuals, domestic wastewater residuals and/or septage as defined in Chapter 62-640 Florida Administrative Code. Biosolids include co-compost with a minimum of 50% biosolids.

Landform - Any physical, recognizable form or feature of the earth's surface, having a characteristic shape and produced by natural causes.

Examples of individual landforms and their definitions are:

Karst - Topography with sinkholes, caves, and underground drainage that is formed in limestone, gypsum, or other rocks by dissolution, and that is characterized by sinkholes, caves, and underground drainage.

Knoll - A small, low, rounded hill rising above adjacent landforms.

Subsurface Drainage – Lowering of the water table in order to improve vegetative growth, remove surface runoff from wet areas, or relieve artesian pressure. Subsurface drainage can be achieved by either using drainage tile or drainage ditches, typically spaced at regular intervals.

REFERENCES

Belz, D.J., L.J. Carter, D.A. Dearstyne, and J.D. Overing. 1990. Soil Survey of Hendry County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Brown, R.B., A.G. Hornsby, and G.W. Hurt. 1991. Soil ratings for crop production and water quality protection. Circular 959. Florida Cooperative Extension Service, University of Florida, Gainesville, FL.

Carter, L., D. Lewis, and J. Vega. 2000. Soil Survey of Glades County, Florida. USDA/NRCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services. http://soils.usda.gov/survey/online_surveys/florida/

Carter, L.J., D.L. Lewis, L. Crockett, and J.A. Vega. 1989. Soil Survey of Highlands County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Doolittle, J.A. and G. Schellentrager. 1989. Soil Survey of Orange County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Florida Ecological Sciences Staff. 1999. Florida Agronomy Field Handbook, Chapter 6. USDA, NRCS, Gainesville, FL.

Florida Phosphorus Index Work Group. 2000. The Florida phosphorus index. <http://efotg.nrcs.usda.gov/treemenuFS.aspx?Fips=12001&MenuName=menuFL.zip> (The Florida Phosphorus Index sheets are located in Section IV of the Table of Contents under C.Tools.)

Ford, R.D., J.M. Robbins, Jr., J.T. Werner, W.D. Cowherd, C.N. Gordon, W.B. Warmack, M.K. Brown, K.W. Monroe, W.G. George, T.E. Sanders, and P.M. Basch. 1990. Soil Survey of Polk County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Furman, A.L., H.O. White, O.E. Cruz, W.E. Russell, and B.P. Thomas. 1975. Soil Survey of Lake County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Henderson, W.G. Jr. 1984. Soil Survey of Charlotte County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Lewis, D.L., K.J. Liudahl, C.V. Noble, and L.J. Carter. 2001. Soil Survey of Okeechobee County, Florida. USDA/NRCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.
http://soils.usda.gov/survey/online_surveys/florida/okeechobee/fl_okeechobee.pdf

MuCullum, S.H. and O.E. Cruz, Sr. 1981. Soil Survey of Martin County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

MuCullum, S.H., O.E. Cruz, Sr., L.T. Stem, W.H. Wittstruck, R.D. Ford, and F.C. Watts. 1978. Soil Survey of Palm Beach County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Readle, E.L. 1979. Soil Survey of Osceola County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil and Water Science Department; and Florida Department of Agriculture and Consumer Services.

Watts, F.C., and D.L. Stankey. 1980. Soil Survey of St. Lucie County, Florida. USDA/SCS in cooperation with the University of Florida, Institute of Food and Agricultural Sciences, Agricultural Experimental Stations and Soil Science Department; and Florida Department of Agriculture and Consumer Services.

APPENDIX I.

Runoff, Leaching Potentials and K-Factors for County Soils.

CHARLOTTE COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Charlotte County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
002	1	Canaveral	High	High ^b		0.10
004	1	Canaveral	High	High ^b		0.10
004	2	Urban Land	Very High	Variable		No Value
005	1	Captiva	Very High	Very High	Low ^c	0.10
006	1	Hallandale	Very High	Very High	Low ^c	0.10
007	1	Matlacha	High	Very High		0.15
007	2	Urban Land	Very High	Variable		No Value
008	1	Hallandale	Very High	Very High		0.10
009	1	EauGallie	Very High	High ^d	Medium	0.10
010	1	Pompano	Very High	Very High	Low ^c	0.10
011	1	Myakka	Very High	High	Medium	0.10
012	1	Felda	Very High	Low	Medium	0.10
013	1	Boca	Very High	Very High ^d	Medium	0.10
014	1	Valkaria	Very High	Very High	Low ^c	0.10
015	1	Estero	Very High	Very High		0.02
016	1	Peckish	Very High	Very High		0.10
017	1	Daytona	Medium	Very High		0.10
018	1	Matlacha	High	Very High		0.15
019	1	Gator	Very High	Very High	Medium	0.02
020	1	Terra Ceia	Very High	Very High	Medium	0.02
022	1	Beaches	Very High	Very High		0.05
023	1	Wulfert	Very High	Very High		0.02
024	1	Kesson	Very High	Very High		0.10
025	1	St. Augustine	High	Very High ^e		0.10
025	2	Urban Land	Very High	Variable		No Value
026	1	Pineda	Very High	Low	High	0.10
027	1	Pompano	Very High	Very High	Low ^c	0.10
028	1	Immokalee	Very High	High	Medium	0.10
029	1	Punta	Very High	High	Medium	0.10
033	1	Oldsmar	Very High	Low	Medium	0.10
034	1	Malabar	Very High	Low	Medium	0.10
035	1	Wabasso	Very High	Low	High	0.10
036	1	Immokalee	Very High	High	Medium	0.10
036	2	Urban Land	Very High	Variable		No Value

Table 13 (cont.). Runoff, Leaching Potentials and K-Factors for Charlotte County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
037	1	Satellite	High	Very High		0.10
038	1	Isles	Very High	Very High ^d	Medium	0.10
039	1	Isles	Very High	Very High ^d	Medium	0.10
040	1	Anclote	Very High	Very High	Low ^c	0.10
041	1	Valkaria	Very High	Very High	Low ^c	0.10
042	1	Wabasso	Very High	Low	High	0.10
043	1	Smyrna	Very High	High	Medium	0.10
044	1	Malabar	Very High	Low	Medium	0.10
045	1	Copeland	Very High	Very High ^d	High	0.10
048	1	St. Augustine	High	Very High ^e		0.10
049	1	Felda	Very High	Low	Medium	0.10
050	1	Oldsmar	Very High	Low	Medium	0.10
051	1	Floridana	Very High	Low	High	0.10
053	1	Myakka	Very High	High	Medium	0.10
055	1	Cocoa	Low ^c	High ^d		0.10
056	1	Isles	Very High	Very High ^d		0.02
057	1	Boca	Very High	Very High ^d		0.10
059	1	Urban Land	Very High	Variable		No Value
061	1	Orsino	Low ^c	High ^b		0.10
062	1	Winder	Very High	Very Low	High	0.10
063	1	Malabar	Very High	Low	Medium	0.10
064	1	Hallandale	Very High	Very High	Low ^c	0.10
064	2	Urban Land	Very High	Variable		No Value
066	1	Caloosa	High	Low		0.10
067	1	Smyrna	Very High	High	Medium	0.10
067	2	Urban Land	Very High	Variable		No Value
069	1	Matlacha	High	Very High		0.15
070	1	Heights	Very High	Very High ^d	Medium	0.10
072	1	Bradenton	Very High	Very Low	High	0.10
073	1	Pineda	Very High	Low	High	0.10
074	1	Boca	Very High	Very High ^d	Medium	0.10
075	1	Hallandale	Very High	Very High	Low ^c	0.10
076	1	Electra	High	Low		0.10
077	1	Pineda	Very High	Low	High	0.10
078	1	Chobee	Very High	Very Low	High	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^c Rate Very Low where percent ground cover is greater than 75%.

^d Rate Low if thickness of loamy/clayey layers is more than 10 inches.

^e Rate High if combined thickness of layers with chroma 3 or more and Bh horizons is more than 20 inches.

GLADES COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Glades County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
002	1	Hallandale	Very High	Very High	Low ^b	0.10
004	1	Valkaria	Very High	Very High	Low ^b	0.10
005	1	Smyrna	Very High	High ^c	Medium	0.10
006	1	Malabar	Very High	Low	Medium	0.10
007	1	Pople	Very High	Very High ^d	High	0.10
008	1	Gator	Very High	Low	Medium	0.02
009	1	Sanibel	Very High	Very High	Low ^b	0.10
010	1	Felda	Very High	Low	Medium	0.10
011	1	Tequesta	Very High	Very High ^d	Medium	0.02
012	1	Chobee	Very High	Very Low	High	0.20
013	1	Boca	Very High	Very High ^d	High	0.10
014	1	Basinger	Very High	Very High	Low ^b	0.10
015	1	Pineda	Very High	High ^d	High	0.10
016	1	Floridana	Very High	Low	High	0.05
017	1	Okeelanta	Very High	Very High	Medium	0.02
019	1	Terra Ceia	Very High	Very High	Medium	0.02
020	1	EauGallie	Very High	High ^d	Medium	0.10
022	1	Astor	Very High	Very High	Low ^b	0.10
023	1	Oldsmar	Very High	Low	Medium	0.10
024	1	Hallandale	Very High	Very High	Low ^b	0.10
024	2	Pople	Very High	Very High ^d	High	0.10
026	1	Immokalee	Very High	High	Medium	0.10
027	1	Ft. Drum	Very High	Very High ^d	Medium	0.10
028	1	Pomello	High	High ^c		0.10
029	1	Myakka	Very High	High	Medium	0.10
032	1	Floridana	Very High	Low		0.05
032	2	Astor	Very High	Very High		0.10
032	3	Felda	Very High	Low		0.10
034	1	Basinger	Very High	Very High	Low ^b	0.10

GLADES COUNTY (cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Glades County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
035	1	Arents	Very High	Variable		Variable ^e
036	1	Malabar	Very High	Low	Medium	0.10
037	1	Lauderhill	Very High	Very High	Medium	0.02
038	1	Pahokee	Very High	Very High	Medium	0.02
040	1	Plantation	Very High	Very High	Low ^b	0.02
041	1	Dania	Very High	Very High	Medium	0.02
042	1	Okeelanta	Very High	Very High	Medium	0.02
042	2	Dania	Very High	Very High	Medium	0.02
043	1	Sanibel	Very High	Very High	Low ^b	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^d Rate Low if thickness of loamy/clayey layers is more than 10 inches.

^e The following K-Factors should be used for the following on-site surface textures: sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, and clay = 0.37

HENDRY COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Hendry County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
001	1	Boca	Very High	Very High ^b	Medium	0.10
002	1	Pineda	Very High	Low	High	0.10
004	1	Oldsmar	Very High	Very High ^b	Medium	0.10
006	1	Wabasso	Very High	Low	High	0.10
007	1	Immokalee	Very High	High	Medium	0.10
008	1	Malabar	Very High	Low	Medium	0.10
009	1	Riviera	Very High	Very High ^b	High	0.10
010	1	Pineda	Very High	Low	High	0.10
012	1	Winder	Very High	Very Low	High	0.10
013	1	Gentry	Very High	Low	Medium	0.10
014	1	Wabasso	Very High	Low	High	0.10
015	1	Myakka	Very High	High	Medium	0.10
017	1	Basinger	Very High	Very High	Low ^c	0.10
018	1	Pompano	Very High	Very High	Low ^c	0.10
019	1	Gator	Very High	Low	Medium	0.02
020	1	Okeelanta	Very High	Very High	Medium	0.02
021	1	Holopaw	Very High	Very High ^b	Medium	0.02
022	1	Valkaria	Very High	Very High	Low ^c	0.10
023	1	Hallandale	Very High	Very High	Low ^c	0.10
024	1	Pomello	High	High ^d		0.10
026	1	Holopaw	Very High	Very High ^b	Medium	0.10
027	1	Riviera	Very High	Very High ^b	High	0.10
028	1	Boca	Very High	Very High ^b	Medium	0.10
029	1	Oldsmar	Very High	Very High ^b	Medium	0.10
032	1	Riviera	Very High	Very High ^b	High	0.10
033	1	Holopaw	Very High	Very High ^b	Medium	0.10
034	1	Chobee	Very High	Very Low	High	0.15
037	1	Tuscawilla	Very High	Very Low	High	0.10
039	1	Udifluvents	High	Variable		0.10
042	1	Riviera	Very High	Very High ^b	High	0.10
044	1	Jupiter	Very High	Very High	Low ^c	0.10
045	1	Pahokee	Very High	Very High	Medium	0.02
047	1	Udorthents	Low ^a	Variable		0.10
050	1	Delray	Very High	Very High ^b	Medium	0.10
051	1	Malabar	Very High	Low	Medium	0.10
053	1	Adamsville	High	High ^d		0.10
056	1	Terra Ceia	Very High	Very High	Medium	0.02
057	1	Chobee	Very High	Very Low	High	0.15

HENDRY COUNTY (cont.).

Table 13. Runoff, Leaching Potentials and K-Factors for Hendry County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
058	1	Oldsmar	Very High	Very High ^b	Medium	0.10
059	1	Winder	Very High	Very Low	High	0.10
060	1	Myakka	Very High	High	Medium	0.10
061	1	Malabar	Very High	Low	Medium	0.10
062	1	Pineda	Very High	Low	High	0.10
063	1	Jupiter	Very High	Very High	Low ^c	0.10
063	2	Ochopee	Very High	Very High ^b	Medium	0.17
063	3	Rock Outcrop	Very High	Very High		0.02
064	1	Hallandale	Very High	Very High	Low ^c	0.10
065	1	Plantation	Very High	Very High	Low ^c	0.02
066	1	Margate	Very High	Very High	Low ^c	0.10
067	1	Lauderhill	Very High	Very High	Medium	0.02
068	1	Dania	Very High	Very High	Medium	0.02
069	1	Denaud	Very High	Very High ^b	Medium	0.10
069	2	Gator	Very High	Very High ^b	Medium	0.02
070	1	Denaud	Very High	Very High ^b	Medium	0.10
073	1	Adamsville Variant	High	High ^d		0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Low if thickness of loamy/clayey layers is more than 10inches.

^c Rate Low where percent ground cover is greater than 75%.

^d Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

HIGHLANDS COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Highlands County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
001	1	Paola	Low ^b	High ^c		0.10
002	1	St. Lucie	Low ^b	Very High		0.10
003	1	Basinger	Very High	Very High	Low ^b	0.10
004	1	Duette	Low ^b	High ^c		0.10
005	1	Daytona	Low ^b	High ^c		0.10
006	1	Tavares	Low ^b	High		0.10
007	1	Placid	Very High	Very High	Low ^b	0.10
008	1	Immokalee	Very High	High	Medium	0.10
009	1	Astatula	Low ^b	High		0.10
010	1	Myakka	Very High	High	Medium	0.10
011	1	Orsino	Low ^b	High ^c		0.10
012	1	Basinger	Very High	Very High	Low ^b	0.10
013	1	Felda	Very High	Low	Medium	0.10
014	1	Satellite	High	Very High		0.10
015	1	Bradenton	Very High	Very Low	High	0.15
016	1	Valkaria	Very High	Very High	Low ^b	0.10
017	1	Malabar	Very High	Low	Medium	0.10
018	1	Kaliga	Very High	Low	Medium	0.02
019	1	Hicoria	Very High	Low	Medium	0.10
020	1	Samsula	Very High	Very High	Medium	0.02
021	1	Hontoon	Very High	Very High	Medium	0.02
022	1	Brighton	Very High	Very High	Medium	0.02
024	1	Pineda	Very High	Low	High	0.10
025	1	Chobee	Very High	Very Low	High	0.15
026	1	Tequesta	Very High	Very Low	High	0.10
028	1	Archbold	Low ^b	Very High		0.10
029	1	Pomona	Very High	Low	Medium	0.10
030	1	Oldsmar	Very High	Low	Medium	0.10
031	1	Felda	Very High	Low	Medium	0.10
032	1	Arents	Very High	Variable		0.10
033	1	Basinger	Very High	Very High	Low ^b	0.10
033	2	St. Johns	Very High	High	Medium	0.10
033	3	Placid	Very High	Very High	Low ^b	0.10
034	1	Tavares	Low ^b	High		0.10
034	2	Basinger	Very High	Very High	Low ^b	0.10

HIGHLANDS COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Highlands County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
034	3	Sanibel	Very High	Very High	Low ^b	0.10
035	1	Sanibel	Very High	Very High	Low ^b	0.10
036	1	Pomello	High	High		0.10
037	1	Malabar	Very High	Low	Medium	0.10
038	1	EauGallie	Very High	Low	Medium	0.10
039	1	Smyrna	Very High	High	Medium	0.10
040	1	Arents	Variable	Variable		Variable ^e
041	1	Anclote	Very High	Very High		0.10
041	2	Basinger	Very High	Very High		0.10
042	1	Astatula	Low ^b	High		0.10
042	2	Urban Land	Very High	Variable		No Value
043	1	Urban Land	Very High	Variable		No Value
044	1	Satellite	High	Very High		0.10
044	2	Basinger	Very High	Very High	Low ^b	0.10
044	3	Urban Land	Very High	Variable		No value
045	1	Paola	Low ^b	High		0.10
045	2	Basinger	Very High	Very High	Low ^b	0.10
046	1	Kaliga	Very High	Low		0.02

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^d Rate Low if thickness of loamy/clayey layers is more than 10inches.

^e K-Factors to be used: muck = 0.02, mucky sand = 0.05, sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, and clay = 0.37

LAKE COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Lake County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
AbB	1	Albany	High	Low		0.10
AbD	1	Albany	Very High	Low		0.10
An	1	Anclote	Very High	Very High	Low ^b	0.10
Am	1	Anclote	Very High	Very High	Low ^b	0.10
Am	2	Myakka	Very High	High	Medium	0.10
ApB	1	Apopka	Low ^b	Low		0.10
ApD	1	Apopka	Medium ^c	Low		0.10
AsB	1	Astatula	Low ^b	High		0.10
AtB	1	Astatula	Low ^b	High		0.10
AtD	1	Astatula	Medium ^c	High		0.10
AtF	1	Astatula	Medium	High		0.10
Br	1	Brighton	Very High	Very High	Medium	0.10
Ca	1	Cassia	High	High ^d		0.10
Em	1	Emeralda	Very High	Very Low	Very High	0.10
Eu	1	Eureka	Very High	Very Low	Very High	0.15
Fd	1	Felda	Very High	Low	Medium	0.10
Fe	1	Fellowship	Very High	Very Low	Very High	0.20
Fm	1	Fill Land	Variable	Variable		0.28
Ib	1	Iberia	Very High	Very Low	Very High	0.32
Im	1	Iberia	Very High	Very Low		0.28
Im	2	Manatee	Very High	Very Low		0.10
Is	1	Immokalee	Very High	High	Medium	0.10
LaB	1	Lake	Low ^b	High		0.10
LaD	1	Lake	Medium ^c	High		0.10
LaE	1	Lake	Medium	High		0.10
Ma	1	Manatee	Very High	Very Low	High	0.10
Md	1	Montverde	Very High	Very High		0.02
Mk	1	Myakka	Very High	High	Medium	0.10
MpC	1	Myakka	Very High	High	High ^e	0.10
Oc	1	Ocilla	High	Low		0.10
Oe	1	Ocoee	Very High	Very High		0.02
Oh	1	Oklawaha	Very High	Very High		0.02
On	1	Ona	Very High	High	Medium	0.10
Or	1	Orlando	Low ^b	High		0.10

LAKE COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Lake County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
PaB	1	Paola	Low ^a	High ^d		0.10
PaD	1	Paola	Medium ^c	High ^d		0.10
Pd	1	Pelham	Very High	Low	Medium	0.10
Pe	1	Placid	Very High	Very High	Low ^b	0.10
Pg	1	Placid	High	Very High		0.10
PmA	1	Placid	Very High	Very High	Low ^d	0.10
Pn	1	Pomello	High	High ^d		0.10
Po	1	Pompano	Very High	Very High	Low ^b	0.10
Sc	1	St. Lucie	Low ^b	Very High		0.10
Sw	1	Swamp	Very High	Variable		0.10
Ta	1	Tavares	Low ^b	High		0.10
Te	1	Tavares Variant	Low ^b	High ^d		0.10
Va	1	Vaocluse	Medium	Very Low		0.15
Wa	1	Wabasso	Very High	Low	High	0.10
Wc	1	Wauchula	Very High		High	0.10
<p>^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.</p> <p>^b Rate Very Low where percent ground cover is greater than 75%.</p> <p>^c Rate Low or Very Low (where percent ground cover is greater than 75%) where slope is greater than 8 percent.</p> <p>^d Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.</p> <p>^e Rate Medium where slope is 5 percent or less.</p>						

MARTIN COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Martin County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
002	1	Lawnwood	Very High	High	High	0.10
003	1	Lawnwood	Very High	High	High	0.10
004	1	Waveland	Very High	High	High	0.10
005	1	Waveland	Very High	High	High	0.10
006	1	Paola	Low ^b	Very High ^c		0.10
007	1	St. Lucie	Low ^b	Very High		0.10
008	1	Palm Beach	Low ^b	Very High ^c		0.10
009	1	Pomello	High	High ^d		0.10
010	1	Basinger	Very High	Very High	Low ^b	0.10
012	1	St. Johns Variant	Very High	High ^d	Medium	0.10
013	1	Placid	Very High	Very High	Low ^b	0.10
014	1	Satellite Variant	High	Very High		0.15
015	1	Electra	High	Low		0.10
016	1	Oldsmar	Very High	Low	Medium	0.10
017	1	Wabasso	Very High	Low	High	0.10
019	1	Winder	Very High	Very Low	High	0.10
020	1	Riviera	Very High	Low	High	0.10
021	1	Pineda	Very High	High ^e	High	0.10
022	1	Okeelanta	Very High	Very High	Medium	0.02
023	1	Urban Land	Very High	Variable		No Value
024	1	Orsino	Low ^b	High ^d		0.10
025	1	Beaches	Very High	Very High		0.05
026	1	Pompano	Very High	Very High	Low ^b	0.10
027	1	Arents	Medium	Variable		0.10
028	1	Canaveral	High	High ^d		0.10
030	1	Bessie	Very High	Very High		0.02
031	1	Cocoa Variant	Low ^b	High ^d		0.15
032	1	Udorthents	Variable	Variable		0.10
033	1	Paola	Low ^b	Very High ^c		0.10
033	2	Urban Land	Very High	Variable		No value
034	1	St. Lucie	Low ^b	Very High		0.10
034	2	Urban Land	Very High	Variable		No Value
035	1	Salerno	Very High	High	Medium	0.10
038	1	Floridana	Very High	Low	High	0.10
039	1	Quartzipsamments	Variable	Variable		0.10
040	1	Sanibel	Very High	Very High	Low ^b	0.10
041	1	Jonathan	Medium	High		0.10
042	1	Hallandale	Very High	Very High	Low ^b	0.10
044	1	Boca	Very High	Very High ^e	Medium	0.10
045	1	Hilolo	Very High	Very Low	High	0.10

MARTIN COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Martin County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
047	1	Pinellas	Very High	Low	Medium	0.10
048	1	Jupiter	Very High	Very High	Low ^b	0.10
049	1	Riviera	Very High	Low	High	0.10
050	1	Okeelanta Variant	Very High	Very High	Medium	0.02
051	1	Pompano	Very High	Very High	Low ^b	0.10
052	1	Malabar	Very High	Low	Medium	0.10
053	1	Arents	Low ^b	Variable		0.10
054	1	Oldsmar	Very High	Very High	Medium	0.10
055	1	Basinger	Very High	Very High	Low ^b	0.10
056	1	Wabasso	Very High	Low	High	0.10
057	1	Chobee	Very High	Very Low	High	0.15
058	1	Gator	Very High	Low	Medium	0.02
060	1	Tequesta Variant	Very High	Very High ^e	Medium	0.02
060	2	Tequesta	Very High	Very High ^e	Medium	0.02
061	1	Hobe	Low ^b	Very High ^f		0.10
062	1	Nettles	Very High	High	High	0.10
063	1	Nettles	Very High	High	High	0.10
064	1	EauGallie	Very High	Low	Medium	0.10
065	1	Tuscawilla	Very High	Very Low	High	0.10
066	1	Holopaw	Very High	Very High ^e	Medium	0.10
067	1	Arents	Very High	Variable		0.10
068	1	Pits	Variable	Variable		0.10
069	1	Hontoon	Very High	Very High	Medium	0.10
070	1	Canova	Very High	Low	High	0.02
072	1	Adamsville Variant	High	Very High ^c		0.10
073	1	Samsula	Very High	Very High	Medium	0.02
074	1	Torry	Very High	Very High	Medium	0.02
075	1	Ft. Drum	Very High	Very High ^e	Medium	0.10
076	1	Valkaria	Very High	Very High	Low ^b	0.10
077	1	St, Lucie	Low ^b	Very High		0.10
078	1	Pomello Variant	High	High ^d		0.10
079	1	Terra Ceia Variant	Very High	Very High		0.02
086	1	Paola	Medium	Very High ^c		0.10
086	2	Astatula	Medium	High		0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate High if combined thickness of layers with chroma 3 or more and Bh horizons is more than 20 inches.

^d Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^e Rate Low if thickness of loamy/clayey layers is more than 10 inches.

^f Rate Very Low if thickness of loamy/clayey layers is more than 10 inches.

OKEECHOBEE COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Okeechobee County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
002	1	Basinger	Very High	Very High	Low ^b	0.10
003	1	Basinger	Very High	Very High	Low ^b	0.10
003	2	Placid	Very High	Very High	Low ^b	0.10
004	1	Bradenton	Very High	Very Low	High	0.15
005	1	Valkaria	Very High	Very High	Low ^b	0.10
006	1	Manatee	Very High	Very Low	High	0.10
007	1	Floridana	Very High	Low	High	0.10
007	2	Riviera	Very High	Low	High	0.10
007	3	Placid	Very High	Very High	Low ^b	0.10
008	1	Pineda	Very High	Low	High	0.10
009	1	Riviera	Very High	Low	High	0.10
010	1	Ft. Drum	Very High	Low	Medium	0.10
011	1	Immokalee	Very High	High	Medium	0.10
012	1	Udorthents	Very High	Very Low		0.15
013	1	Manatee	Very High	Very Low		0.10
013	2	Floridana	Very High	Low		0.10
013	3	Tequesta	Very High	Low		0.02
014	1	Myakka	Very High	High	Medium	0.10
015	1	Okeelanta	Very High	Very High	Medium	0.02
017	1	Orsino	Low ^b	High ^c		0.10
018	1	Parkwood	Very High	Very High ^d	High	0.10
019	1	Floridana	Very High	Low		0.10
019	2	Placid	Very High	Very High		0.10
019	3	Okeelanta	Very High	Very High		0.02
020	1	Pomello	High	High		0.10
021	1	Adamsville	High	High		0.10
023	1	St. Johns	Very High	High	Medium	0.10
024	1	Terra Ceia	Very High	Very High	Medium	0.02
025	1	Wabasso	Very High	Low	High	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^d Rate Very Low if thickness of loamy/clayey layers is more than 10 inches.

ORANGE COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Orange County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
001	1	Arents	Variable	Variable		0.10
002	1	Archbold	Low ^b	Very High		0.10
003	1	Basinger	Very High	Very High	Low ^b	0.10
004	1	Candler	Low ^b	High		0.10
005	1	Candler	Medium ^c	High		0.10
006	1	Candler	Medium ^c	High		0.10
006	2	Apopka	Medium ^c	Low		0.10
007	1	Candler	Low ^b	High		0.10
007	2	Urban Land	Very High	Variable		No Value
008	1	Candler	Medium ^c	High		0.10
008	2	Urban Land	Very High	Variable		No value
009	1	Canova	Very High	Low	High	0.02
010	1	Chobee	Very High	Very Low		0.15
011	1	Floridana	Very High	Low		0.10
011	2	Chobee	Very High	Very Low		0.15
012	1	Holopaw	Very High	Very High ^d		0.10
013	1	Felda	Very High	Low	Medium	0.10
014	1	Felda	Very High	Low	Medium	0.10
015	1	Felda	Very High	Low		0.10
016	1	Floridana	Very High	Low		0.10
017	1	Floridana	Very High	Low	Medium	0.10
018	1	Gator	Very High	Low	Medium	0.02
019	1	Hontoon	Very High	Very High	Medium	0.02
020	1	Immokalee	Very High	High	Medium	0.10
021	1	Lake	Low ^b	High		0.10
022	1	Lochloosa	High	Low		0.10
023	1	Malabar	Very High	Low	Medium	0.10
024	1	Millhopper	Low ^b	Low		0.10
024	2	Urban Land	Very High	Variable		No Value
025	1	Okeelanta	Very High	Very High	Medium	0.02
026	1	Ona	Very High	High	Medium	0.10
027	1	Ona	Very High	High	Medium	0.10
027	2	Urban Land	Very High	Variable		No Value
028	1	Florahome	Low ^b	High		0.10
029	1	Florahome	Low ^b	High		0.10
029	2	Urban Land	Very High	Variable		No Value
030	1	Pineda	Very High	Low	High	0.10
031	1	Pineda	Very High	Low		0.10
032	1	Pinellas	Very High	High	Medium	0.10

ORANGE COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Orange County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
033	1	Pits	Variable	Variable		Variable ^e
034	1	Pomello	High	High		0.10
035	1	Pomello	High	High		0.10
035	2	Urban Land	Very High	Variable		No Value
036	1	Pompano	Very High	Very High	Low ^b	0.10
037	1	St. Johns	Very High	High	Medium	0.10
038	1	St Lucie	High	Very High		0.10
039	1	St. Lucie	High	Very High		0.10
039	2	Urban Land	Very High	Variable		No Value
040	1	Samsula	Very High	Very High	Medium	0.02
041	1	Samsula	Very High	Very High	Medium	0.02
041	2	Hontoon	Very High	Very High	Medium	0.02
041	3	Basinger	Very High	Very High	Low ^b	0.10
042	1	Sanibel	Very High	Very High	Low ^b	0.10
043	1	Seffner	High	High		0.10
044	1	Smyrna	Very High	High	Medium	0.10
045	1	Smyrna	Very High	High	Medium	0.10
045	2	Urban Land	Very High	Variable		No Value
046	1	Tavares	Low ^b	High		0.10
047	1	Tavares	Low ^b	High		0.10
047	2	Millhopper	Low ^b	Low		0.10
048	1	Tavares	Low ^b	High		0.10
048	2	Urban Land	Very High	Variable		No Value
049	1	Terra Ceia	Very High	Very High	Medium	0.02
050	1	Urban Land	Very High	Variable		No Value
051	1	Wabasso	Very High	Low	High	0.10
052	1	Wabasso	Very High	Low	High	0.10
052	2	Urban Land	Very High	Variable		No Value
053	1	Wauberg	Very High	Low	High	0.15
054	1	Zolfo	High	High		0.10
055	1	Zolfo	High	High		0.10
055	2	Urban Land	Very High	Variable		No Value

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Low or Very Low (where percent ground cover is greater than 75%) if slope is less than 8 percent.

^d Rate Very Low if thickness of loamy/clayey layers is more than 10 inches.

^e The following K-factors should be used for the following on-site surface textures: sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, and clay = 0.37.

OSCEOLA COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Osceola County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
001	1	Adamsville	High	High		0.10
002	1	Adamsville Variant	High	High		0.10
003	1	Ankona	Very High	Low	Medium	0.10
004	1	Arents	High	Variable		0.10
005	1	Basinger	Very High	Very High	Low ^b	0.10
006	1	Basinger	Very High	Very High	Low ^b	0.10
007	1	Candler	Low ^b	High		0.10
008	1	Candler	Medium ^c	High		0.10
009	1	Cassia	High	High ^d		0.10
010	1	Delray	Very High	Low	Medium	0.10
011	1	EauGallie	Very High	Low	Medium	0.10
012	1	Floridana	Very High	Low	High	0.10
013	1	Gentry	Very High	Low		0.10
014	1	Holopaw	Very High	Very High ^e	Medium	0.10
015	1	Hontoon	ery High	Very High	Medium	0.02
016	1	Immokalee	Very High	High	Medium	0.10
017	1	Kaliga	Very High	Low	Medium	0.02
017	2	Terra Ceia	Very High	Very High	Medium	0.02
018	1	Lokosee	Very High	Very High ^f	Medium	0.10
019	1	Malabar	Very High	Low	Medium	0.10
020	1	Malabar	Very High	Low	Medium	0.10
021	1	Malabar	Very High	Low	Medium	0.10
021	2	Pineda	Very High	High ^e	High	0.10
022	1	Myakka	Very High	High	Medium	0.10
023	1	Myakka	Very High	High	Medium	0.10
023	2	Urban Land	Very High	Variable		Variable ^g
024	1	Narcoossee	High	High ^d		0.10
025	1	Nittaw	Very High	Very Low		0.02
026	1	Oldsmar	Very High	Low	Medium	0.10
027	1	Ona	Very High	High	Medium	0.10
028	1	Paola	Low ^b	Very High ^f		0.10
029	1	Parkwood	Very High	Very High ^h	High	0.10
030	1	Pineda	Very High	Low	High	0.10
032	1	Placid	Very High	Very High	Low ^b	0.10

OSCEOLA COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Osceola County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
033	1	Placid Variant	High	Very High		0.10
034	1	Pomello	High	High		0.10
035	1	Pomona	Very High	Low	Medium	0.10
036	1	Pompano	Very High	Very High	Low ^b	0.10
037	1	Pompano	Very High	Very High	Low ^b	0.10
038	1	Riviera	Very High	Low	High	0.10
039	1	Riviera	Very High	Low	High	0.10
040	1	Samsula	Very High	Very High	Medium	0.02
041	1	Satellite	High	Very High		0.10
042	1	Smyrna	Very High	High	Medium	0.10
043	1	St. Lucie	Low ^b	Very High		0.10
044	1	Tavares	Low ^b	High		0.10
045	1	Vero	Very High	Low	High	0.10
046	1	Wauchula	Very High	Low	High	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Low or Very Low (where percent ground cover is greater than 75%) if slope is less than 8 percent.

^d Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^e Rate Low if thickness of loamy/clayey layers is more than 10 inches thick.

^f Rate High if combined thickness of layers with chroma 3 or more and Bh horizons is more than 20 inches.

^g The following K-factors should be used for the following on-site surface textures: sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, sandy clay = 0.32, and clay = 0.37.

^h Rate Very Low if thickness of loamy/clayey layers is more than 10 inches.

PALM BEACH COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Palm Beach County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
An	1	Anclote	Very High	Very High	Low ^b	0.10
AsF	1	Arents	Medium	High		0.10
AU	1	Arents	High	Variable		0.10
AU	2	Urban Land	Very High	Variable		No Value
AX	1	Arents, Organic	High	Variable		0.10
AX	2	Urban Land	Very High	Variable		No Value
Ba	1	Basinger	Very High	Very High	Low ^b	0.10
Bc	1	Basinger	Very High	Very High	Low ^b	0.10
Bc	2	Urban Land	Very High	Variable		No Value
Bm	1	Basinger	Very High	Very High	Low ^b	0.10
Bm	2	Myakka	Very High	High	Medium	0.10
Be	1	Beaches	Very High	Very High		0.05
Bo	1	Boca	Very High	Very High ^c	Low ^b	0.10
Cc	1	Canaveral	High	High ^d		0.10
Cc	2	Urban Land	Very High	Variable		No Value
Ch	1	Chobee	Very High	Very Low	High	0.15
CuB	1	Cocoa	Low ^b	High ^c		0.10
CuB	2	Urban Land	Very High	Variable		No Value
Da	1	Dania	Very High	Very High	Medium	0.02
Fa	1	Floridana	Very High	Low	High	0.10
Ha	1	Hallandale	Very High	Very High	Low ^b	0.10
Ho	1	Holopaw	Very High	Very High ^c	Medium	0.10
Im	1	Immokalee	Very High	High	Medium	0.10
Ju	1	Jupiter	Very High	Very High	Medium	0.10
La	1	Lauderhill	Very High	Very High	Medium	0.02
Mk	1	Myakka	Very High	High	Medium	0.10
Mu	1	Myakka	Very High	High	Medium	0.10
Oc	1	Okeechobee	Very High	Very High	Medium	0.02
On	1	Okeelanta	Very High	Very High	Medium	0.02
Os	1	Oldsmar	Very High	Low	Medium	0.10
Pa	1	Pahokee	Very High	Very High	Medium	0.02
PbB	1	Palm Beach	Low ^b	Very High ^c		0.10
PbB	2	Urban Land	Very High	Very High		No Value
PcB	1	Paola	Low ^b	Very High		0.10

PALM BEACH COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Palm Beach County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
Pd	1	Pineda	Very High	High ^c	High	0.10
Pe	1	Pinellas	Very High	Low	Medium	0.10
Pf	1	Pits	Variable	Variable		0.10
Pg	1	Placid	Very High	Very High	Low ^b	0.10
PhB	1	Pomello	High	High ^d		0.10
Po	1	Pompano	Very High	Very High	Low ^b	0.10
QaB	1	Quartzipsammets	Low ^b	Very High		0.10
Ra	1	Riviera	Very High	High ^c	High	0.10
Rd	1	Riviera	Very High	High ^c	High	0.10
Ru	1	Riviera	Very High	High ^c	High	0.10
Ru	2	Urban Land	Very High	Variable		No Value
Sa	1	Sanibel	Very High	Very High	Low ^b	0.02
ScB	1	St. Lucie	Low ^b	Very High		0.10
SuB	1	Paola	Low ^b	High ^d		0.10
SuB	2	Urban Land	Very High	Variable		No Value
Ta	1	Tequesta	Very High	Very High ^c	Low ^b	0.02
Tc	1	Terra Ceia	Very High	Very High	Medium	0.02
Tm	1	Kesson	Very High	Very High		0.10
To	1	Wulfert	Very High	Very High		0.02
Tr	1	Torry	Very High	Very High	Medium	0.02
Ud	1	Udorthents	Low ^b	Variable		0.10
Ur	1	Urban Land	Very High	Variable		No Value
Wa	1	Wabasso	Very High	Low	High	0.10
Wn	1	Winder	Very High	Low	High	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Low if thickness of loamy/clayey layers is more than 10 inches thick.

^d Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^e Rate High if combined thickness of layers with chroma 3 or more and Bh horizons is more than 20 inches.

POLK COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for Polk County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
002	1	Apopka	Low ^b	Low		0.10
003	1	Candler	Low ^b	High		0.10
004	1	Candler	Low ^b	High ^c		0.10
005	1	EauGallie	Very High	Low	Medium	0.10
006	1	Eaton	Very High	Low	High	0.10
007	1	Pomona	Very High	Low	Medium	0.10
008	1	Hydraquents	Very High	Very Low	High	0.37
009	1	Lynne	Very High	Low	High	0.10
010	1	Malabar	Very High	Low	Medium	0.10
011	1	Arents	Low ^b	Variable		0.10
012	1	Neilhurst	Low ^b	High ^c		0.10
013	1	Samsula	Very High	Very High	Medium	0.02
014	1	Sparr	High	Low		0.10
015	1	Tavares	Low ^b	High		0.10
016	1	Urban Land	Very High	Variable		No Value
017	1	Smyrna	Very High	High	Medium	0.10
017	2	Myakka	Very High	High	Medium	0.10
019	1	Floridana	Very High	Low	High	0.10
020	1	Fort Meade	Low ^b	High		0.15
021	1	Immokalee	Very High	High	Medium	0.10
022	1	Pomello	High	High ^c		0.10
023	1	Ona	Very High	High	Medium	0.10
024	1	Nittaw	Very High	Very Low		0.24
025	1	Placid	Very High	Very High	Low ^b	0.10
025	2	Myakka	Very High	High	Medium	0.10
026	1	Lochloosa	High	Low		0.10
027	1	Kendrick	Low ^b	Low		0.10
029	1	St. Lucie	Low ^b	Very High		0.10
030	1	Pompano	Very High	Very High	Low ^b	0.10
031	1	Adamsville	High	High		0.10
032	1	Kaliga	Very High	Low	Medium	0.02
033	1	Holopaw	Very High	Very High ^d	Medium	0.10
034	1	Anclote	Very High	Very High	Low ^b	0.10
035	1	Hontoon	Very High	Very High	Low ^b	0.02
036	1	Basinger	Very High	Very High	Low ^b	0.10
037	1	Placid	Very High	Very High		0.10

POLK COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Polk County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
038	1	Electra	High	Low		0.10
039	1	Arents	High	Variable		0.10
040	1	Wauchula	Very High	Low	High	0.10
041	1	St. Johns	Very High	High	Medium	0.10
041	2	Basinger	Very High	Very High		0.10
042	1	Felda	Very High	Low	Medium	0.10
043	1	Oldsmar	Very High	Low	Medium	0.10
044	1	Paisley	Very High	Very Low	Very High	0.15
046	1	Astatula	Low ^b	High		0.10
047	1	Zolfo	High	High ^c		0.10
048	1	Chobee	Very High	Very Low	High	0.15
049	1	Adamsville	High	High		0.10
049	2	Urban Land	Very High	Variable		No Value
050	1	Candler	Low ^b	High		0.10
050	2	Urban Land	Very High	Variable		No Value
051	1	Pomona	Very High	Low	Medium	0.10
051	2	Urban Land	Very High	Variable		No Value
053	1	Myakka	Very High	High	Medium	0.10
053	2	Immokalee	Very High	High	Medium	0.10
053	3	Urban Land	Very High	Variable		No Value
054	1	Pomello	High	High ^c		0.10
054	2	Urban Land	Very High	Variable		No Value
055	1	Sparr	High	Low		0.10
055	2	Urban Land	Very High	Variable		No Value
057	1	Haplaquents	Very High	Variable		0.32
058	1	Pits	Very High	Variable		Variable ^e
059	1	Arents	High	Variable		0.10
059	2	Urban Land	Very High	Variable		No Value
060	1	Arents	Medium	Variable		0.17
061	1	Arents	Medium	Variable		0.10
061	2	Urban Land	Very High	Variable		No Value
062	1	Wabasso	Very High	Low	High	0.10
063	1	Tavares	Low ^b	High		0.10
063	2	Urban Land	Very High	Variable		No Value
064	1	Neilhurst	Low ^b	High ^c		0.10
064	2	Urban Land	Very High	Variable		0.10

POLK COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for Polk County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
066	1	Fort Meade	Low ^b	High		0.15
066	2	Urban Land	Very High	Variable		No Value
067	1	Bradenton	Very High	Very Low	High	0.15
068	1	Arents	Low ^b	Variable		0.10
070	1	Duette	Low ^b	High ^c		0.10
072	1	Bradenton	Very High	Very Low		0.15
072	2	Felda	Very High	Low		0.10
072	3	Chobee	Very High	Very Low		0.15
073	1	Gypsum Spoil	High	Variable		Variable ^e
074	1	Narcossee	High	High ^c		0.10
075	1	Valkaria	Very High	Very High	Low ^b	0.10
076	1	Millhopper	Low ^b	Low		0.10
077	1	Satellite	High	Very High		0.10
078	1	Paisley	Very High	Very Low	Very High	0.15
080	1	Chobee	Very High	Very Low		0.15
081	1	St. Augustine	High	Very High ^f		0.10
082	1	Felda	Very High	Low		0.10
083	1	Archbold	Low ^b	Very High		0.10
085	1	Winder	Very High	Very Low	High	0.10
086	1	Felda	Very High	Low	Medium	0.10
087	1	Basinger	Very High	Very High	Low ^b	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.

^b Rate Very Low where percent ground cover is greater than 75%.

^c Rate Very High if combined thickness of layers with chroma 3 or more and Bh horizons is less than 20 inches.

^d Rate Low if thickness of loamy/clayey layers is more than 10 inches thick.

^e The following K-factors should be used for the following on-site surface textures: sand = 0.10, loamy sand = 0.15, sandy loam = 0.20, sandy clay loam = 0.24, sandy clay = 0.32, and clay = 0.37.

^f Rate High if combined thickness of layers with chroma 3 or more and Bh horizons is more than 20 inches.

ST. LUCIE COUNTY

Table 13. Runoff, Leaching Potentials and K-Factors for St. Lucie County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
001	1	Anclote	Very High	Very High	Low ^b	0.10
002	1	Ankona	Very High	Low	Medium	0.10
003	1	Ankona	Very High	Low	Medium	0.10
003	2	Urban Land	Very High	Variable		No Value
004	1	Arents	High	Variable		0.10
005	1	Arents	Low ^b	Variable		0.10
006	1	Arents	Medium	Variable		0.10
007	1	Astatula	Low ^b	High		0.10
008	1	Basinger	Very High	Very High	Low ^b	0.10
009	1	Beaches	Very High	Very High		0.05
010	1	Canaveral	High	Very High ^c		0.10
011	1	Chobee	Very High	Very Low	High	0.15
012	1	Electra	High	Low		0.10
013	1	Floridana	Very High	Low	High	0.10
014	1	Fluvaquents	Very High	Variable		0.37
015	1	Hallandale	Very High	Very High	Low ^b	0.10
016	1	Hilolo	Very High	Very Low	High	0.10
017	1	Hobe	Low ^b	Very High ^d		0.10
018	1	Hontoon	Very High	Very High	Medium	0.02
019	1	Jonathan	Medium	High		0.10
020	1	Kaliga	Very High	Low	Medium	0.02
021	1	Lawnwood	Very High	High	High	0.10
022	1	Lawnwood	Very High	High	High	0.10
022	2	Urban Land	Very High	Variable		No Value
023	1	Malabar	Very High	Low	Medium	0.10
024	1	Myakka	Very High	High	Medium	0.10
025	1	Nettles	Very High	High	High	0.10
026	1	Oldsmar	Very High	Low	Medium	0.10
027	1	Palm Beach	Low ^b	Very High ^c		0.10
028	1	Paola	Low ^b	Very High ^c		0.10
029	1	Pendarvis	High	High ^e		0.10
030	1	Pendarvis	High	High ^e		0.10
030	2	Urban Land	Very High	Variable		No Value
031	1	Pepper	Very High	Low	High	0.10
032	1	Pineda	Very High	High ^f	High	0.10
033	1	Pits	Very High	Variable		0.10
034	1	Pompano	Very High	Very High	Low ^b	0.10
035	1	Pompano Variant	Very High	Very High		0.10
035	1	Kaliga Variant	Very High	Low		0.02

ST. LUCIE COUNTY (Cont.)

Table 13. Runoff, Leaching Potentials and K-Factors for St. Lucie County Soils.

Map Unit	Seq. No. ^a	Soil Name	Undrained Runoff Potential	Undrained and Drained Leaching Potential	Drained Runoff Potential	K-Factor
036	1	Pople	Very High	Very High ^c	High	0.10
037	1	Riviera	Very High	Low	High	0.10
038	1	Riviera	Very High	Low	High	0.10
039	1	Salerno	Very High	High	Medium	0.10
040	1	Samsula Variant	Very High	Very High	Medium	0.02
041	1	Satellite	High	Very High		0.10
042	1	St. Lucie	Low ^b	Very High		0.10
043	1	Susanna	Very High	Very High ^f	High	0.10
044	1	Tantile	Very High	High ^e	High	0.10
045	1	Terra Ceia	Very High	Very High		0.10
046	1	Turnbull Variant	Very High	Very High		0.32
047	1	Urban Land	Very High	Variable		No Value
048	1	Wabasso	Very High	Low	High	0.10
049	1	Wabasso Variant	Very High	Low	High	0.10
050	1	Waveland	Very High	High	High	0.10
051	1	Waveland	Very High	High	High	0.10
051	2	Lawnwood	Very High	High	High	0.10
052	1	Waveland	Very High	High	High	0.10
052	2	Urban Land	Very High	Variable		No Value
053	1	Welaka Variant	Low ^b	High		0.10
054	1	Winder	Very High	Very Low	High	0.10
054	2	Winder	Very High	Very Low	High	0.10
055	1	Winder	Very High	Very Low	High	0.10
056	1	Winder Variant	Very High	Very Low	High	0.10
049	1	Wabasso Variant	Very High	Low	High	0.10
050	1	Waveland	Very High	High	High	0.10
051	1	Waveland	Very High	High	High	0.10
051	2	Lawnwood	Very High	High	High	0.10
052	1	Waveland	Very High	High	High	0.10
052	2	Urban Land	Very High	Variable		No Value
053	1	Welaka Variant	Low ^b	High		0.10
054	1	Winder	Very High	Very Low	High	0.10
054	2	Winder	Very High	Very Low	High	0.10
055	1	Winder	Very High	Very Low	High	0.10
056	1	Winder Variant	Very High	Very Low	High	0.10

^a Seq. No. indicates a particular soil series name among one or more names constituting a map unit name.
^b Rate Very Low where percent ground cover is greater than 75%.
^c Rate High if combined thickness of layers, with chroma 3 or more and Bh horizons, is more than 20 inches.
^d Rate Very Low if thickness of loamy/clayey layers is more than 10 inches.
^e Rate Very High if combined thickness of layers, with chroma 3 or more and Bh horizons, is less than 20 inches.
^f Rate Low if thickness of loamy/clayey layers is more than 10 inches.