

Crataegus laevigata: English Hawthorn¹

Edward F. Gilman and Dennis G. Watson²

Introduction

English Hawthorn reaches a height 20 to 25 feet. It grows rapidly in a pyramidal form to about 20 feet, then the crown expands to become oval or irregular. The tree tolerates most soils, growing well in clay, but prefers heavy, dry loam. The main ornamental feature is white or pink flowers borne in spring and good fall color. Some types produce ornamental scarlet fruits. Though quite ornamental, Hawthorns are severely affected by insect and disease problems.




Figure 1. Middle-aged *Crataegus laevigata*: English Hawthorn

General Information

Scientific name: *Crataegus laevigata*

Pronunciation: kruh-TEE-gus lee-vih-GAY-tuh

Common name(s): English Hawthorn

Family: *Rosaceae*

USDA hardiness zones: 4B through 8B (Fig. 2)

Origin: not native to North America

Invasive potential: little invasive potential

Uses: urban tolerant; street without sidewalk; reclamation; screen; espalier; sidewalk cutout (tree pit); tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; highway median; Bonsai

Availability: not native to North America




Figure 2. Range

1. This document is ENH369, one of a series of the Environmental Horticulture, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Reviewed May 2011. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

Description

Height: 20 to 25 feet

Spread: 15 to 25 feet

Crown uniformity: irregular

Crown shape: pyramidal, oval

Crown density: moderate

Growth rate: moderate

Texture: fine

Foliage

Leaf arrangement: alternate (Fig. 3)

Leaf type: simple

Leaf margin: lobed, serrate

Leaf shape: ovate, obovate

Leaf venation: pinnate

Leaf type and persistence: deciduous

Leaf blade length: less than 2 inches

Leaf color: green

Fall color: no color change

Fall characteristic: not showy

Flower

Flower color: lavender, white/cream/gray, pink

Flower characteristics: showy

Fruit

Fruit shape: round

Fruit length: less than .5 inch

Fruit covering: fleshy

Fruit color: red

Fruit characteristics: attracts birds; showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches droop; not showy; typically multi-trunked; no thorns

Pruning requirement: needed for strong structure

Breakage: resistant

Current year twig color: brown

Current year twig thickness: medium

Wood specific gravity: unknown

Culture

Light requirement: full sun

Soil tolerances: clay; sand; loam; alkaline; acidic; occasionally wet; well-drained

Drought tolerance: high

Aerosol salt tolerance: low

Other

Roots: not a problem

Winter interest: yes

Outstanding tree: no

Ozone sensitivity: unknown

Verticillium wilt susceptibility: resistant

Pest resistance: sensitive to pests/diseases




Figure 3. Foliage

Use and Management

The tree casts heavy shade, and turf can not be grown underneath, if lower branches are left on the trunk. Persistent lower branches add to the ornamental characteristic of the tree and it makes a nice specimen in a lawn for all seasons when left unpruned. When lower branches are removed, this tough tree can be used as a street or parking lot tree where overhead space is limited by powerlines or other features. Branches bear sharp thorns which can inflict pain as they meet flesh. Grows well in tree pits and other confined soil spaces but probably should not be planted there due to sharp thorns. Plant in the open with good air circulation to help reduce disease, since leaf diseases can be common.

A number of improved cultivars are offered in garden centers including 'Crimson Cloud' with red flowers. This cultivar is nearly thornless and is preferred over the species. A number of other cultivars are offered in garden centers: 'Flore-Plena' - double, white flowers; 'Gireoudii' - foliage variegated, white flowers, large red fruit; 'Masekii' - pale

rose, double flowers; 'Paulii' ('Paul's Scarlet') - deep pink, double flowers; 'Pendula' - weeping growth habit; 'Punica' - single, pink flowers; 'Rosea' - pink flowers; 'Stricta' - upright growth habit.

Pests

Aphids on small trees can be partially controlled with strong sprays of water from a garden hose, if the colony is in the lower branches. Sometimes the aphids themselves are not seen but the distorted growth, honeydew on the leaves, and sooty mold growing on the honeydew are obvious.

Borer attacks may be prevented if the trees are kept in good vigor with regular fertilization.

Leaf miners symptoms are brown blotches on the leaves.

Lace bugs can be a serious, though occasional, problem. The insect feeding on the undersides of the leaves causes chlorotic flecks on the upper leaf surfaces. The lower sides of the leaves are covered with small, brown, sticky flecks.

The pear slug skeletonizes Hawthorn leaves and these sawfly larvae have a slimy appearance. A few insects can be washed off with a garden hose.

Tent caterpillar nests can be pruned out while still small. Sprays of *Bacillus thuringiensis* may be used. Do not burn nests while the nests are in the tree. The injury from the fire may exceed that caused by the insects.

Scales may be controlled with horticultural oil sprays.

Spider mites are so small they can cause much foliage discoloration before being detected.

Diseases

Fire blight: This disease can be severe in some parts of the country. The first noticeable symptom of fire blight is the browning of branch tips. The tips appear to be burned or scorched and the dead, brown leaves droop but hang on the tree. Cankers form and the bacteria is washed farther down the branch by rain. The bacteria, *Erwinia amylovora*, are spread from diseased to healthy twigs by rain, bees, and other mechanical means. There is no satisfactory chemical control. The disease is less of a problem if trees are not located near apple or pear orchards. Prune out blighted branch tips by cutting a foot or two beyond the diseased wood. Over-fertilizing with nitrogen fertilizer may increase tree susceptibility to fire blight.

Leaf blight attacks most Hawthorns but especially English Hawthorn. The symptoms are small reddish brown spots on the leaves which may run together. Infected leaves drop in August and severely infected trees may be completely bare.

Cedar Hawthorn rust causes orange or rust colored spots on the leaves leading to early defoliation. The fruits and twigs are also attacked. Juniper is an alternate host. Cedar-quince rust attacks fruits. Washington, Lavelle and Cockspur Hawthorn are resistant to rust diseases.

Scab causes leaf spotting and defoliation. The fruit have black raised spots on them.

Powdery mildew causes a white powdery growth on the leaves.