

Ilex opaca: American Holly¹

Edward F. Gilman and Dennis G. Watson²

Introduction

A popular landscape plant since the beginning of American history, this broad-leaved evergreen has served a variety of uses through the years (Fig. 1). The American Indians used preserved Holly berries as decorative buttons and were much sought after by other tribes who bartered for them. The wood has been used for making canes, scroll work and furniture, and has even been substituted for ebony in inlay work when stained black.



Figure 1. Middle-aged *Ilex opaca*: American Holly
Credits: Ed Gilman

General Information

Scientific name: *Ilex opaca*

Pronunciation: EYE-lecks oh-PAY-kuh

Common name(s): American Holly

Family: *Aquifoliaceae*

USDA hardiness zones: 5B through 9 (Fig. 2)

Origin: native to North America

Invasive potential: little invasive potential

Uses: street without sidewalk; specimen; hedge; reclamation; screen; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; sidewalk cutout (tree pit); urban tolerant; highway median; Bonsai

Availability: generally available in many areas within its hardiness range



Figure 2. Shaded area represents potential planting range.

1. This document is ENH463, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date November 1993. Revised December 2006. Reviewed February 2014. Visit the EDIS webwite at <http://edis.ifas.ufl.edu>.
2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Description

Height: 35 to 50 feet

Spread: 15 to 25 feet

Crown uniformity: symmetrical canopy with a regular (or smooth) outline, and individuals have more or less identical crown forms

Crown shape: pyramidal

Crown density: dense

Growth rate: slow

Texture: medium

Foliage

Leaf arrangement: alternate (Fig. 3)

Leaf type: simple

Leaf margin: pectinate, entire, spiny

Leaf shape: elliptic (oval), lanceolate

Leaf venation: banchidrome; pinnate

Leaf type and persistence: evergreen, broadleaf evergreen

Leaf blade length: less than 2 inches, 2 to 4 inches

Leaf color: green

Fall color: no color change

Fall characteristic: not showy




Figure 3. Foliage of American Holly.

Flower

Flower color: green, white

Flower characteristics: pleasant fragrance; inconspicuous and not showy

Fruit

Fruit shape: round

Fruit length: less than .5 inch

Fruit covering: fleshy

Fruit color: red

Fruit characteristics: attracts birds; no significant litter problem; persistent on the tree; showy

Trunk and Branches

Trunk/bark/branches: bark is thin and easily damaged from mechanical impact; droop as the tree grows, and will require pruning for vehicular or pedestrian clearance

beneath the canopy; not particularly show; should be grown with a single leader; no thorns

Pruning requirement: needs little pruning to develop a strong structure

Breakage: resistant

Current year twig color: green, brown

Current year twig thickness: medium

Wood specific gravity: 0.61

Culture

Light requirement: tree grows in the shade; tree grows in full sun

Soil tolerances: sand; loam; clay; acidic; slightly alkaline; extended flooding; well-drained

Drought tolerance: high

Aerosol salt tolerance: high

Soil salt tolerance: moderate

Other

Roots: not a problem

Winter interest: tree has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

Outstanding tree: not particularly outstanding

Ozone sensitivity: tolerant

Verticillium wilt susceptibility: not known to be susceptible

Pest resistance: long-term health usually not affected by pests

Use and Management

American Holly is a beautifully shaped tree, with a symmetrical, dense, wide pyramidal form. The spiny, dull green leaves are accented with clusters of red berries which persist throughout the fall and winter. Male and female flowers appear on separate trees and trees of both sexes must be located in the same neighborhood to ensure production of berries on the female plants. American Holly is ideal for use as a street or courtyard tree (with lower branches removed), framing tree, specimen, barrier planting or screen. Roots are shallow and finely branched, and rarely invasive due to their great number and relatively small diameter. This native tree is ideal for naturalizing on moist, slightly acid soils, and the fruit is very attractive to wildlife, serving as an excellent food source. A 35-foot-tall tree can be 20 feet wide in 40 years.

Growing well in full sun to partial shade, American Holly should be located on fertile, well-drained but moist, slightly acid soils below 6.5 pH. Berry production is highest in full

sun on female trees. American Holly foliage thins during drought but insect and disease infestations are usually minimal.

Hundreds of cultivars of American Holly have been developed and hybridized over the years, providing variety of form, leaf characteristics, and fruit color. The following is a list of some available cultivars and hybrids: 'Carolina #2' has dark green leaves and abundant fruit; 'George E. Hart' has a narrow conical growth habit with small dark green leaves; 'Hume No. 2' has compact, dark green foliage, and heavy fruit; 'Croonenburg' has dark green, slightly glossy foliage and abundant fruit; 'Howard' has dense, glossy green leaves with few spines, large fruit, and a more compact form; 'Greenleaf' is softer in form than 'Croonenburg', fast-growing, and responds well to shearing; 'Jersey Knight', a male cultivar, is very hardy and has excellent foliage; 'Jersey Princess', a female cultivar, has excellent form and shining dark green leaves; 'Rotunda' has an upright growth habit, smooth, entire, glossy green leaves, and is profusely fruiting; 'Ft. McCoy', 'Dupre', 'Lake City', 'Savannah', and 'Taber' all have quite spiny leaves. 'Savannah' also has wavy curved foliage and dark, heavy fruit; 'East Palatka', a female cultivar, is actually *Ilex x attenuata*, a cross of *Ilex cassine* x *Ilex opaca*, and has only a small spine at the leaf tip. Those with yellow berries include: 'Xanthocarpa', 'Canary', and 'Morgan Gold'.

Those cultivars particularly adapted for the south include: 'Amy' - female, abundant fruit; 'Bountiful' - cone-shaped, compact, dark red fruit annually; 'Calloway' - yellow fruit; 'Miss Helen' - dark red, abundant fruit; 'Slim Jim' - open, slender Holly with narrow leaves; 'Steward's Cream Crown' - creamy, marginal venation; 'Yellow Jacket' - cadmium orange fruit.

Propagation is by cuttings or grafting.

Pests

Holly leaf miner larvae mines out the leaf middle leaving yellow or brown trails.

Scales of various types may infest Holly.

Spider mites cause discoloration and speckling of Holly foliage.

Diseases

Tar spot may occasionally cause small yellow spots on the leaves in early summer. Eventually the spots turn reddish brown with narrow yellow borders. Leaves may not drop prematurely but the infected areas drop out leaving holes in the leaves. Gather up and destroy badly infected leaves.

Many different fungi cause leaf spots on Holly. Reduce the injury caused by leaf spots by keeping trees healthy. Dispose of diseased leaves.

Cankers caused by several different fungi lead to sunken areas on stems and plant dieback. Keep trees healthy and prune out infected branches.

Spine spot is small gray or yellow spots with purple margins and is caused by spines of one leaf puncturing an adjacent leaf.

Chlorosis symptoms are light green or yellowish leaves with darker green veins. This problem is often due to a high pH leading to iron deficiency. Use acidifying fertilizers and sulfur to bring down the pH. Sprays of iron chelate will green up plants.

In northern climates, Hollies sometimes scorch during the late winter due to rapid and wide temperature fluctuations. Shade plants during the winter to prevent the problem.

Purple blotches on the leaves are caused by some environmental factor such as nutrient deficiencies, drought, and winter injury.

Black root rot can be damaging.