

Lagerstroemia indica: Crapemyrtle¹

Edward F. Gilman and Dennis G. Watson²

Introduction

A long period of striking summer flower color, attractive fall foliage, and good drought-tolerance all combine to make Crape-Myrtle a favorite small tree for either formal or informal landscapes. It is highly recommended for planting in urban and suburban areas.




Figure 1. Young *Lagerstroemia indica*: Crapemyrtle

General Information

Scientific name: *Lagerstroemia indica*

Pronunciation: lay-ger-STREE-mee-uh IN-dih-kuh

Common name(s): Crapemyrtle

Family: *Lythraceae*

USDA hardiness zones: 7A through 9A (Fig. 2)

Origin: not native to North America

Invasive potential: has been evaluated using the IFAS Assessment of the Status of Non-Native Plants in Florida's Natural Areas (Fox et al. 2005). This species is not documented in any undisturbed natural areas in Florida. Thus, it is not considered a problem species and may be used in Florida.

Uses: street without sidewalk; specimen; deck or patio; container or planter; trained as a standard; parking lot island < 100 sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; urban tolerant; highway median; shade

-
1. This document is ENH-501, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised March 2007. Reviewed October 2003. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
 2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

Availability: not native to North America




Figure 2. Range

Description

Height: 10 to 30 feet

Spread: 15 to 25 feet

Crown uniformity: symmetrical

Crown shape: vase

Crown density: moderate

Growth rate: moderate

Texture: medium

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)

Leaf type: simple

Leaf margin: entire

Leaf shape: elliptic (oval), obovate, oblong

Leaf venation: pinnate

Leaf type and persistence: deciduous

Leaf blade length: less than 2 inches, 2 to 4 inches

Leaf color: green

Fall color: orange, red, yellow

Fall characteristic: showy

Flower

Flower color: white/cream/gray, pink, purple, lavender, red

Flower characteristics: very showy

Fruit

Fruit shape: oval, round

Fruit length: less than .5 inch

Fruit covering: dry or hard

Fruit color: brown

Fruit characteristics: does not attract wildlife; showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches droop; showy; typically multi-trunked; thorns

Pruning requirement: little required

Breakage: resistant

Current year twig color: brown, green

Current year twig thickness: thin

Wood specific gravity: unknown

Culture

Light requirement: full sun

Soil tolerances: clay; sand; loam; alkaline; acidic; well-drained

Drought tolerance: high

Aerosol salt tolerance: moderate

Other

Roots: not a problem

Winter interest: yes

Outstanding tree: no

Ozone sensitivity: unknown

Verticillium wilt susceptibility: resistant

Pest resistance: resistant to pests/diseases

Use and Management

Available in all shades of white, pink, red, or lavender, the 6- to 12-inch-long clustered blooms appear on the tips of branches during late spring and summer in USDA hardiness zones 9 and 10, and summer in other areas. The individual flowers are ruffled and crinkly as to appear made of crepe paper. The smooth, peeling bark and multi-branched, open habit of Crape-Myrtle make it ideal for specimen planting where its bright red to orange-colored fall



Figure 3. Foliage



Figure 4. Flower

leaves add further interest. Most forms of the tree are upright, upright-spreading, or vase-shaped, spreading out as they ascend. Most tree types grow to 20 to 25 feet tall although there are more dwarf types available. The upright, vase-shaped crown makes the tall-growing selections well-suited for street tree planting.

Pruning should be done in late winter or early in the spring before growth begins because it is easier to see which branches to prune. New growth can be

pinched during the growing season to increase branchiness and flower number. Pruning methods vary from topping to cutting Crape-Myrtle nearly to the ground each spring to the removal of dead wood and old flower stalks only. Topping creates several long, thin branches from each cut which droop down under the weight of the flowers. This practice disfigures the nice trunk and branch structure. Lower branches are often thinned to show off the trunk form and color. You can remove the spent flower heads to encourage a second flush of flowers and to prevent formation of the brown fruits. Since cultivars are now available in a wide range of growth heights, severe pruning should not be necessary to control size. Severe pruning or topping can stimulate basal sprouting which can become a constant nuisance, requiring regular removal. Some trees sprout from the base of the trunk and roots even without severe heading. This can be a maintenance nuisance.

Crape-Myrtle grows best in full sun with rich, moist soil but will tolerate less hospitable positions in the landscape just as well, once it becomes established. It grows well in limited soil spaces in urban areas such as along boulevards, in parking lots, and in small pavement cutouts if provided with some irrigation until well established. They tolerate clay and alkaline soil well. However, the flowers of some selections may stain car paint. Insect pests are few but Crape-Myrtle is susceptible to powdery mildew damage, especially when planted in some shade or when the leaves are kept moist. There are new cultivars (many developed by the USDA) available which are resistant to powdery mildew and aphids.

Many cultivars of Crape-Myrtle are available: hybrid 'Acoma', 14 to 16 feet tall, white flowers, purple-red fall foliage, mildew resistant; hybrid 'Biloxi', 25 feet tall, pale pink blooms, orange-red fall foliage, hardy and mildew resistant; 'Cherokee', 10 to 12 feet, bright red flowers; 'Powhatan', 14 to 20 feet, clear yellow fall foliage, medium purple flowers. The hybrid cultivars 'Natchez', 30 feet tall, pure white flowers, resistant to aphids, one of the best Crape-Myrtles; 'Muskogee', 24 feet tall, light lavender flowers, and 'Tuscarora', 16 feet tall, dark coral pink blooms, are hybrids between *Lagerstroemia indica* and *Lagerstroemia fauriei* and have greater resistance to mildew. The cultivar

'Crape-Myrtlettes' have the same color range as the species but only grow to three to four feet high. The National Arboretum releases are generally superior because they have been selected for their disease resistance. These releases may prove more resistant to powdery mildew in the Deep South, although further testing needs to be done to confirm this.

Propagation is by cuttings or seed.

Pests

Aphids often infest the new growth causing an unsightly but harmless sooty mold to grow on the foliage. Heavy aphid infestations cause a heavy black sooty mold which detracts from the tree's appearance.

Diseases

Powdery mildew can severely affect Crape-Myrtle. Select resistant cultivars and hybrids to avoid this disease. Leaf spots are only a minor concern and do not require treatment.

Literature Cited

Fox, A.M., D.R. Gordon, J.A. Dusky, L. Tyson, and R.K. Stocker (2005) IFAS Assessment of the Status of Non-Native Plants in Florida's Natural Areas. Cited from the Internet (November 3, 2006), <http://plants.ifas.ufl.edu/assessment.html>