

*Lagerstroemia x 'Cherokee': 'Cherokee' Crapemyrtle*¹

Edward F. Gilman and Dennis G. Watson²

Introduction

A long period of striking summer flower color, attractive fall foliage, and good drought-tolerance all combine to make this cultivar of Crape-Myrtle a favorite very small tree or large shrub for either formal or informal landscapes. Most are seen seven to nine feet tall with a similar spread. It is highly recommended for planting in urban areas.




Figure 1. Young *Lagerstroemia x 'Cherokee': 'Cherokee'* Crapemyrtle

General Information

Scientific name: *Lagerstroemia x*

Pronunciation: lay-ger-STREE-mee-uh

Common name(s): 'Cherokee' Crapemyrtle

Family: *Lythraceae*

USDA hardiness zones: 7A through 9A (Fig. 2)

Origin: not native to North America

Invasive potential: little invasive potential

Uses: tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; street without sidewalk; parking lot island < 100 sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; container or planter; trained as a standard; deck or patio; specimen; highway median

Availability: somewhat available, may have to go out of the region to find the tree

1. This document is ENH-505, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.




Figure 2. Range

Description

Height: 10 to 15 feet

Spread: 6 to 10 feet

Crown uniformity: symmetrical

Crown shape: vase

Crown density: moderate

Growth rate: moderate

Texture: medium

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)

Leaf type: simple

Leaf margin: entire

Leaf shape: oblong, elliptic (oval), obovate

Leaf venation: pinnate

Leaf type and persistence: deciduous

Leaf blade length: less than 2 inches, 2 to 4 inches

Leaf color: green

Fall color: orange, red, yellow

Fall characteristic: showy

Flower

Flower color: red

Flower characteristics: very showy

Fruit

Fruit shape: oval, round

Fruit length: less than .5 inch

Fruit covering: dry or hard

Fruit color: brown

Fruit characteristics: does not attract wildlife; showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches droop; showy; typically multi-trunked; thorns

Pruning requirement: little required

Breakage: resistant

Current year twig color: brown, green

Current year twig thickness: thin

Wood specific gravity: unknown

Culture

Light requirement: full sun

Soil tolerances: sand; loam; clay; acidic; slightly alkaline; well-drained

Drought tolerance: high

Aerosol salt tolerance: moderate

Other

Roots: not a problem

Winter interest: yes

Outstanding tree: no

Ozone sensitivity: unknown

Verticillium wilt susceptibility: resistant

Pest resistance: resistant to pests/diseases

Use and Management

The 6- to 12-inch-long clustered bright red blooms appear on the tips of branches during late spring and summer in USDA hardiness zones 9 and 10 and summer in other areas. If flowers appear pink instead of red, then you have not purchased the true 'Cherokee' cultivar. The individual flowers are ruffled and crinkly as to appear made of crepe paper. The smooth, peeling bark and multi-branched, open habit make it ideal for specimen planting where its bright red to orange-colored fall leaves add further interest.



Figure 3. Foliage



Figure 4. Flower

Pruning should be done in late winter or early in the spring before growth begins because it is easier to see which branches to prune. New growth can be pinched during the growing season to increase branchiness and flower number. Pruning methods vary from topping to cutting Crape-Myrtle nearly to the ground each spring to the removal of dead wood and old flower stalks only. Lower branches are often thinned to show off the trunk form and color. You can remove the spent flower heads to encourage a second flush of flowers and to prevent formation of

the brown fruits. Since cultivars are now available in a wide range of growth heights, severe pruning should not be necessary to control size. Severe pruning can stimulate basal sprouting which can become a constant nuisance, requiring regular removal. Some Crape-Myrtle trees sprout from the base of the trunk and roots even without severe heading.

Crape-Myrtle grows best in full sun with rich, moist soil but will tolerate less hospitable positions in the landscape just as well, once it becomes established. It grows well in limited soil spaces in urban areas such as along boulevards, in parking lots, and in small pavement cutouts if provided with some irrigation. They tolerate clay and alkaline soil well. However, the flowers of some selections may stain car paint. Insect pests are few (except for the Crape-Myrtle aphid) and Crape-Myrtle is susceptible to powdery mildew damage, especially when planted in some shade or when the leaves are kept moist. There are other new cultivars (many developed by the USDA) available which are resistant to powdery mildew.

Many other cultivars of Crape-Myrtle are available: hybrid 'Acoma', 14 to 16 feet tall, white flowers, purple-red fall foliage, mildew resistant; hybrid 'Biloxi', 25 feet tall, pale pink blooms, orange-red fall foliage, hardy and mildew resistant; 'Powhatan', 14 to 20 feet, clear yellow fall foliage, medium purple flowers. The hybrid cultivars 'Natchez', 30 feet tall, pure white flowers, 'Muskogee', 24 feet tall, light lavender flowers, and 'Tuscarora', 16 feet tall, dark coral pink blooms, are hybrids between *Lagerstroemia indica* and *Lagerstroemia fauriei* and have greater resistance to mildew. The cultivar 'Crape Myrtlettes' have the same color range as the species but only grow to three to four feet high. The National Arboretum releases are generally superior because they have been selected for their disease resistance.

Propagation is by cuttings.

Pests

Aphids often infest the new growth causing an unsightly but harmless sooty mold to grow on the

foliage. Heavy aphid infestations cause a heavy black sooty mold which detracts from the tree's appearance.

Diseases

Powdery mildew can severely affect Crape-Myrtle. Select resistant cultivars and hybrids to avoid this disease. Leaf spots are only a minor concern and do not require treatment.