


## *Lagerstroemia* x 'Muskogee': 'Muskogee' Crapemyrtle<sup>1</sup>

Edward F. Gilman and Dennis G. Watson<sup>2</sup>

### Introduction

A long period of striking summer flower color, attractive fall foliage, and good drought-tolerance all combine to make Crape-Myrtle a favorite small tree for either formal or informal landscapes. It is highly recommended for planting in urban and suburban areas.



**Figure 1.** Young *Lagerstroemia* x 'Muskogee': 'Muskogee' Crapemyrtle

### General Information

**Scientific name:** *Lagerstroemia* x

**Pronunciation:** lay-ger-STREE-mee-uh

**Common name(s):** 'Muskogee' Crapemyrtle

**Family:** *Lythraceae*

**USDA hardiness zones:** 7A through 10A (Fig. 2)

**Origin:** not native to North America

**Invasive potential:** little invasive potential

**Uses:** urban tolerant; street without sidewalk; specimen; deck or patio; container or planter; trained as a standard; parking lot island < 100 sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; highway median; shade

**Availability:** somewhat available, may have to go out of the region to find the tree

- 
1. This document is ENH-506, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. . Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
  2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean




Figure 2. Range

**Description**

**Height:** 20 to 25 feet

**Spread:** 15 to 25 feet

**Crown uniformity:** symmetrical

**Crown shape:** vase

**Crown density:** moderate

**Growth rate:** moderate

**Texture:** medium

**Foliage**

**Leaf arrangement:** opposite/subopposite (Fig. 3)

**Leaf type:** simple

**Leaf margin:** entire

**Leaf shape:** oblong, elliptic (oval), obovate

**Leaf venation:** pinnate

**Leaf type and persistence:** deciduous

**Leaf blade length:** less than 2 inches, 2 to 4 inches

**Leaf color:** green

**Fall color:** orange, red

**Fall characteristic:** showy

**Flower**

**Flower color:** lavender

**Flower characteristics:** very showy

**Fruit**

**Fruit shape:** oval, round

**Fruit length:** less than .5 inch

**Fruit covering:** dry or hard

**Fruit color:** brown

**Fruit characteristics:** does not attract wildlife; showy; fruit/leaves not a litter problem

### Trunk and Branches

**Trunk/bark/branches:** branches droop; showy; typically multi-trunked; thorns

**Pruning requirement:** little required

**Breakage:** resistant

**Current year twig color:** brown, green

**Current year twig thickness:** thin

**Wood specific gravity:** unknown

### Culture

**Light requirement:** full sun

**Soil tolerances:** sand; loam; clay; acidic; slightly alkaline; well-drained

**Drought tolerance:** high

**Aerosol salt tolerance:** moderate

### Other

**Roots:** not a problem

**Winter interest:** yes

**Outstanding tree:** no

**Ozone sensitivity:** unknown

**Verticillium wilt susceptibility:** resistant

**Pest resistance:** resistant to pests/diseases

### Use and Management

The 6- to 12-inch-long clustered lavender or pink blooms appear on the tips of branches during late spring and summer in USDA hardiness zones 9 and 10 and summer in other areas. The individual flowers are ruffled and crinkly as to appear made of crepe paper. The smooth, peeling bark and multi-branched, open habit of Crape-Myrtle make it ideal for specimen planting where its bright red to



Figure 3. Foliage



Figure 4. Flower

orange-colored fall leaves add further interest. The tree forms an upright to upright-spreading crown, the branches spreading out as they ascend. The tree grows 20 to 25 feet tall and almost as wide with an upright, vase-shaped crown making it well-suited for street tree planting.

Pruning should be done in late winter or early in the spring before growth begins because it is easier to see which branches to prune. New growth can be pinched during the growing season to increase

branchiness and flower number. Pruning methods vary from topping to cutting Crape-Myrtle nearly to the ground each spring to the removal of dead wood and old flower stalks only. Topping creates several long, thin branches from each cut which droop down under the weight of the flowers. This practice disfigures the nice trunk and branch structure. Lower branches are often thinned to show off the trunk form and color. You can remove the spent flower heads to encourage a second flush of flowers and to prevent formation of the brown fruits. Since cultivars are now available in a wide range of growth heights, severe pruning should not be necessary to control size. Severe pruning can stimulate basal sprouting which can become a constant nuisance, requiring regular removal. Some trees sprout from the base of the trunk and roots even without severe heading. This can be a maintenance nuisance.

Crape-Myrtle grows best in full sun with rich, moist soil but will tolerate less hospitable positions in the landscape just as well, once it becomes established. It grows well in limited soil spaces in urban areas such as along boulevards, in parking lots, and in small pavement cutouts if provided with some irrigation. They tolerate clay and alkaline soil well. However, the flowers may stain car paint. Aphids are the main insect pest of Crape-Myrtle but 'Muskogee' is resistant to powdery mildew. There are other new cultivars (many developed by the USDA) available which are resistant to powdery mildew and aphids.

Many other cultivars of Crape-Myrtle are available: hybrid 'Acoma', 14 to 16 feet tall, white flowers, purple-red fall foliage, mildew resistant; hybrid 'Biloxi', 25 feet tall, pale pink blooms, orange-red fall foliage, hardy and mildew resistant; 'Cherokee', 10 to 12 feet, bright red flowers; 'Powhatan', 14 to 20 feet, clear yellow fall foliage, medium purple flowers. The hybrid cultivars 'Natchez', 30 feet tall, pure white flowers, 'Tuscarora', 16 feet tall, dark coral pink blooms, are hybrids between *Lagerstroemia indica* and *Lagerstroemia fauriei* and have greater resistance to mildew. The cultivar 'Crape Myrtlettes' have the same color range as the species but only grow to three to four feet high. The National Arboretum releases are generally superior because they have been selected for their disease resistance.

Propagation is by cuttings or seed.

### Pests

Aphids often infest the new growth causing an unsightly but harmless sooty mold to grow on the foliage. Heavy aphid infestations cause a heavy black sooty mold which detracts from the tree's appearance.

### Diseases

Powdery mildew can severely affect Crape-Myrtle but 'Muskogee' is highly resistant.