

## ***Magnolia grandiflora* 'Majestic Beauty': 'Majestic Beauty' Southern Magnolia<sup>1</sup>**

---

Edward F. Gilman and Dennis G. Watson<sup>2</sup>

### **Introduction**

This is a rather large cultivar of Southern Magnolia with profuse flowering. Capable of growing at a moderate rate to a height of 50 feet with a 20 to 25-foot spread, this Southern Magnolia forms an open canopy displaying a coarse habit and is probably best suited for large properties. Leaves are large and dark shiny dark green making this a striking specimen tree. The lower branches often bend to the ground forming a pyramidal crown.

### **General Information**

**Scientific name:** *Magnolia grandiflora*


**Pronunciation:** mag-NO-lee-uh gran-dih-FLOR-uh

**Common name(s):** 'Majestic Beauty' Southern Magnolia

**Family:** *Magnoliaceae*

**USDA hardiness zones:** 7A through 10A (Fig. 2)

**Origin:** native to North America



**Figure 1.** Middle-aged *Magnolia grandiflora* 'Majestic Beauty': 'Majestic Beauty' Southern Magnolia

**Invasive potential:** little invasive potential

**Uses:** shade; specimen; street without sidewalk; screen; espalier; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; highway median

1. This document is ENH-535, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

**Availability:** somewhat available, may have to go out of the region to find the tree




Figure 2. Range

### Description

**Height:** 40 to 50 feet

**Spread:** 20 to 25 feet

**Crown uniformity:** symmetrical

**Crown shape:** pyramidal, oval

**Crown density:** dense

**Growth rate:** moderate

**Texture:** coarse

### Foliage

**Leaf arrangement:** alternate (Fig. 3)

**Leaf type:** simple

**Leaf margin:** entire

**Leaf shape:** elliptic (oval), ovate

**Leaf venation:** brachidodrome, pinnate

**Leaf type and persistence:** evergreen, broadleaf evergreen

**Leaf blade length:** 4 to 8 inches, 8 to 12 inches

**Leaf color:** green

**Fall color:** no color change

**Fall characteristic:** not showy

### Flower

**Flower color:** white/cream/gray

**Flower characteristics:** very showy

### Fruit

**Fruit shape:** elongated

**Fruit length:** 1 to 3 inches, 3 to 6 inches

**Fruit covering:** dry or hard

**Fruit color:** brown, red

**Fruit characteristics:** attracts birds; showy; fruit/leaves a litter problem

### Trunk and Branches

**Trunk/bark/branches:** branches droop; not showy; typically one trunk; thorns

**Pruning requirement:** little required

**Breakage:** resistant

**Current year twig color:** green

**Current year twig thickness:** thick

**Wood specific gravity:** 0.50

### Culture

**Light requirement:** full sun, partial sun or partial shade

**Soil tolerances:** sand; loam; clay; acidic; slightly alkaline; well-drained; occasionally wet

**Drought tolerance:** moderate

**Aerosol salt tolerance:** moderate

### Other

**Roots:** not a problem

**Winter interest:** no

**Outstanding tree:** yes


**Ozone sensitivity:** unknown

**Verticillium wilt susceptibility:** susceptible

**Pest resistance:** resistant to pests/diseases

### Use and Management

The 8-inch-long, leathery, oblong, shiny leaves are shed as new leaves emerge but the debris is well-hidden by the foliage of the lower limbs, if they



**Figure 3.** Foliage

are left on the tree. But some people consider this a litter nuisance when the large, slowly-decomposing leaves drop on the sidewalk, lawn, or patio. The underside of the leaves is covered with a fine, red-brown fuzz. In late spring and sporadically throughout the summer, huge, 8-inch-diameter, waxy, fragrant, white blossoms open to perfume the entire garden. Fuzzy brown cones follow these blooms, ripening in fall and winter to reveal bright red seeds which are used by a variety of wildlife.

Used as a striking garden specimen for large landscapes, Southern Magnolia can also serve as a street tree (with lower limbs removed). The only objection to this tree as a street tree might be the falling leaves and fruit. Its ease of growth and carefree nature make Southern Magnolia ideal for the low-maintenance landscape. With proper pruning, Southern Magnolia trees can also be used as an interesting espalier.

If moist, peaty soils are available, Southern Magnolia will thrive in full sun and hot conditions once established. If irrigation cannot be provided periodically, plants located in partial shade for several years after planting seem to grow better. Very drought tolerant when grown in areas with plenty of soil for root expansion. Only moderately drought tolerant in restricted-soil areas or in areas with poor, dry soil. Southern Magnolia prefers acid soil but will tolerate a slightly basic, even wet or clay soil. It is generally too hot and dry in central and western Texas

and Oklahoma, and the soil pH is often too alkaline for this tree.

The root system is wider spreading than most other trees, extending from the trunk a distance equal to about four times the canopy width. This makes it very difficult to save existing Magnolia trees on construction sites.

Be sure that there are no roots circling close to the trunk, as Magnolia is prone to girdling roots. Cut any circling roots prior to planting. Field-grown trees recover slowly from transplanting due to the wide-spreading root system in the nursery, and trees often transplant best in winter and spring, not in the fall.

The species germinated from seed is quite variable in growth rate and form with some trees dense and compact, others loose and open. A number of other cultivars are available: 'Bracken's Brown Beauty' has an unusually dark brown lower leaf surface; 'Cairo' has an early and long flowering period; 'Charles Dickens' has broad, nearly blunt leaves, large flowers and large red fruit; 'Edith Bogue' is the hardiest of the cultivars and will bloom when only two to three-years-old; 'Glen St. Mary' has a compact form, will bloom when young, is slow-growing, and the leaves have a bronze underside; 'Gloriosa' has large flowers and leaves; 'Goliath' has flowers up to 12 inches across, a long blooming period, and a bushy habit of growth; 'Hasse' can be used for a compact, dense hedge; 'Lanceolata' has a narrow pyramidal form, narrower leaves with rusty undersides; 'Little Gem' has a dwarf upright form, probably to 30 feet tall, small leaves and flowers, is very slow-growing, flowers heavily at an early age and for a long time during the summer (5-months), and has bronze leaf-undersides. It will bloom when only three to four feet tall and is excellent as a pruned evergreen hedge, for use as a small street tree or for use as an espalier. 'Praecox Fastigiata' has upright, narrow growth habit; 'Samuel Sommer' has an upright, rapid growth habit and flowers up to 14 inches across; 'Victoria' is very hardy, has small flowers, and rust-red leaf-undersides. There are others, often difficult to see real differences among a number of cultivars.

Propagation is by cuttings (for the cultivars), grafting, or seed.

### **Pests**

Scales of various types will infest twigs and leaves. Magnolia scale is the most common scale and can be one half-inch-across. Overwintering scales can be controlled with horticultural oil. Trees appear to grow fine even with heavy infestations, although they can be unsightly.

Tulip-poplar weevil (sassafras weevil) feeds as a leaf miner when young and chews holes in the leaves as an adult.

Magnolia borer is a problem on young nursery stock. It girdles the trunk usually just below the soil surface. Control is difficult but attainable with the proper material.

### **Diseases**

Magnolia may be subject to leaf spots, blights, scabs and black mildews caused by a large number of fungi, or a bacterium but they rarely require chemical controls. Raking up and disposing infected leaves may reduce leaf spots next year. Algae can also cause leaf spots.

Canker diseases will kill branches. Cankers on branches can be pruned out. Keep trees healthy with regular fertilization and by watering in dry weather.

Verticillium wilt may cause death of a few branches or, rarely, may kill the tree. Prune out dead branches and fertilize.