


## ***Syringa reticulata* 'Summer Snow': 'Summer Snow' Japanese Tree Lilac<sup>1</sup>**

---

Edward F. Gilman and Dennis G. Watson<sup>2</sup>

### **Introduction**

Although a Lilac, this member of the species is quite different in appearance than those with which gardeners are more familiar. Its rounded habit varies from symmetrical to irregular. Cultivars including 'Ivory Silk' and 'Summer Snow' could be used instead of the species due to the more consistent habit and more flowers. 'Summer Snow' is spectacular in flower and the persistent seed pods carry ornamental interest into the fall. This is a very large shrub or small tree, reaching a height of about 20 to 30 feet with a 20 to 25-foot-spread. The huge clusters of creamy white flowers, borne in early summer for about two weeks, are the main ornamental feature but lack the fragrance of the spring-blooming Lilacs -- this Lilac's fragrance is more suggestive of privet.



**Figure 1.** Mature *Syringa reticulata* 'Summer Snow': 'Summer Snow' Japanese Tree Lilac

### **General Information**

**Scientific name:** *Syringa reticulata*

**Pronunciation:** sih-RING-guh reh-tick-yoo-LAY-tuh

**Common name(s):** 'Summer Snow' Japanese Tree Lilac

**Family:** *Oleaceae*

**USDA hardiness zones:** 3A through 7A (Fig. 2)

**Origin:** not native to North America

1. This document is ENH-770, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

**Invasive potential:** little invasive potential

**Texture:** medium

**Uses:** urban tolerant; screen; street without sidewalk; specimen; deck or patio; container or planter; trained as a standard; parking lot island < 100 sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft; sidewalk cutout (tree pit); tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; highway median

### Foliage

**Leaf arrangement:** opposite/subopposite (Fig. 3)

**Leaf type:** simple

**Leaf margin:** entire, undulate

**Leaf shape:** ovate

**Availability:** somewhat available, may have to go out of the region to find the tree



Figure 2. Range

### Description

**Height:** 20 to 30 feet

**Spread:** 20 to 25 feet

**Crown uniformity:** symmetrical

**Crown shape:** upright/erect, round

**Crown density:** dense

**Growth rate:** moderate

**Leaf venation:** pinnate, brachidodrome


**Leaf type and persistence:** deciduous

**Leaf blade length:** 2 to 4 inches, 4 to 8 inches

**Leaf color:** green

**Fall color:** no color change

**Fall characteristic:** not showy

**Flower****Flower color:** white/cream/gray**Flower characteristics:** very showy**Fruit****Fruit shape:** elongated, oval**Fruit length:** .5 to 1 inch**Fruit covering:** dry or hard**Fruit color:** yellow, green**Fruit characteristics:** does not attract wildlife; showy; fruit/leaves not a litter problem**Trunk and Branches****Trunk/bark/branches:** branches don't droop; showy; typically multi-trunked; thorns**Pruning requirement:** little required**Breakage:** resistant**Current year twig color:** brown**Current year twig thickness:** thick**Wood specific gravity:** unknown**Culture****Light requirement:** full sun**Soil tolerances:** clay; sand; loam; slightly alkaline; acidic; well-drained**Drought tolerance:** moderate**Aerosol salt tolerance:** high**Other****Roots:** not a problem**Winter interest:** yes**Outstanding tree:** yes**Ozone sensitivity:** unknown**Verticillium wilt susceptibility:** susceptible**Pest resistance:** resistant to pests/diseases**Figure 3.** Foliage**Use and Management**

It is being used as a street tree in some parts of the country, particularly in areas with overhead power lines. Japanese Tree Lilac is also popular as a garden specimen or as an accent in a shrub border. It deserves to be in any landscape. It provides shade and a colorful spring show for a deck or patio area. Green fruit clusters are somewhat showy when viewed from close range.

The tree is sold as a multi-stemmed specimen or as a single-trunked street tree. The trunk is often trained fairly straight to 10 feet and then it branches into a stiff, upright, rounded head of foliage. The bark is somewhat showy with prominent lenticels, being reminiscent of Black Cherry. As with other Lilacs, the plant as a shrub may need rejuvenation by pruning every few years as it becomes overgrown. It is perhaps the most pest-resistant Lilac, but that does not mean it is pest-free. Regular irrigation during dry spells help make this a pest-resistant tree.

Japanese Tree Lilac is tolerant of urban conditions, growing in poor, clay or alkaline soil. The gorgeous flowers are most showy and prolific when the tree is located in full sun with good drainage. Plants in partial shade can be infected with powdery mildew which can cause some defoliation.

Another available cultivar is 'Ivory Silk' which grows in USDA hardiness zones 3 to 6 into an upright oval with nice flowers which are borne in alternate years.

### **Pests**

If properly located on an appropriate site, there are few problems, but borers can severely affect trees in certain areas.

Lilac borer larvae tunnel in the branches, causing wilting, particularly on drought-stressed trees. Severely infested branches may break off. Remove and destroy infested stems. Keep plants healthy with regular waterings during dry weather and by fertilizing.

Lilac leaf miner tunnels in the leaves in early summer. After mining the leaf, the caterpillars emerge and web leaves together and skeletonize the foliage. Light infestation can be controlled by hand picking.

Scales are most often found infesting the lower stems and often blend in with the bark. Inspect unhealthy-looking plants for scale infestations. Spray with horticultural oil for some control.

### **Diseases**

Usually free of serious disease, but can be affected in certain regions by disease.

Bacterial blight is most serious on white flowered varieties. The young shoots develop black stripes or one side of the shoot turns black. Spots develop on the leaves, forming a water-soaked blotch. Young leaves turn black and die quickly. On older shoots, the spots enlarge more slowly. The flowers wilt and darken. The disease is worse when wet weather occurs as the new shoots are developing. Thin plants to increase air circulation. Remove and

destroy diseased shoots and avoid excessive nitrogen fertilizer.

Phytophthora blight kills stems to the ground. The leaves turn black and shoots have brown lesions on them.

Leaf blotch causes zoned, brown spots. The infected area drops out, leaving a hole in the leaf.

Many fungi cause leaf spots.

Powdery mildew coats the leaves with white powder. During wet weather, Lilacs mildew easily. Mildew is especially severe on shade-grown plants. Ignore late season infections.

Verticillium wilt causes wilting and premature leaf drop. The disease may kill one, several or all the branches. Try fertilizing regularly to help prevent diseases.

Bacterial crown gall causes round, warty galls on the stems near the soil line. Remove infected plants and do not replant in the same spot.