

Taxodium ascendens: Pondcypress¹

Edward F. Gilman and Dennis G. Watson²

Introduction

Similar to Baldcypress in that the trunk is perfectly straight 50 to 60 feet tall, Pondcypress has a narrower crown, is smaller, and has a more open habit. It is found along the edges of streams and around the edge of swampy ground where water is standing; whereas Baldcypress is usually found along stream banks. The bright green, awl-shaped leaves are arranged in an upright row formation along the branches when young, giving a somewhat stiffer and more upright appearance than Baldcypress. The leaves turn an attractive light brown in fall before dropping but the bare branches and light brown, ridged bark provide much landscape interest during the winter. The trunk grows unusually thick toward the base, even on young trees. This is thought to provide support for the tree in its wet habitat. The small seeds are used by some birds and squirrels.

General Information

Scientific name: *Taxodium ascendens*

Pronunciation: tack-SO-dee-um uh-SEN-denz

Common name(s): Pondcypress

Family: *Taxodiaceae*

USDA hardiness zones: 5B through 9B (Fig. 2)

Origin: native to North America

Invasive potential: little invasive potential

Uses: sidewalk cutout (tree pit); tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; street without sidewalk; specimen; reclamation; parking lot island < 100



Figure 1. Middle-aged *Taxodium ascendens*: Pondcypress

sq ft; parking lot island 100-200 sq ft; parking lot island > 200 sq ft

Availability: somewhat available, may have to go out of the region to find the tree

Description

Height: 50 to 60 feet

Spread: 10 to 15 feet

Crown uniformity: symmetrical

Crown shape: pyramidal, upright/erect, columnar

Crown density: open

1. This document is ENH-777, one of a series of the Environmental Horticulture, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date November 1993. Revised December 2006. Reviewed May 2011. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Edward F. Gilman, professor, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.



Figure 2. Range

Growth rate: fast

Texture: fine

Foliage

Leaf arrangement: alternate (Fig. 3)

Leaf type: simple

Leaf margin: entire

Leaf shape: awl-like

Leaf venation: none, or difficult to see

Leaf type and persistence: deciduous

Leaf blade length: less than 2 inches

Leaf color: green

Fall color: copper

Fall characteristic: showy

Flower

Flower color: unknown

Flower characteristics: not showy

Fruit

Fruit shape: round, oval, cone

Fruit length: .5 to 1 inch, 1 to 3 inches

Fruit covering: dry or hard

Fruit color: brown

Fruit characteristics: attracts birds; not showy; fruit/leaves not a litter problem

Trunk and Branches

Trunk/bark/branches: branches don't droop; showy; typically one trunk; thorns

Pruning requirement: little required

Breakage: resistant

Current year twig color: green

Current year twig thickness: thin

Wood specific gravity: unknown

Culture

Light requirement: full sun, partial sun or partial shade

Soil tolerances: clay; sand; loam; slightly alkaline; acidic; extended flooding; well-drained

Drought tolerance: high

Aerosol salt tolerance: moderate

Other

Roots: not a problem

Winter interest: yes

Outstanding tree: yes

Ozone sensitivity: unknown

Verticillium wilt susceptibility: resistant

Pest resistance: resistant to pests/diseases




Figure 3. Foliage

Use and Management

Although often seen at water's edge where it will develop "knees", or root projections, that will extend above the water for gas exchange, Pondcypress can also be grown in dry locations and could make an attractive street tree for a very narrow space. Cypress knees do not generally form on these drier sites. The "knees" do not form as readily as on Baldcypress, even on wet sites. It provides a good vertical accent to the landscape and should be used more often in urban areas. The roots do not appear to lift sidewalks and curbs as readily as some other species. Its delicate foliage affords light, dappled shade, and the heartwood is quite

strong and resistant to rot. However, most lumber available at lumber yards today is sapwood and is not resistant to rot.

Pondcypress is ideal for wet locations, such as its native habitat of stream banks and mucky soils, but the trees will also grow quite well on almost any soil, including clay, silt and sand, except alkaline soils with a pH above 7.5. Its drought-avoidance mechanism allows it to drop leaves in extended dry periods but little harm appears to come to the tree. Pondcypress is relatively maintenance-free, requiring pruning only to remove dead wood and unwanted lower branches which persist on the tree. It maintains a desirably straight trunk without pruning and does not form double or multiple leaders as do many large trees.

The cultivar 'Prairie Sentinel' is narrower than the species.

Propagation is by seed.

Diseases

No diseases are serious.

Twig blight is caused by a weak pathogen and is usually present on dead or dying tissue. When the tree is stressed the fungus can kill branch tips. Dead tips can be pruned off. Do not let dead or diseased branches remain on the tree. Keep trees healthy with regular fertilization.