

# Wild Radish--Biology and Control<sup>1</sup>

J. A. Ferrell, B. Sellers, and G. E. MacDonald<sup>2</sup>

Wild radish is one of the most common and problematic pasture weeds in the Florida Panhandle. It is found throughout the state and can be a serious pest in other crops including peanut, corn and winter vegetables. This publication provides information concerning the biology and growth of wild radish, the problems associated with its presence in wheat and other small grains and methods of control and management.

## Biology

Wild radish (*Raphanus raphanistrum*) is a member of the *Brassicaceae* plant family with cabbage, turnip, and mustard. Most of these species are cool-season plants and have been introduced from eastern Europe and Asia. Wild radish has traditionally been classified as a winter annual. Generally, these species germinate during the fall months when soil temperatures drop below 65° F. Studies indicate a chilling requirement is necessary to break dormancy. In addition, wild radish has a thick fruit pod from which the seed does not shatter free easily. Therefore, the pod must decay before the seed can be released to germinate.

After emergence, wild radish forms a rosette of leaves throughout the winter and early spring (Figure 1). Seedling wild radish possess heart-shaped cotyledons and the first true leaves will be slightly serrated and indented about two to three times as long as wide. As the leaves mature, the

serrations will be more jagged and more deeply indented. In addition, the leaves are covered with stiff hairs, giving a bristly feel to the touch.



Figure 1. Wild Radish (rosette).

The wild radish plant remains in rosette form through most of the winter, reaching approximately 10 to 14 inches across at the base. In the late winter to early spring, as the temperature and day length increase, the plant bolts (Figure 2). Bolting is a process in which the internodes (regions of the stem between leaves) begin to lengthen and a flower stalk forms at the top. In wild radish, multiple flower heads form on several branches arising from a single flower stalk.

1. This document is SS-AGR-236, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published September 2005. Revised November 2010. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. J.A. Ferrell, associate professor, Agronomy Department; B. Sellers, assistant professor; G.E. MacDonald, professor, Agronomy Department; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. Use herbicides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean



Figure 2. Wild Radish (bolt).

The flowers are generally yellow but occasionally may be white.

## Control

### Prevention

The first step in controlling wild radish is prevention. Prevention can be addressed by two strategies. The first is reduction of the seed population in the soil, preventing the weed from infesting an area. This is accomplished by preventing the wild radish from going to seed (i.e., good early season control) and not allowing re-introduction from other sources. For example, sowing winter pastures with certified seed will avoid the possibility of accidentally spreading weed seeds.

The second strategy is to maintain a dense pasture. Wild radish will grow most vigorously where there is a break in the sward. Therefore, proper grazing and soil fertility will likely minimize the impact of this particular weed.

### Mechanical

Mechanical control of wild radish is a consideration, but rarely is it effective. Mowing will generally not harm the basal leaves and allow regrowth to occur within a short period of time. Additionally, a single wild radish plant will often produce many flushes of flowers and set several seed crops. Regular mowing cycles will reduce seed production, but will not eliminate seed production entirely. Frequent mowing will also reduce the productivity and forage yield of the pasture. Therefore, mowing to control wild radish is often more expensive and less effective than other options.

### Chemical

One of the most common and cost effective methods of controlling wild radish is through the use of herbicides.

Some of the most effective and inexpensive herbicides for wild radish control are growth regulators such as 2,4-D and dicamba (Banvel, Clarity, etc.). These herbicides provide excellent control of wild radish (Figure 3) when properly applied. The growth regulating herbicides are generally considered safe on grasses. However, grass crops can be injured if these herbicides are applied incorrectly, or at the wrong developmental stage of the crop. Applications of phenoxy-type herbicides should be made to grain crops after two to three tillers have formed; plants are often 4 to 6 inches tall at this time. Applications of these materials before this stage of growth cause a “rat-tail” effect, whereby the leaf does not form and unfurl properly. In addition, the plant may appear stunted and delayed in maturity. Conversely, applications made after the jointing may result in malformed seedheads. Tolerance of cool season forages to herbicides will vary according to species. Generally, wheat is the most tolerant and oats are least tolerant to 2,4-D applications.




Figure 3. Effect of 2,4-D applied at 1 pt/A on wild radish control as impacted by weed size at time of application.

The timing of an herbicide application is critical for effective wild radish control. Research has shown that >90% wild radish control can be consistently achieved when 2,4-D is applied to plants less than 6 inches in height. By delaying the application until the plant reaches 12 inches, control drops to approximately 70%. However, if wild radish begins to flower before 2,4-D is applied, less than 50% control should be expected. Therefore, herbicides should be applied early to achieve the greatest wild radish control while avoiding herbicide injury to winter forage.

For bermudagrass hay fields, control of wild radish is usually attempted well after flowering and seed development. Control of fully mature plants with 2,4-D can be very difficult. In these situations, metsulfuron (MSM 60, others) at 0.2 oz product/A is most effective. Depending

on temperature at time of application, metsulfuron may require 3 to 5 weeks to control mature wild radish. But this herbicide is highly effective on wild radish and is safe on bermudagrass at any stage from dormant to full greenup.

## Applications with Nitrogen

It is a common practice for many growers to tank mix phenoxy materials with liquid nitrogen to reduce application costs. This practice is acceptable, but certain factors need consideration. The first is the size and density of the wild radish. Nitrogen should be applied near full tiller to maximize nitrogen needed for grain development. If the wild radish density and size is large, waiting this late may result in yield loss, and quite possibly a reduction in control. Applying nitrogen too early to gain a weed control advantage may result in nitrogen deficiency in late season during grain fill. The best rule of thumb is to apply each at its proper time and if these times coincide, a tank mix may be employed. If not, the factors above need to be weighed before such a practice is performed.

If planning to use 2,4-D in combination with nitrogen, herbicide formulation is important to note. The ester formulation of 2,4-D will readily mix with nitrogen solutions while the amine formulation must be mixed with water before adding to the nitrogen solution. Failure to preslurry with water can result in an uneven distribution of herbicide in the spray tank. It should also be noted that spraying the ester formulation of 2,4-D with a nitrogen carrier will often result in greater leaf burning as compared with amine formulation.

## Summary

Wild radish is a common weed problem in winter forages. Managing wild radish depends on several factors. The most important factors are often weed size and crop stage at time of application. Regularly scouting fields will allow you to know the level of weed pressure in each field. Knowing this information will often make the herbicide application decisions much easier. All herbicides applied in grazing areas carry specific grazing restrictions. Please consult the herbicide label, or your local extension office, for more information on specific grazing restrictions.