

Weeds and Epidemiology of Bacterial Leaf Spot of Lettuce in the Everglades Agricultural Area¹

D. C. Odero and G. Sandoya²

Causal Organism and History

Bacterial leaf spot of lettuce, caused by *Xanthomonas campestris* pv. *vitians*, was first reported in the United States in 1918 on head lettuce in New York. In Florida, bacterial leaf spot was first reported in the 1992–93 lettuce growing season. All major types of lettuce (crisphead, butterhead, and leaf) were affected, but the disease was more severe in romaine lettuce. So far the disease has not been observed on endive lettuce.

Symptoms

Symptoms of bacterial leaf spot are black, angular, water-soaked lesions that occur primarily on mature, fully expanded leaves (Figure 1). These lesions coalesce as the disease develops, resulting in large necrotic areas and collapse of the leaf. Occasionally, the pathogen may also infect stem tissue, causing stem rot, stunting, and collapse of young plants.

Weeds and Disease Epidemiology

Infected weeds and epiphytic populations on weeds growing in close proximity to lettuce can be possible sources



Figure 1. Bacterial leaf spot of lettuce.

Credits: Nikol Havranek, UF/IFAS

of initial *Xanthomonas campestris* pv. *vitians* inoculum. *Xanthomonas campestris* pv. *vitians* causes bacterial leaf spot symptoms in broadleaf weeds, including prickly lettuce, trumpet firewood, annual sowthistle, field bindweed, panicle willoweed, shepherd's purse, pineapple weed, netleaf goosefoot, common knotweed, little mallow, and common groundsel. However, it does not cause any bacterial leaf spot symptoms on grass weeds. The most common broadleaf

1. This document is SS-AGR-347, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date May 2011. Reviewed July 2014 and September 2018. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. D. C. Odero, associate professor, Agronomy Department, Everglades REC—Belle Glade, FL, Florida; and G. Sandoya, assistant professor, Horticultural Sciences Department, Everglades REC—Belle Glade, FL, Florida. Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition. All chemicals should be used in accordance with directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 2. Spiny amaranth 4 weeks after inoculation with *Xanthomonas campestris* pv. *vitians* strain.
Credits: Nikol Havranek, UF/IFAS



Figure 4. Common purslane 4 weeks after inoculation with *Xanthomonas campestris* pv. *vitians* strain.
Credits: Nikol Havranek, UF/IFAS



Figure 3. Common lambsquarters 4 weeks after inoculation with *Xanthomonas campestris* pv. *vitians* strain.
Credits: Nikol Havranek, UF/IFAS




Figure 5. Growth of *Xanthomonas campestris* pv. *vitians* following streaking with inoculum from common purslane (top left petri dish), spiny amaranth (top right petri dish), common lambsquarters (bottom left petri dish), and lettuce (bottom right petri dish) previously inoculated with *Xanthomonas campestris* pv. *vitians* strain (lettuce showing and confirming *Xanthomonas campestris* pv. *vitians* growth while the weeds show no growth).
Credits: Nikol Havranek, UF/IFAS

weed species in and around lettuce fields in the Everglades Agricultural Area (EAA) include common lambsquarters, *Amaranthus* spp., common purslane, common ragweed, ragweed parthenium, and horse purslane. *Xanthomonas campestris* pv. *vitians* or its epiphytes do not cause any symptoms on these weed species (Figure 2, 3, and 4). Similarly, no growth of *Xanthomonas campestris* pv. *vitians* occurs on glucose nutrient agar following streaking with inoculum from weeds inoculated with *Xanthomonas campestris* pv. *vitians* strains (Figure 5). Thus, these weed species may not be sources of *Xanthomonas campestris* pv. *vitians* inoculum around lettuce fields in the EAA. However, further studies that are more comprehensive should be conducted to corroborate these observations. Nonetheless, control programs for these weed species in and around lettuce fields should be practiced to forestall other negative effects on production.

References

- Barak, J. D., S. T. Koike, and R. L. Gilbertson. 2001. "Role of crop debris and weeds in the epidemiology of bacterial leaf spot of lettuce in California." *Plant Disease* 85: 169-78.
- Davis, R. M., K. V. Subbarao, R. N. Raid, and E. A. Kurtz. 1997. *Compendium of Lettuce Diseases*. St. Paul: APS Press.
- Pernezny, K., R. N. Raid, R. E. Stall, N. C. Hodge, and J. Collins. 1995. "An outbreak of bacterial spot of lettuce in Florida caused by *Xanthomonas campestris* pv. *vitians*." *Plant Disease* 79: 359-60.