

Seed Quality and Seeding Technology¹

Shinsuke Agehara and Bielinski M. Santos²

Seed

A seed is an encapsulated plant embryo developed from a fertilized ovule. Seed quality plays an important role in the production of agronomic and horticultural crops. High quality seeds are genetically and varietally pure with a high germination percentage, free from disease and disease organisms, and with a proper moisture content and weight.

High quality seeds insure good germination, rapid emergence, and vigorous growth. These aspects translate to a good stand (whether greenhouse or field). Poor quality seeds result in “skips,” excessive thinning, or yield reductions due to overcrowding, all of which diminish profitability.

Vigor is often implied when discussing seed quality, and most growers have come to use the terms “quality” and “vigor” interchangeably. “Vigor” is defined as those properties that determine the potential performance of seed during germination and establishment. Seed vigor is generally related to yield in vegetables. Therefore, high-vigor seed should be used in all instances to ensure good stand establishment under varying field conditions.

Different seed lots sown in the same environment may act differently, and seedbed factors (temperature, water

content, etc.) strongly affect seed performance. Fluctuations (airspace, moisture, temperature, etc.) in the seedbed environment are the most important factors in determining final seedling emergence. Furthermore, large seeds of a particular species frequently produce more vigorous plants and uniform stands than small seeds.

Most seeds are fairly hardy; however, seeds in the bean and pea family (snap, lima, southern pea, English pea, etc.) are fragile and should be handled with extreme care. Dropping these seeds from any height while loading or unloading or pouring into seed hoppers will crack their seed coats and decrease germination.

National and international seed companies strive to provide high-quality vegetable seed through various milling processes and stringent disease screening. These techniques reduce the total tonnage of raw seed but increase the overall quality. Federal minimum germination standards regulate the seed industry (Table 1). However, most companies attempt to exceed these minimum standards.

Organic Seed and Sources

Seed used for organic vegetable production must meet specifications of the USDA's National Organic Program. Recordkeeping is required for growers during all phases

1. This document is HS713, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date June 1995. Revised July 1996, September 2001, December 2005, September 2007, and March 2017 (minor revisions by S. Agehara). Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Shinsuke Agehara, assistant professor; and Bielinski M. Santos, former associate professor; Horticultural Sciences Department, UF/IFAS Gulf Coast Research and Education Center, Wimauma, FL 33598.

The use of trade names in this publication is solely for the purpose of providing specific information. It is not a guarantee or warranty of the products named, and does not signify that they are approved to the exclusion of others of suitable composition. Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

of organic production, as is certification by an accredited certifying agency. Use of conventionally produced seed is allowed only when preferred cultivars are not available from organic sources. In no case are seeds treated with non-approved substances, or seeds of genetically modified organisms or derived from artificial seed technologies, allowed in organic production.

Primed Seed

Improved seed quality and vigor may be obtained with primed or enhanced seed. Primed seed is biologically altered through the addition of just enough water (and often some hormones) to allow the seed to undergo the early stages of germination without actually germinating. In this “primed” state, seed will germinate more rapidly and emerge more uniformly over a greater range of temperatures and soil moisture conditions. This translates to greater seedling vigor, uniformity, and rapid establishment.

Primed seed is best suited for direct seeding where adverse conditions may prevail during germination and emergence. However, enhanced seed use in the greenhouse is also cost effective. Uniformity of emergence alleviates “playing catch-up” within a flat due to uneven emergence. Rapid emergence means production programs (fertilizer, pesticides, etc.) can be quickly implemented. The reduction in wasted space from poor germination and in labor for thinning overseeded flats increases production efficiency. Greater efficiency means greater profitability.

Seed Storage

Care must be taken with seed to avoid injury and provide proper storage. Unused seed is best stored if the hermetic seal has not been broken. However, if open cans are to be stored, it is best to seal these cans in ziplock bags and store at 40°F–50°F preferably in a refrigerator used solely for seed storage. High temperatures and humidity are very harmful to seed. Humid conditions lead to increases in seed moisture, which reduce shelf life. Under proper storage conditions, vegetable seeds can remain viable for several years. The relative life expectancy of vegetable seed is presented in Table 2.

Field Seeding

Prior to seeding, the field should be leveled and worked to obtain a fine textured soil, free of clods and debris. Seedbed firmness, depth of planting, and available moisture are important considerations. Treated seeds should be used for protection against soil-borne diseases and insects during germination and seedling development. New seed

film-coating techniques apply fungicides and pesticides in “no dust” formulations, which decrease the potential for worker injury by reducing absorption and inhalation risk.

See Table 3 for the number of seeds per unit weight for the individual crops.

Direct seeding can be accomplished with a variety of planters. The particular type of planter used will depend on the grower’s preference, field conditions, equipment, and experience. Tractor speed is the key to success or failure of many stands. Lower speeds reduce injury to seeds (especially bean and pea) as they pass through the planter. A planting speed of 2–3 miles per hour is preferable.

In precision seeding, a single seed is planted at an exact plant spacing to achieve a uniform stand. Precision seeders vary in their approach to seed singulation (punched belts, vacuum plates, cups on armatures, etc.), and, again, choice depends on grower preference, seed to be planted, and seedbed conditions. See Table 4 for the number of plants per acre at various row spacings.

Precision seeding requires uniformity in seed size both between species and within species. For example, large seed (snap bean) cannot be planted with the same belt/plate/cup as small seed (broccoli). Within species, two small seeds may be picked up where only one seed should fall. Sizing within seed lots may be accomplished during commercial separation or through pelletizing techniques. Pelletizing, where a seed is encased in a material that can be formed to produce a uniformly sized pellet, has improved over the years to reduce the occurrence of slow and erratic germination compared to raw seed.

While pelletized seed is required by some precision seeders, many planters are capable of planting raw seed. Sized raw seed used in these planters has been an advantage for some growers. Plants from large seed will emerge first and grow faster than from small seed.

Table 1. Minimum official federal germination standards.¹

| Seed | (%) | Seed | (%) | Seed | (%) |
|------------------|-----|-----------------------|-----|----------------------|-----|
| Artichoke | 60 | Corn, Sweet | 75 | Onion | 70 |
| Asparagus | 70 | Corn salad | 70 | Onion, Welsh | 70 |
| Bean, Asparagus | 75 | Cowpea (Southern pea) | 75 | Pak-choi | 75 |
| Bean, Broad | 75 | Cress, Garden | 75 | Parsley | 60 |
| Bean, Garden | 70 | Cress, Upland | 60 | Parsnip | 60 |
| Bean, Lima | 70 | Cress, Water | 40 | Pea | 80 |
| Bean, Runner | 75 | Cucumber | 80 | Pepper | 55 |
| Beet | 65 | Dandelion | 60 | Pumpkin | 75 |
| Broccoli | 75 | Dill | 60 | Radish | 75 |
| Brussels sprouts | 70 | Eggplant | 60 | Rhubarb | 60 |
| Cabbage | 75 | Endive | 70 | Rutabaga | 75 |
| Cardoon | 60 | Kale | 75 | Sage | 60 |
| Carrot | 55 | Kohlrabi | 75 | Salsify | 75 |
| Cauliflower | 75 | Leek | 60 | Sorrel | 65 |
| Celery/celeriac | 55 | Lettuce | 80 | Spinach | 60 |
| Chard, Swiss | 65 | Muskmelon | 75 | Spinach, New Zealand | 40 |
| Chicory | 65 | Mustard | 75 | Squash | 75 |
| Chinese cabbage | 75 | Mustard, Spinach | 75 | Tomato | 75 |
| Chives | 50 | Mustard, Vegetables | 75 | Tomato, Husk | 50 |
| Citron | 65 | Okra | 50 | Turnip | 80 |

¹Adapted from Donald N. Maynard and George J. Hochmuth, *Knott's Handbook for Vegetable Growers*, 5th Edition (2006). Reprinted by permission of John Wiley & Sons.

Table 2. Approximate life expectancy of vegetable seeds stored under favorable conditions.¹

| Vegetable | Years | Vegetable | Years | Vegetable | Years |
|------------------|-------|---------------|-------|----------------------|-------|
| Asparagus | 3 | Corn salad | 5 | Parsnip | 1 |
| Bean | 3 | Cress, Garden | 5 | Pea | 3 |
| Beet | 4 | Cress, Water | 5 | Pepper | 2 |
| Broccoli | 3 | Cucumber | 5 | Pumpkin | 4 |
| Brussels sprouts | 4 | Dandelion | 2 | Radish | 5 |
| Cabbage | 4 | Eggplant | 4 | Roselle | 3 |
| Cardoon | 5 | Endive | 5 | Rutabaga | 4 |
| Carrot | 3 | Fennel | 4 | Salsify | 1 |
| Cauliflower | 4 | Kale | 4 | Scorzonera | 2 |
| Celeriac | 3 | Kohlrabi | 3 | Sea kale | 1 |
| Celery | 3 | Leek | 2 | Sorrel | 4 |
| Chard, Swiss | 4 | Lettuce | 6 | Southern pea | 3 |
| Chervil | 3 | Martynia | 2 | Spinach | 3 |
| Chicory | 4 | Muskmelon | 5 | Spinach, New Zealand | 3 |
| Chinese cabbage | 3 | Mustard | 4 | Squash | 4 |
| Ciboule | 2 | Okra | 2 | Tomato | 4 |
| Collards | 5 | Onion | 1 | Turnip | 4 |
| Corn, sweet | 2 | Parsley | 1 | Watermelon | 4 |

¹Adapted from Donald N. Maynard and George J. Hochmuth, *Knott's Handbook for Vegetable Growers*, 5th Edition (2006). Reprinted by permission of John Wiley & Sons

Table 3. Number of seeds per unit weight.

| Crop | Seeds/unit weight | Crop | Seeds/unit weight |
|------------------|--------------------------|-------------------------|--------------------------|
| Asparagus | 13,000–20,000/lb | Onion | |
| Bean | | Bulb | 105,000–144,000/lb |
| Baby lima | 1,150–1,450/lb | Bunching | 180,000–200,000/lb |
| Fordhook lima | 440–550/lb | Parsley | 240,000–288,000/lb |
| Snap | 1,600–2,200/lb | Parsnip | 7,500–12,000/lb |
| Beet | 24,000–26,000/lb | Pea | 1,440–2,580/lb |
| Broccoli | 8,500–9,000/oz | Pepper | 4,000–4,700/oz |
| Brussels sprouts | 8,500–9,000/oz | Pumpkin | 1,900–3,200/lb |
| Cabbage | 8,500–9,000/oz | Radish | 40,000–50,000/lb |
| Carrot | 300,000–400,000/lb | Rutabaga | 150,000–192,000/lb |
| Cauliflower | 8,900–10,000/oz | Southern pea | 3,000–4,200/lb |
| Celery | 60,000–72,000/oz | Spinach | 40,000–50,000/lb |
| Collards | 7,500–8,500/oz | Squash | |
| Cucumber | 15,000–16,000/lb | Summer | 3,500–4,800/lb |
| Eggplant | 6,000–6,500/oz | Winter | 1,600–4,000/lb |
| Endive, Escarole | 22,000–26,000/oz | Sweet corn | |
| Kale | 7,500–8,900/oz | Normal, sugary enhanced | 1,800–2,500/lb |
| Leek | 170,000–180,000/lb | Supersweet | 3,000–5,000/lb |
| Lettuce | | Tomato | |
| Leaf | 25,000–31,000/oz | Fresh | 10,000–11,400/oz |
| Head | 20,000–25,000/oz | Processing | 160,000–190,000/lb |
| Muskmelon | 16,000–19,000/lb | Turnip | 15,000/oz |
| Mustard | 15,000–17,000/oz | Watermelon | |
| Okra | 450–550/oz | Small seed | 8,000–10,400/lb |
| | | Large seed | 3,200–4,800/lb |

Table 4. Plants per acre at various between and in-row spacings.

| Distance between rows (inches) | In-row spacing (inches) | | | | | | | | | | | | |
|--------------------------------|---------------------------|---------|---------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | 2 | 4 | 6 | 8 | 10 | 12 | 14 | 16 | 18 | 24 | 30 | 36 | 48 |
| | Number of plants per acre | | | | | | | | | | | | |
| 7 | 448,046 | 224,022 | 149,348 | 112,011 | 89,609 | 74,674 | 64,006 | | | | | | |
| 12 | 261,360 | 130,680 | 87,120 | 64,340 | 52,272 | 43,560 | 37,337 | 32,670 | 29,040 | 21,780 | 17,424 | 14,520 | 10,890 |
| 18 | 174,240 | 87,911 | 58,080 | 43,560 | 34,848 | 29,040 | 24,891 | 21,780 | 19,360 | 14,520 | 11,616 | 9,680 | 7,260 |
| 21 | 149,354 | 74,675 | 49,782 | 37,337 | 29,870 | 24,891 | 21,335 | 18,669 | 16,594 | 12,446 | 9,957 | 8,297 | 6,223 |
| 24 | 130,860 | 65,405 | 43,560 | 32,670 | 26,136 | 20,908 | 17,424 | 16,335 | 15,520 | 10,890 | 8,712 | 7,260 | 5,445 |
| 30 | 104,544 | 52,272 | 34,848 | 26,136 | 20,908 | 17,424 | 14,934 | 13,068 | 11,616 | 8,712 | 6,970 | 5,808 | 4,356 |
| 36 | 87,120 | 43,560 | 29,040 | 21,780 | 17,424 | 14,520 | 12,445 | 10,890 | 9,680 | 7,260 | 5,808 | 4,840 | 3,630 |
| 42 | 74,674 | 37,337 | 24,891 | 18,668 | 14,934 | 12,446 | 10,667 | 9,334 | 8,297 | 6,223 | 4,978 | 4,148 | 3,111 |
| 48 | 65,340 | 32,673 | 21,780 | 16,335 | 13,068 | 10,890 | 9,334 | 8,167 | 7,790 | 5,445 | 4,356 | 3,630 | 2,723 |
| 60 | | | 17,424 | 13,068 | 10,538 | 8,712 | 7,467 | 6,534 | 5,808 | 4,356 | 3,484 | 2,901 | 2,178 |
| 72 | | | 14,520 | 10,890 | 8,712 | 7,260 | 6,223 | 5,445 | 4,840 | 3,630 | 2,904 | 2,420 | 1,815 |
| 84 | | | 12,455 | 9,334 | 7,467 | 6,222 | 5,334 | 4,667 | 4,148 | 3,111 | 2,489 | 2,074 | 1,555 |
| 96 | | | 10,890 | 8,167 | 6,534 | 5,445 | 4,667 | 4,084 | 3,630 | 2,722 | 2,178 | 1,815 | 1,361 |