

# Chapter 19. Biopesticides and Alternative Disease and Pest Management Products<sup>1</sup>

Natalia Peres, Gary Vallad, Johan Desaeger, and Hugh Smith<sup>2</sup>

Table 1. Biopesticides labeled for management of arthropod pests of many crops.

Trade Name (Active Ingredient)	MOA Code <sup>1</sup>	Insect or Mite Pest	Days to Harvest	REI (Hours)	OMRI-Listed	Remarks
<b>Agree WG, Xentari</b> ( <i>Bacillus thuringiensis</i> subspecies <i>aizawai</i> ) <b>Biobit HP, Deliver, DiPel DF, Javelin WG</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	11A	Caterpillars	0	4	Yes	Apply when larvae are small for best control. Can be used in greenhouse.
<b>Aza-Direct</b> (azadirachtin)	-	Aphids, beetles, caterpillars, leafhoppers, leafminers, thrips, whiteflies	0	4	Yes	Antifeedant, repellent, insect growth regulator.
<b>Azera</b> (pyrethrins + azadirachtin)	3A + -	Aphids, beetles, caterpillars, leafminers, whiteflies	0	12	Yes	Apply when larvae are small for best control. Can be used in greenhouse.
<b>BotaniGard ES</b> ( <i>Beauveria bassiana</i> strain GHA)	-	Aphids, leafhoppers, mealybugs, mites, plant bugs, planthoppers, thrips, whiteflies	0	4	No	Works by contact. Spores attach to insects, germinate, and penetrate insect cuticle. May be used in greenhouses.
<b>BotaniGard Maxx</b> ( <i>Beauveria bassiana</i> strain GHA + pyrethrins)	3A + -	Aphids, leafhoppers, leafminers, plant bugs, thrips, whiteflies	0	12	No	
<b>Captiva</b> ( <i>Capsicum oleoresin</i> extract, garlic oil, soybean oil)	-	Leafhoppers, mites, thrips, whiteflies	0	4	No	

1. This document is CV295. Original publication date August 2013. Revised May 2021. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Natalia Peres, professor, Plant Pathology Department, UF/IFAS Gulf Coast Research and Education Center; Gary E. Vallad, professor, Plant Pathology Department, UF/IFAS Gulf Coast REC; Johan Desaeger, assistant professor, Entomology and Nematology Department, UF/IFAS Gulf Coast REC; and Hugh A. Smith, associate professor, Entomology and Nematology Department, UF/IFAS Gulf Coast REC; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Trade Name (Active Ingredient)	MOA Code <sup>1</sup>	Insect or Mite Pest	Days to Harvest	REI (Hours)	OMRI- Listed	Remarks
<b>Captiva Prime</b> ( <i>Capsicum oleoresin</i> extract, garlic oil, canola oil)	-	Leafhoppers, mites, thrips, whiteflies	0	4	Yes	
<b>Des-X</b> (soap, insecticidal)	-	Aphids, mites, plant bugs, whiteflies	0	12	Yes	
<b>Entrust SC</b> (spinosad)	5	Beetles, caterpillars, thrips	See label	4	Yes	See label for resistance management and for maximum allowed amount per season.
<b>Grandevo WDG</b> ( <i>Chromobacterium</i> <i>subtsugae</i> strain PRAA4- 1)	-	Aphids, mites, plant bugs, whiteflies	0	4	Yes	
<b>JMS Stylet oil</b> (paraffinic oil)	-	Aphids, leafhoppers, leafminers, mites, whiteflies	0	4	No	
<b>Met52 EC</b> ( <i>Metarhizium anisopliae</i> strain F52)		Mites, thrips	0	12	Yes	Composed of spores of insect pathogenic fungus <i>Metarhizium anisopliae</i> strain F52.
<b>M-Pede 49% EC</b> (soap, insecticidal)	-	Aphids, leafhoppers, mites, thrips, whiteflies	0	12	Yes	
<b>Mycotrol ESO</b> ( <i>Beauveria bassiana</i> strain GHA)	-	Aphids, leafhoppers, whiteflies	0	4	Yes	
<b>Mycotrol WPO</b> ( <i>Beauveria bassiana</i> strain GHA)	-	Aphids, leafhoppers, whiteflies	0	4	Yes	
<b>Neemix 4.5</b> (azadirachtin)	-	Aphids, leafminers, whiteflies	0	12	Yes	IGR and feeding repellent. Greenhouse and field.
<b>Organic JMS Stylet Oil</b> (paraffinic oil)	-	Aphids, leafhoppers, leafminers, mites, whiteflies	0	4	Yes	
<b>PFR-97 20% WDG</b> ( <i>Isaria fumosoroseus</i> Apopka Strain 97)	-	Aphids, leafminers, mites, plant bugs, psyllids, weevils, whiteflies	0	4	Yes	Can be used on vegetables grown for transplant. Do not mix with fungicides or apply within 5 days of fungicide applications other than copper. Dust/mist respirator must be used for mixing and applying.
<b>Pyganic 5.0</b> (pyrethrins)	3A	Aphids, beetles, leafhoppers, leafminers, mealybugs, mites, plant bugs, planthoppers, psyllids, stink bugs, thrips, weevils, whiteflies	0	12	Yes	Harmful to bees. Can be used in greenhouses.
<b>Seduce Insect Bait</b> (spinosad)	5	Caterpillars	See specific crops on label	4	Yes	
<b>Suffoil-X</b> (mineral oil)	-	Aphids, leafhoppers, thrips, whiteflies	0	4	Yes	Thorough coverage is essential.
<b>Trilogy</b> (extract of neem oil)	-	Aphids, mites, thrips, whiteflies	0	4	Yes	Apply morning or evening to reduce potential for leaf burn. Toxic to bees exposed to direct treatment.

Trade Name (Active Ingredient)	MOA Code <sup>1</sup>	Insect or Mite Pest	Days to Harvest	REI (Hours)	OMRI- Listed	Remarks
<b>Venerate XC</b> (heat-killed <i>Burkholderia</i> spp. strain A396 cells and spent fermentation media)	-	Corn earworm, mealybugs, mites, stink bugs, weevils, whiteflies	0	4	Yes	

<sup>1</sup> Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 8.2 March 2020 Number codes (1 through 29) are used to distinguish the main insecticide mode of action groups, with additional letters for certain subgroups within each main group. All insecticides within the same group (with same number) indicate the same active ingredient or similar mode of action. This information must be considered for the insecticide resistance management decisions. - = unknown, or a mode of action that has not been classified yet.

Table 2. Biopesticides and other alternative products labeled for plant disease management.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
Labels change frequently. Be sure to read a current product label before applying any pesticide. Refer to crop-specific tables for conventional pesticides labeled for disease and pest management.						
<b>ActinoGrow</b> <b>Actinovate</b> ( <i>Streptomyces lydicus</i> WYEC 108), NC	Vegetables & Strawberries	Broad-spectrum fungicide and bactericide	0	1 hr	Yes	See label for specific rates and application recommendations.
<b>Afla-guard GR</b> ( <i>Aspergillus flavus</i> NRRL 21882), NC	Sweet Corn	Aspergillus	-	4 hr	No	Do not exceed 20 lb/acre per growing season. See label for specific application instructions.
<b>AgriPhage</b> (bacteriophage), NC	Tomato, Pepper	Bacterial spot, Bacterial speck	0	0	No	Bacterial strains must be characterized periodically by manufacturer to correctly formulate the bacteriophage mixture.
<b>Aviv</b> ( <i>Bacillus subtilis</i> strain IAB/BS03)	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	Yes	See label for specific rates and application recommendations.
( <i>Bacillus subtilis</i> strain QST 713), 44 <b>Many brands available:</b> Minuet, Rhapsody, Serenade ASO, Serenade Opti,	Vegetables & Strawberries	Broad-spectrum fungicide and bactericide	0	4 hr	Yes	For foliar applications, mix with copper compounds or other effective fungicides for improved disease control. See label for details regarding specific foliar and soil applications and target diseases.
<b>BioCover</b> (Oil, petroleum), NC	Vegetables & Strawberries	Powdery mildew, Rust	0	4 hr	No	See label for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>BIO-TAM</b> ( <i>Trichoderma asperellum</i> strain ICC 012 + <i>Trichoderma gamsii</i> strain ICC 080), NC	Vegetables & Strawberries	<i>Fusarium</i> spp., <i>Phytophthora</i> spp., <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., <i>Sclerotinia</i> spp., <i>Sclerotium</i> <i>rolfsii</i> , <i>Thielaviopsis basicola</i> , and <i>Verticillium</i> spp.	-	1 hr	Yes	See label for additional rates and recommendations for transplant production and details for specific diseases. Check label for product incompatibility with certain chemical fungicides.
<b>Botector</b> ( <i>Aureobasidium pullulans</i> strain DSM 14940, 14941)	Tomatoes & Strawberries	Anthracnose, Botrytis, Phomopsis, Rhizopus	0	4 hr	Yes	See label for specific rates and application recommendations. Check label for precautions regarding incompatibility with fungicides.
<b>BotryStop</b> ( <i>Ulocladium oudemansii</i> U3 strain), NC	Vegetables & Strawberries	Botrytis, Sclerotinia	0	4 hr	No	See label for specific rates, application recommendations, and precautions regarding fungicide compatibility.
<b>Cinnerate</b> (cinnamon oil), NC	Vegetables & Strawberries	Powdery mildew, Botrytis, Rusts	0	0	Yes	See label for specific rates and application recommendations.
<b>Contans WG</b> ( <i>Coniothyrium minitans</i> strain CON/M/91-08)	Vegetables & Strawberries	<i>Sclerotinia sclerotiorum</i> and <i>Sclerotinia minor</i>	0	4 hr	Yes	See label for specific rates and application recommendations.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
<b>Cyclops</b> (olive and cinnamon oils), NC	Vegetables & Strawberries	Broad-spectrum fungicide	0	0	Yes	See label for specific rates and application recommendations. Ingredients are exempt from FIFRA.
<b>Debug ON</b> <b>Debug Optimo</b> <b>Debug Tres</b> <b>Debug Turbo</b> (fats and glyceridic oils Mongosa, Azadirachtin), NC	Vegetables & Strawberries	Powdery mildew, Rust, <i>Rhizoctonia</i> , <i>Sclerotinia</i> , <i>Sclerotium rolfsii</i>	0	4 hr	Yes	See labels for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>Double Nickel 55</b> <b>Double Nickel LC</b> <b>Triathlon BA</b> ( <i>Bacillus amyloliquifaciens</i> strain D747), 44	Vegetables & Strawberries	<i>Alternaria</i> spp., Anthracnose, Bacterial diseases, Botrytis, Early blight, <i>Fusarium</i> spp., Late blight, <i>Phytophthora</i> spp., Powdery mildew, <i>Pythium</i> spp., <i>Rhizoctonia</i>	0	4 hr	Yes	See label for additional rates and recommendations for foliar and soil application rates and details for specific diseases. Use as a soil drench at transplant and periodically throughout the season. Can also be used as a seed treatment. See label for details.
<b>EcoSwing</b> (extract of <i>Swinglea glutinosa</i> ), NC	Vegetables & Strawberries	<i>Alternaria alternata</i> , <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> , <i>Botrytis cinerea</i> , <i>Erysiphaceae</i> spp., <i>Erwinia carotovora</i> subspp., <i>Aspergillus niger</i> , <i>Alternaria tenuis</i> , <i>Cladosporium herbarum</i> , <i>Rhizopus arrhizus</i> , <i>Penicillium</i> spp.	0	4 hr	Yes	See label for specific rates, application recommendations, and greenhouse use.
<b>Glacial Spray Fluid</b> (Oil, petroleum), NC	Vegetables & Strawberries	Powdery mildew, Rust	0	4 hr	Yes	See label for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>GreenClean PRO</b> <b>PerCarb</b> (sodium carbonate peroxyhydrate), NC	Vegetables & Strawberries	Broad-spectrum fungicide/bactericide	0	0	No	See label for specific rates and application recommendations.
<b>Howler</b> ( <i>Pseudomonas chlororaphis</i> strain AFS009), NC	Vegetables & Strawberries	<i>Rhizoctonia</i> , <i>Pythium</i> , <i>Fusarium</i> , <i>Phytophthora</i> , <i>Sclerotinia</i> , <i>Colletotrichum</i> , and <i>Botrytis</i> .	0	4 hr	Yes	See label for specific rates and application recommendations.
(Hydrogen peroxide + peroxyacetic acid), NC <b>Many brands available:</b> GreenClean Liquid, Jet-Ag, OxiDate, Purpose Plus, Rendition, SaniDate, SporeQuell, StorOx, TerraClean, TerraStart, ZeroTol	Vegetables & Strawberries	Broad-spectrum fungicide	0	1 hr for enclosed areas; until spray dries in open-field areas.	No	See labels for specific rates, application recommendations, and precautions regarding use with other pesticides. Some formulations can be used as a soil drench at transplant and periodically throughout the season. Can also be used as a seed treatment or for postharvest treatments.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
<b>JMS Stylet-Oil</b> <b>Organic JMS Stylet-Oil</b> (paraffinic oil), NC	Vegetables & Strawberries	Potato Virus Y, Tobacco Etch Virus, Cucumber Mosaic Virus	0	4 hr	Yes, but only for one label	See label for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>LEAPES</b> ( <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> , strain ABTS-351 and methyl salicylate)	Tomato & Pepper	<i>Xanthomonas</i> spp.	0	0.5	No	Apply before disease is observed in the field; should be used as part of a plant pathogen control program. See label for details regarding insecticidal activity.
<b>LifeGard</b> ( <i>Bacillus mycoloides</i> isolate J)	Vegetables	Broad-spectrum fungicide		4 hr	Yes	See label for specific rates and application recommendations.
<b>Mildew Cure</b> (cotton, corn, and garlic oils), NC	Vegetables & Strawberries	Powdery mildew	0	0	Yes	See label for specific rates and application recommendations. Ingredients are exempt from FIFRA.
(mineral oil), NC <b>Many brands:</b> 440 Superior Spray Oil, BioCover, Glacial Spray Oil, Omni Supreme Oil, PureSpray Green, SuffOil-X, TriTek, Ultra- Pure Oil	Vegetables & Strawberries	Powdery mildew	0	4 hr	Yes	See label for specific rates, application recommendations, precautions, and fungicide compatibility.
<b>Mycostop</b> ( <i>Streptomyces</i> sp. strain K61)	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	No	See label for specific rates and application recommendations.
<b>Nutrol</b> (potassium phosphate), NC	Cucurbits, Peppers & Tomatoes	Powdery mildew	0	0	No	Do not mix with copper fungicides or other products affected by low pH (<5.5). See label for specific rates and application recommendations.
<b>OxiPhos</b> (mono- and dipotassium salts of phosphorus acid + hydrogen peroxide), 33 & NC	Vegetables & Strawberries	<i>Alternaria</i> spp., Anthracnose, Bacterial diseases, Botrytis, Early blight, <i>Fusarium</i> spp., Late blight, <i>Phytophthora</i> spp., Powdery mildew, <i>Pythium</i> spp., Rhizoctonia	0	1 hr for enclosed areas; until spray dries in open-field areas.	No	See label for additional rates and recommendations for transplant production and details for specific diseases. Use as a soil drench at transplant and periodically throughout the season. Can also be used as a seed treatment.
<b>PlantShield HC</b> ( <i>Trichoderma harzianum</i> Rifai strain KRL-AG2), NC	Vegetables & Strawberries	<i>Fusarium</i> spp., <i>Pythium</i> spp., Rhizoctonia	0	4 hr	Yes	Can be applied to plant as a direct drench, furrow spray, chemigation, or in transplant starter solution. See label for details.
(potassium bicarbonate), NC <b>Many brands available:</b> Armcarb 100, Armicarb "O", Carb-O-Nator, Eco- mate, Kaligreen, Millstop	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	Yes	See label for specific rates and application recommendations.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
(potassium phosphite; mono- and dipotassium salts of phosphorous acid), 33 <b>Many brands available:</b> Alude, Apear, Confine Extra T&O, Fosphite, Fungi-Phite, Helena Prophyt, K-Phite 7LP AG, Phiticide, Phorcephite, Phostrol, Rampart, Reveille	Vegetables & Strawberries	<i>Alternaria</i> spp., Anthracnose, Bacterial diseases; Downy mildew, <i>Fusarium</i> spp., Late blight, Leaf blights caused by <i>Cercospora</i> and <i>Septoria</i> spp., <i>Phytophthora</i> spp., Powdery mildew, <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., Root rots	0	4 hr	No	See label for details, specific recommendations, and precautions for tank mixing with copper-based fungicides.
<b>Prev-Am</b> (sodium tetraborohydrate decahydrate), NC	Vegetables & Strawberries	Downy mildew, Botrytis, Powdery mildew, Late blight, White rust, White mold	0	0.5	No	See label for details, crop-specific recommendations, and precautions.
<b>ProBlad Verde</b> (Banda de Lupinus albus doce (BLAD))	Vegetables & Strawberries	Anthracnose, Botrytis, Powdery mildew, Southern blight, White mold	1	4 hr	Yes	See label for details, crop specific recommendations, and precautions.
<b>Procidic, BerryCare</b> (citric acid), NC	Vegetables & Strawberries	Broad-spectrum fungicide	0	0	No	See label for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>Purespray Green</b> (oil, petroleum), NC	Vegetables & Strawberries	Powdery mildew, Rust	0	4 hr	Yes	See label for specific rates, application recommendations, and precautions regarding use.
<b>Pvent</b> <b>LALSTOP G46 WG</b> ( <i>Gliocladium catenulatum</i> strain J1446), NC	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	Yes	See label for details, crop specific recommendations, and precautions.
<b>Regalia</b> <b>Regalia Rx</b> <b>Pacesetter</b> (extract of <i>Reynoutria sachalinensis</i> ), P	Vegetables & Strawberries	Bacterial canker, Bacterial speck, Bacterial spot, Botrytis, Early blight, Late blight, <i>Phytophthora</i> spp., Powdery mildew, Target spot	0	4 hr	Yes	Tank-mix with other effective fungicides for improved disease control under heavy pressure. See label for details.
<b>Romeo</b> (Cerevisane; cell walls of <i>Saccharomyces cerevisiae</i> strain LAS117)	Vegetables & Strawberries	<i>Alternaria</i> leaf spot, Anthracnose, Botrytis, Downy mildew, Fusarium, Late blight, <i>Phytophthora</i> , Powdery mildew, <i>Rhizoctonia</i> , Scab, Sour rot	0	4 hr	Yes	See label for details, crop specific recommendations, and precautions.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
<b>RootShield Granular</b> <b>RootShield Home &amp; Garden</b> ( <i>Trichoderma harzianum</i> Rifai strain T-22), NC	Vegetables & Strawberries	<i>Cylindrocladium</i> spp., <i>Fusarium</i> spp., <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., <i>Thielaviopsis</i> spp.	0	0	Yes	See label for specific rates, application recommendations, and precautions regarding use and fungicide compatibility.
<b>RootShield Plus Granular</b> <b>RootShield Plus WP</b> ( <i>Trichoderma harzianum</i> Rifai strain T-22 and <i>Trichoderma virens</i> strain G-41), NC	Vegetables & Strawberries	<i>Cylindrocladium</i> spp., <i>Fusarium</i> spp., <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., <i>Thielaviopsis</i> spp.	0	4 hr	Yes	See label for specific rates, application recommendations, and precautions.
<b>RootShield WP</b> ( <i>Trichoderma harzianum</i> Rifai strain KRL-AG2), NC	Vegetables & Strawberries	<i>Cylindrocladium</i> spp., <i>Fusarium</i> spp., <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., <i>Thielaviopsis</i> spp.	0	0	Yes	See label for specific rates, application recommendations, and precautions regarding use and fungicide compatibility.
<b>Seican</b> (cinnamaldehyde)	Vegetables & Strawberries	Botrytis, Powdery mildew, Rusts	0	4 hr	No	See label for details, crop specific recommendations, and precautions.
<b>Serifel Biofungicide</b> <b>Serifel NG Biological Fungicide</b> <b>Subtlex NG</b> ( <i>Bacillus amyloliquetaciens</i> strain MBI 600), NC	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	Yes	See label for details, crop specific recommendations, and precautions.
<b>Sil-Matrix</b> (potassium silicate), NC	Vegetables & Strawberries	Broad-spectrum fungicide	0	4 hr	No	Must be used in a rotational program with other fungicides when conditions are conducive for disease development. See label for details.
<b>Soilgard</b> ( <i>Gliocladium virens</i> Gl-21), NC	Vegetables & Strawberries	Fusarium root and crown rot, <i>Phytophthora capsici</i> , <i>Pythium</i> spp., <i>Rhizoctonia</i> , <i>Sclerotinia</i> spp., <i>Sclerotium</i> spp.	0	0	Yes	For best results, apply to transplants or as a drench during transplanting. Subsequent applications can be made as drench, directed spray, or by chemigation. See label for precautions regarding use and fungicide compatibility.
<b>Sonata</b> ( <i>Bacillus pumilus</i> QST 2808), NC	Vegetables & Strawberries	Early blight, Downy mildew, Late blight, Powdery mildew, Rust	0	4 hr	Yes	Mix or alternate with other effective fungicides for improved disease control. See label for details.
<b>Sporatec</b> (oils of clove, rosemary and thyme), NC	Vegetables & Strawberries	Bacterial spot, Botrytis, Early blight, Gray mold, Late blight, Powdery mildew	0	0	Yes	Exercise care when applying. Begin applications once disease is observed. Use of a spreader and/or penetrant adjuvant recommended for improved performance. Do not apply when temps are above 90°F. See label for details. Ingredients are exempt from FIFRA.



Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
<b>Stargus</b> ( <i>Bacillus amyloliquefaciens</i> strain F727 cells and spent fermentation media)	Vegetables & Strawberries	Broad-spectrum fungicide and bactericide	0	4 hr	Yes	See label for specific rates and application recommendations.
<b>Tagro ECO</b> ( <i>Bacillus amyloliquefaciens</i> strain FZB24), NC	Vegetables	<b>Foliar diseases:</b> Downy mildew, Powdery mildew, <i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp. <b>Soilborne diseases:</b> <i>Fusarium</i> spp., <i>Phytophthora</i> spp., <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., <i>Sclerotinia</i> spp.	-	1	No	See label for specific instructions regarding soil injected, spray, or incorporated applications. Maximum of 12 applications per season. For best efficacy, product should be applied prior to disease or disease establishment. May be applied to greenhouse produced crops.
<b>Thymox</b> (thyme oil)	Vegetables & Strawberries	Botrytis and Powdery mildew	0	0	-	See label for specific rates and application recommendations. Ingredients are exempt from FIFRA.
<b>Timorex ACT</b> <b>Timorex Gold</b> (tea tree oil), 46	Vegetables & Strawberries	Broad-spectrum fungicide	2	1	No	See label for specific rates and application recommendations.
<b>Thyme Guard</b> (thyme oil extract), NC	Vegetables & Strawberries	Broad-spectrum bactericide and fungicide	0	0	-	See label for specific rates, application recommendations, and precautions regarding use with other pesticides. Ingredients are exempt from FIFRA.
<b>Trilogy</b> (clarified hydrophobic extract of neem oil), NC	Vegetables & Strawberries	<i>Alternaria</i> spp., Anthracnose, Botrytis, Early blight, Powdery mildew	0	4 hr	Yes	See label for specific rates, application recommendations, and precautions regarding use with other pesticides.
<b>Ultra-Pure Oil</b> (mineral oil), NC	Vegetables & Strawberries	Powdery mildew	0	4 hr	Yes	See label for specific rates, application recommendations, precautions, and fungicide compatibility.
<b>Vacciplant</b> (laminarin), P	Vegetables & Strawberries	Anthracnose, Bacterial speck, Bacterial spot, Early blight, Phytophthora blight, Powdery mildew	0	4 hr	No	Start applications preventatively, when weather conditions are favorable for disease development. Repeat applications until disease conditions end. Add a labeled copper product to VacciPlant if the disease symptoms appear.
<b>Veg'Lys, Brandt</b> (garlic oil), NC	Vegetables & Strawberries	Broad-spectrum bactericide and fungicide	0	0	Yes	See label for specific rates and application recommendations and precautions. Do not apply when temps are above 90°F. Ingredients are exempt from FIFRA.

Product (Active Ingredient), Fungicide Group <sup>1</sup>	Crops	Pertinent Diseases or Pathogens	Minimum Days to:		OMRI- Listed	Remarks <sup>2</sup>
			Harvest	Reentry		
Zayin (geraniol)	Vegetables & Strawberries	Powdery mildew	0	0		See label for specific rates and application recommendations.

<sup>1</sup>FRAC code (fungicide group): Number (33 and 44) and letters (NC and P) are used to distinguish the fungicide mode of action groups. All fungicides within the same group (with same number or letter) indicate the same active ingredient or similar mode of action. This information must be considered for the fungicide resistance management decisions. However, products with NC or P are considered low risk and don't require any rotation unless specifically directed on the label. NC = not classified, includes mineral oils, organic oils, potassium bicarbonate, and other materials of biological origin; P = host plant defense inducers. Source: FRAC Code List 2021 <http://www.frac.info/> (FRAC = Fungicide Resistance Action Committee).

<sup>2</sup>Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned.

Table 3. Biopesticides labeled for management of nematode pests of many crops.

Pest	MOA Code	Active Ingredient	Product(s)	Rate	Days to Harvest	REI (Hours)	Remarks
Nematodes	n/a	<i>Burkholderia rinojensis</i> heat-killed	Majestene	4–8 qt	0	4	OMRI-listed.
	n/a	<i>Purpureocillium lilicanus</i>	Melocon	2–4 lb	0	4	OMRI-listed.
	n/a	<i>Myrothecium verrucaria</i> fermentation product	DiTera	See label	0	4	OMRI-listed
	n/a	Mustard oil + capsaicin	Dazitol	6.25 gal	0	4	OMRI-listed. Apply 3–5 days preplant or 1/4 rate postplant.
	n/a	Azadirachtin	AzaGuard, Ecozine Plus, Molt-X	See label	0	4	OMRI-listed.
	n/a	Thyme oil	Nemakill, Promax, Thyme Guard	See label	0	4	OMRI-listed.