

Validation of an Area-Wide Extension Program to Estimate the Seasonal Abundance of Adult Citrus Root Weevils with Unbaited Pyramidal Traps¹

S. H. Futch, L. W. Duncan, and M. Zekri²

One hundred modified Tedders traps were monitored weekly for 2.5 years in each of seven citrus orchards located throughout Florida. Temporal patterns of the prevalence of adult *Diaprepes abbreviatus* at each site were compared to determine the feasibility of using insect abundance in sentinel orchards to schedule pest management activities within a region. Weevil abundance in three orchards on Florida's central ridge exhibited no consistent temporal patterns between years or among sites. In three central and eastern flatwoods sites, the abundance of weevils tended to be highest during the spring and early summer, but the variation in this pattern was high between sites and years. The mean abundance of weevils in the flatwoods sites was more than five fold greater ($P=0.02$) than that for sites on the central ridge. These data suggest the need for site-specific monitoring programs to aid in management of *D. abbreviatus*. Causes of regional variation in the abundance of *D. abbreviatus* merits additional research.

Diaprepes abbreviatus (Diaprepes root weevil) is a serious problem of widespread distribution within the Florida citrus industry. Methods to determine the local abundance of the insect are limited and expensive. Many growers do not know when weevils are present in their groves or

what the seasonal abundance is for their geographical areas. Without this basic information, control tactics may be improperly implemented. Unbaited pyramidal traps, commonly referred to as "Tedders traps," are sometimes employed to determine the seasonal abundance patterns of the weevils (Duncan et al. 2001). After emergence from the soil, the adults are attracted to any nonreflective silhouette (a citrus tree trunk or another structure placed in the grove) that simulates a tree trunk. Grower acceptance of the use of "Tedders traps" has been relatively poor due to the time and cost of regularly servicing and monitoring the traps. However, data from various studies have shown regional similarity in adult emergence and abundance on a seasonal basis. The objective of this study was to determine whether an area-wide scouting program using Tedders traps to estimate the emergence and seasonal abundance of adults can assist growers in instituting control methods for this weevil.

Materials and Methods

A two-and-one-half-year survey was implemented to estimate the seasonal abundance of adult *Diaprepes* root weevils using 100 unbaited pyramidal traps per location in six major regions in Florida. Three study sites were located

1. This document is HS994, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date February 2005. Revised February 2017 and May 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. S. H. Futch, Extension agent IV emeritus; L. W. Duncan, professor, Entomology and Nematology Department, UF/IFAS Citrus Research and Education Center; and M. Zekri, Extension agent IV, UF/IFAS Extension Hendry County; UF/IFAS Extension, Gainesville, FL 32611.

This research was supported by the Florida Agricultural Experiment Station, and approved for publication as Journal Series No. N-02573. This project is supported by the USDA Diaprepes Grant #727384112.

on the Florida central ridge in Lake and Polk Counties, and four sites were in flatwoods conditions in DeSoto, Hendry, Indian River and Dade Counties. Florida's central ridge is characterized by deep, well-drained sandy soils. Depth to groundwater is much shallower in the coastal and central "flatwoods" regions, where trees are generally grown on beds to increase the soil volume suitable for root growth. Soil texture and drainage characteristics vary greatly among flatwoods sites. Initially, all sites were commercially producing citrus groves known to be infested with *Diaprepes* for four to six years and were irrigated with low-volume microsprinkler irrigation systems. The Indian River and Dade County sites were changed to other locations after the first year of the study. The original Indian River site was replaced with another citrus grove that was under the same management whereas the Dade County site was relocated to an ornamental tree nursery. Site descriptions, sample areas and trap spacings are given in Table 1.

Data was collected by grower cooperators who reported the number of weevils captured in each trap during each previous seven-day period. Graphical data summaries were posted weekly and were freely available to growers, who were informed about the survey through Extension citrus newsletters, magazines, and meetings.

Soil samples from each site were processed to determine the particle size distribution. Oven-dried samples (100 g) were dispersed in a dilute alkaline solution of sodium metaphosphate on a mechanical shaker overnight and rinsed thoroughly on a 0.053 mm opening sieve to remove silt and clay particles. Soil remaining on the sieve was dried, and mechanically shaken on a nested series of sieves (0.053, 0.106, 0.25, 0.5, 1.0 mm; Rototap RX-29, W.S. Tyler, Mentor, OH), and fractions retained by each were weighed. Relationships between weevil abundance and soil particle size classes at each site were investigated by graphical examination and calculation of linear correlation coefficients.

Results and Discussion

Although locations were known to have been infested with *Diaprepes* for at least four to six years, the emergence patterns and average weevils collected per trap varied greatly between the study sites. Previous reports have indicated that although adult emergence and abundance can vary throughout the year, the primary emergence period is normally in April to mid-July and/or late-August to mid-October (McCoy et al. 2004). Data from this study agrees with year-round presence; however, the patterns often differed from discreet spring and fall peaks of abundance.

Following is a brief discussion of seasonal abundance patterns for each of the three well-drained sites in Lake and Polk locations (Figure 1).




Figure 1. Numbers of *Diaprepes abbreviatus* collected per trap each week during 31 months in three citrus orchards on the central ridge of Florida. Weekly data for each month were averaged to better illustrate seasonal patterns. Error bars are standard errors of the mean.

Lake County

Weevil abundance increased from early May 2001 until mid-July, when numbers peaked following a major rain event of greater than 6 inches in a one-week period. This spike was somewhat like an event at the DeSoto location below. Nigg et al. (2003) also reported that adult weevil emergence in a central Florida citrus orchard was initiated following the onset of the summer rains in July, following an unusually dry spring and early summer. After July, the weevil abundance remained relatively low (<0.1 weevils per trap per week) for the remainder of the survey year. Abundance during 2002 remained low the entire year at less than 0.1 weevils per trap per week until the fall, when weevil capture increased somewhat from November until February 2003. Thereafter, weevil abundance remained low until the survey terminated in October.

Polk County 1

This location had the lowest weevil population (<0.1 weevils per trap) of any of the six survey sites. Weevil populations declined from May 2001 until September when they increased slightly. Very few weevils were captured early

in 2002, until April. As in Lake County, weevil abundance increased in autumn of 2002, and highest numbers occurred in December. The number of weevils declined sharply in January 2003 and increased gradually thereafter. This grove was known to be infested with weevils for more than six years and is in a location where many other groves are infested with weevils.

Polk County 2

Trapping at the second Polk County site occurred from May 2002 until November 2003, and weevil abundance was low throughout the survey. Populations were slightly higher in May–June and December 2002 and may have increased briefly in September 2003.

The less well-drained sites are typical of the flatwoods and east coast areas include DeSoto, Hendry and Indian River (Figure 2).




Figure 2. Numbers of *Diaprepes abbreviatus* collected per trap each week during 31 months in three citrus orchards in flatwoods regions of Florida. Weekly data for each month were averaged to better illustrate seasonal patterns. Error bars are standard errors of the mean.

DeSoto County

The first major peak in abundance occurred from May through July 2001 followed by a steady decline in numbers for the remainder of the year. An interesting event occurred in early July 2001, when heavy rains triggered a major emergence totaling 2.14 weevils per trap for the weekly period (not shown). These heavy rains (12 inches in two

weeks) cause localized short-term flooding within the grove for several days. This event indicates that a large number of mature weevils can be in the soil and will emerge when forced or encouraged to emerge based upon environmental or other factors (Nigg et al. 2003). Weevil abundance remained relatively low (<0.10 weevils per trap) until May 2002. After July the numbers of weevils declined for the remainder of the year. Abundance again increased in March of 2003, with peak emergence in April, then declining somewhat into the summer.

Hendry County

This location had the highest average weevil levels among the seven sites. Weevil abundance was high in April 2001 at the beginning of the study (>0.5 weevils per trap) and decreased steadily for the remainder of 2001. In 2002, abundance began to increase in January, with peak recovery in March and April, and then declined for the remainder of the year. In 2003, weevil abundance followed a similar pattern, peaking in March and gradually declining thereafter.

Indian River County

Emergence at this location was extremely low (<0.04 weevils per trap) from the beginning of the survey in early April 2001 until December. This pattern did not follow the trends of the other flatwoods locations. Beginning in mid-December 2001, abundance began to increase until mid-April and then began to decline in late April 2002, when this grapefruit grove was removed due to small fruit size and low fruit prices. The grove was in good horticultural condition at the time of tree removal. The site was replaced with a grove in the same general area in October 2002. At the new site, abundance increased in December 2002, with peak emergence in January 2003, then declined until April, when populations levels increased somewhat until June.

Dade County

The abundance pattern varied from 0 to 0.15 weevils per trap per week but lacked any major or strong emergence patterns during 2001 (Figure 3). Weevils were trapped throughout the year at low levels. The location was receiving limited care (irrigation and nutritional inputs), which could have limited the flushing cycle of the trees, thereby reducing or limiting weevil numbers. Peak abundance occurred in June and July 2002. This site was replaced in September 2002 with an ornamental site that supported higher numbers of weevils. Emergence peaked at the ornamental site in January and again in August 2003.




Figure 3. Numbers of *Diaprepes abbreviatus* collected per trap each week during 31 months in a citrus orchard in Dade County, Florida. Weekly data for each month were averaged to better illustrate seasonal patterns. Error bars are standard errors of the mean.

Conclusions

The average weekly abundance of weevils was more than five fold greater ($P=0.02$) in the flatwoods sites (DeSoto, Hendry, Indian River Counties) than that in sites on the central ridge (Lake and Polk Counties) (Figure 4). Adult weevil numbers were highest at the DeSoto and Hendry locations, followed by Indian River. This regional trend for weevil population pressure is consistent with previous reports, although the causes of variation in weevil numbers between sites is unknown (Adair 1994; Stansly et al. 1997; Duncan et al. 2001). Duncan et al. (2003) reported greater natural control of *D. abbreviatus* by entomopathogenic nematodes in an orchard on coarse sandy soil on the ridge compared to one on sandy clay loam in the flatwoods. However, we detected no relationships when weevil abundance was plotted against the proportions of any of the sand particle sizes, and linear correlation coefficients between abundance and soil particle sizes were not significant ($P>0.10$).

When data were pooled within regions, weevil abundance was highest in the spring to mid-summer (March–July) of each year in the flatwoods, which is consistent with other reports (Adair 1994; Duncan et al. 2001). Seasonal abundance was not apparent for the pooled data from the ridge sites, although abundance has been reported to increase consistently in the spring and occasionally in the autumn from individual sites on the ridge (Duncan et al. 2001).




Figure 4. Average numbers of *Diaprepes abbreviatus* collected per trap each week during 31 months in three citrus orchards in flatwoods regions and three orchards on the central ridge of Florida. Weekly data for each month were averaged to better illustrate seasonal patterns. Inset shows the tree-year average for each grove used in the study. Error bars are standard errors of the mean.

While population trends within each geographical region should be somewhat similar, the temporal abundance of weevils varied significantly with location. Weevil pressure tended to be higher in flatwoods orchards compared to those on the central ridge.

Growers were provided an easily accessible source of timely information of weevil abundance trends in selected sites. Knowledge of these population patterns may help growers anticipate seasonal weevil population growth in order to time management strategies more effectively; however, management decisions should be based on supplemental observations regarding site-specific population levels.

References

- Adair, R. C. 1994. "A four-year field trial of entomopathogenic nematodes for control of *Diaprepes abbreviatus* in flatwoods citrus groves." *Proc. Florida State Hort. Soc.* 107: 63–68.
- Duncan, L. W., J. H. Graham, D. C. Dunn, J. Zellers, C. W. McCoy, and K. Nguyen. 2003. "Incidence of endemic entomopathogenic nematodes following application of *Steinernema riobrave* for control of *Diaprepes abbreviatus*." *J. Nematol.* 35:178.
- Duncan, L. W., C. W. McCoy, P. H. Stansly, J. H. Graham, and R. F. Mizell. 2001. "Estimating the relative abundance of citrus root weevils with modified Tedders traps." *Environ. Entomol.* 30:939–946.

McCoy, C. W., S. H. Futch, J. H. Graham, L. W. Duncan, and H. N. Nigg. 2004. *Citrus Root Weevils*. In *2004 Florida Citrus Pest Management Guide*, edited by L. W. Timmer. SP-43. Gainesville: University of Florida Institute of Food and Agricultural Sciences.

Nigg, H. N., S. E. Simpson, R. J. Stuart, L. W. Duncan, C. W. McCoy, and F. G. Gmitter Jr. 2003. "Abundance of *Diaprepes abbreviatus* (L.) (Coleoptera: Curculionidae) neonates falling to the soil under tree canopies in Florida citrus groves." *Horticultural Entomology* 96(3): 835–843.

Stansly, P. A., R. F. Mizell, and C. W. McCoy. 1997. "Monitoring *Diaprepes abbreviatus* with Tedder's traps in southwest Florida citrus." *Proc. Florida State Hort. Soc.* 110:22–26.

Table 1. Basic Site Information per Location for the Diaprepes Emergence Study.

Grove Factor	County Site Locations						
	Lake	Polk 1	Polk 2	Indian River	DeSoto	Hendry & Fallglo	Dade
Variety	Valencia	Hamlin & Pineapple	Mixed	Red grapefruit, switched to different location in Yr 2	Hamlin	Sunburst & Fallglo	Yr 1—Red grapefruit, Yr 2—ornamental nursery
Rootstock	Carrizo	Mixed, lemon, Swingle and Carrizo	Mixed	Yr 1, sour orange Yr 2, mixed rootstocks	Sour orange/bittersweet	Swingle	
Irrigation	Microsprinkler, 10 gph or less	Microsprinkler	Microsprinkler	Microsprinkler, 10 gph per tree	Microsprinkler, 23 gph per tree	Microsprinkler	Microsprinkler, but limited use
Year planted	1985	1950s with interplanted in the 70s & 80s	1960s	Yr 1—1979 Yr 2—mid 90	1990	1992	1990s
Year infested	1998	early 1990s	mid 1990s	1994 or 95	1997 or 98	1995 +/-	Unknown
Soil type	Astatula sand	Sparr sand & Millhopper fine sand		Winder fine sand	Ona and Smyrna	Boca and Chobee	Krome—mixture of marel & Rockdale
Spacing	15' x 25', no beds	15' x 30' with variation	12.5' x 25'	12' x 25', two row beds	10' x 25', four row beds	12.5' x 25', two row beds	12.5' x 25', flat planting with limited mounds for each tree
2001–2002 Yield estimate (boxes/acre)	400 boxes in better grove area with around 200 in poor areas	400 +/-		Est. 350, not harvested due to size and economic factors	675	319 Sunburst, 448 Fallglo	Not harvested due to canker rules and economics
Percent resets	90% mature trees	None within the last 3 years		1%–2%	Minimum 8%, some tree loss due to citrus tristeza virus	Few	Few
General grove conditions, comments	Test area varies in condition with well-drained soils in better condition. Part of the test area is near a water hole with poor drainage and is in poor condition.	The grove is in slightly-less-than average condition for the area.		Grove was in average condition, but removed due to low fruit prices in May of 2002 and scheduled to be replanted in June or July of 2002.	The grove is in average condition for the area with some foliage thinning.	Grove condition varies within block. Better-drained soil is in better condition than poorly drained soil.	Grove is in poor condition with little inputs for pest or nutritional care, little weed control.