

# Spanish Moth or Convict Caterpillar, *Xanthopastis timais* (Cramer) (Insecta: Lepidoptera: Noctuidae)<sup>1</sup>

John B. Heppner, Kathryn A. Barbara, and Eileen A. Buss<sup>2</sup>

## Introduction

The Spanish moth, *Xanthopastis timais* (Cramer), is unmistakable for any other moth in Florida. The larvae are likewise very colorful, and have been called convict caterpillars. The larvae are occasional pests of lilies, mainly in Amaryllidaceae. Spotted larval forms of Spanish moth appear similar to lily borer larvae, *Brithys crini* (Fabricius) (Godfrey 1972) of Europe and the Old World tropics, but only the banded larval form of the Spanish moth occurs in North America.



Figure 1. An adult Spanish moth, *Xanthopastis timais* (Cramer).  
Credits: John B. Heppner, DPI

## Synonymy

*Xanthopastis timais* (Cramer)

*Xanthopastis amaryllidis* Sepp

*Xanthopastis heterocampa* Guenée

*Xanthopastis regnatrix* Grote

*Xanthopastis antillium* Dyar

*Xanthopastis moctezuma* Dyar

*Xanthopastis molinoi* Dyar



Figure 2. An adult Spanish moth, *Xanthopastis timais* (Cramer).  
Credits: Jennifer L. Gillett-Kaufman, UF/IFAS Department of Entomology and Nematology

## Distribution

The Spanish moth, originally described from Surinam, is found throughout lowland areas of South and Central America, and in the Caribbean. The Spanish moth occurs throughout all lowland Neotropical regions of the

1. This document is EENY-271, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date June 2002. Revised June 2009 and February 2017. Reviewed June 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu>.
2. John B. Heppner, Florida Department of Agriculture, Division of Plant Industry; Kathryn A. Barbara; and Eileen A. Buss, associate professor, Department of Entomology and Nematology, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Caribbean, and as far south as northern Argentina. In North America, the species has a southeastern distribution, from the Carolinas to Texas, but strays northward along the Atlantic Coast as far as coastal New York, and inland as far north as Kentucky and Arkansas. It occurs in all of Florida ([Slosson 1894; see also Dyar [1901, 1902], Frost [1964], and Kimball [1965]).

## Description

Adult Spanish moths are relatively uniform in coloration throughout their range from North America to the Neotropics. Adults have rosy-pink and black forewings, spotted with orange along the wing veins, with a black body and gray hindwings. Some varieties have more white than pink on the forewings. Larvae are variable in Latin America, but in Florida are black with cream-white or yellow-white bands and orange head, prolegs, and posterior end. In addition to the whitish band, each body segment also has a dorsal and a lateral patch of cream-white near each band. The head and posterior end each have two eye-like black spots, making the posterior end appear much like the head. Variation in the Neotropics is mostly in the amount of orange on the head and posterior, and with the yellow-white body banding more as spots than bands. Larvae are up to 5 cm long and feed gregariously on leaves, bulbs, and rhizomes of the host plants. Numbers of larvae can damage lily leaves in a short time. The pupa of the Spanish moth is typical for noctuid moths and almost black in color. Eggs are rounded (somewhat flattened) and yellowish, otherwise typical for noctuid moths (Dyar 1901).



Figure 3. Spanish moth larva (convict caterpillar), *Xanthopastis timais* (Cramer), feeding on amaryllis.

Credits: Terry DelValle, UF/IFAS

## Life Cycle

In Florida, Spanish moth adults are active from January to early June, and September to December, but may have nearly continuous generations in the southernmost areas of the state and possibly also in greenhouse production. Females lay several hundred eggs grouped in clusters

(Bourquin 1935), usually on the lower leaf surfaces. Larvae pupate in loose soil. Larvae have six instars (Bourquin 1935). Larval eclosion takes about eight days under optimal conditions, followed by 17 days of larval feeding. The pupal stage lasts about 19 days. Adults usually live about eight to 10 days, including two to three days for adult flight and mating. A total generation time of seven to eight weeks allows up to six generations per year.

## Host Plants

Spanish moth larvae mainly feed on spider lilies and other Amaryllidaceae, plus Iridaceae and Liliaceae (Tietz 1972). Reports of *Ficus* (Moraceae) and *Hibiscus* (Malvaceae) as hosts are probably erroneous, but *Xanthosoma* and *Zantedeschia* (both Araceae) are rare alternate hosts. Incidental records include *Cocoloba uvifera* (Polygonaceae) (DPI record) and *Polianthes tuberosa* (Agavaceae) (Pirone 1970); also *Lactuca* sp. (Compositae) (Covell 1984) in lab rearings.

Host plant records in Amaryllidaceae include *amaryllis*, *Clivia*, *Cooperia*, *Eucharis*, *Haemanthus*, *Hippeastrum*, *Hymenocallis*, *Narcissus*, *Pancratium*, *Polianthes*, and *Zephyranthes*; in Iridaceae, *Iris*; and in Liliaceae, *Crinum*, *Leucojum* and *Lilium*.

## Damage

Spanish moth larvae cause damage by chewing gregariously on leaves, bulbs, and rhizomes of the host plants. Economic damage to lilies by Spanish moth has been noted by Biezanko and Guerra (1975), Bourquin (1935), Bruner et al. (1975), D'Angelo (1941), Figueiredo and Pereira (1944), Gundlach (1881), Martorell (1976), Monte (1932, 1934), Pirone (1970), and Wolcott (1936, 1951).

## Management

### Sampling

Periodically examine leaves for Spanish moth caterpillars. Direct observation of larvae on the plant during the early stages of growth in the spring is the best sampling option due to the plant's small size. Adult moth populations can be sampled using a blacklight. Once moths are detected, searching for eggs and larvae is the next step.

### Insecticides

Larvae can be treated with a bacterial spray (*Bt*), or more immediate results can be obtained from the application of various pesticides. In French Guiana, an ectoparasitic nematode has been reported on larvae of Spanish moth and

other noctuid moths (Rogers et al. 1990). Brunner et al. (1975) note a Tachinid fly parasitoid.

## Florida Insect Management Guide for Landscape Plants

### Selected References

Biezanko CM, Guerra M de S. 1975. Contribuição ao estudo de *Xanthopastis timais* Stoll, 1782 importante praga das amarilidáceas (Lepidoptera, Heterocera, Noctuidae). Arquivos do Museu Nacional 54: 267–272.

Bourquin F. 1935. Metamorfosis de *Xanthopastis timais* Cr. (Lep. Noct.). Revista de la Sociedad Entomologica de Argentina 7: 195–201.

Bruner SC, Scaramuzza LC, Otero AR. 1975. Catalogo de los Insectos que Atacan a las Plantas Economicas de Cuba. 2<sup>nd</sup> ed. Academia de Ciencias de Cuba, Inst. de Zoología, Havana. 399 p.

Covell Jr CV. 1984. Field Guide to the Moths of Eastern North America. Houghton-Mifflin, Boston. 496 p.

D'Angelo WA. 1941. Mariposa dañina para ciertas plantas de jardín *Xanthopastis timais*, Cram. Almanaque de Ministerio de Agricultura Argentinas 16: 169–172.

Dyar HG. 1901. Notes on the winter Lepidoptera of Lake Worth, Florida. Proceedings of the Entomological Society of Washington 4: 446–485

Dyar HG. 1902. The larva of *Euthanisotia timais* Cram. Journal of the New York Entomological Society 10: 125–126.

Figueiredo Jr ER de, Pereira HF. 1944. Notas sobre *Xanthopastis timais* (Cram.) (Lep. Noct.), praga das amarilidáceas. Arquivos do Instituto de Biologica (São Paulo) 15: 289–298.

Frost SW. 1964. Insects taken in light traps at the Archbold Biological station, Highlands County, Florida. [Florida Entomologist](#) 47: 127–161.

Godfrey GL. 1972. A review and reclassification of larvae of the subfamily Hadeninae (Lepidoptera, Noctuidae) of America north of Mexico. U.S. Department of Agriculture Technical Bulletin 1450. 265 p.

Gundlach J. 1881. Contribución á la Entomología Cubana. Havana. 445 p.

Kimball CP. 1965. The Lepidoptera of Florida: an annotated checklist. In Arthropods of Florida and Neighboring Land Areas. Vol. 1. Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville. 363 p., 26pl.

Martorell LF. 1976. Annotated Food Plant Catalog of the Insects of Puerto Rico. Agricultural Experiment Station, University of Puerto Rico, Rio Piedras. 303 p.

Monte O. 1932. Notas biológicas sobre o lepidoptero noctuido *Xanthopastis timais* Cram. O Campo (Rio de Janeiro) 3: 41–42.

Monte O. 1934. Borboletas que vivem em plantas cultivadas. Boletim de Agric. Zootechnia e Veterinária (Ser. Agricola) 7: 111–112.

Pirone PP. 1970. Diseases and Pests of Ornamental Plants. 4<sup>th</sup> ed. Ronald Press, New York, 546 p.

Rogers CE, Marti OG, Simmons AM, Silvain JF. 1990. Host range of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae): an ectoparasite of moths in French Guiana. Environmental Entomology 19: 795–798.

Slosson AT. 1894. Florida field notes. Journal of the New York Entomological Society 2: 106–107.

Tietz HM. 1972. An Index to the Described Life Histories, Early Stages and Hosts of the Macrolepidoptera of the Continental United States and Canada. Allyn Museum of Entomology, Sarasota. 2 v.

Wolcott GN. 1936. Insectae Borinquenses- a revised annotated checklist of the insects of Puerto Rico. Journal of Agriculture, University of Puerto Rico 20: 1–600.

Wolcott GN. 1951. The insects of Puerto Rico. Journal of Agriculture, University of Puerto Rico 32: 1–975.