

# Chapter 7. Cucurbit Production<sup>1</sup>

Josh H. Freeman, Eugene J. McAvoy, Nathan S. Boyd, Mathews Paret, Qingren Wang, Christian F. Miller, Johan Desaeger, Jawwad Qureshi, Xavier Martini, and Phillip B. Williams<sup>2</sup>

## Botany and Planting

**Angled luffa** (silk squash)—*Luffa acutangula*

**Bittermelon**—(Chinese and Indian types)—*Momordica charantia*

**Butternut squash**—*Cucurbita moschata*

**Cantaloupe**—*Cucumis melo*

**Chayote**—*Sechium edule*

**Chinese cucumber**—*Trichosanthes kirilowii*

**Cucumber**—*Cucumis sativus*, Cucurbitaceae

**Fuzzy melon** (immature fruit) and **Winter melon**—*Benincasa hispida*

**Long gourd** (oopoh)—*Lagenaria siceraria*

**Pumpkin** (jack-o-lantern is *C. pepo*; some processing pumpkins are *C. maxima* and *C. moschata*)

**Smooth luffa**—*Luffa aegyptica* (cylindrical)

**Snake gourd**—*Trichosanthes cucumerina*

**Summer squash**—*Cucurbita pepo*

**Tropical pumpkin (calabaza)**—*Cucurbita moschata*

**Watermelon**—*Citrullus lanatus*

**Winter squash**—*Cucurbita maxima*

1. This document is HS725, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date June 1995. Revised April 2020. This is Chapter 7 of the *Vegetable Production Handbook of Florida, 2020–2021 edition*. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Joshua H. Freeman, associate professor, UF/IFAS North Florida Research and Education Center; Eugene McAvoy, Extension agent IV, UF/IFAS Extension Hendry County; Nathan S. Boyd, associate professor, UF/IFAS Gulf Coast REC; Mathews Paret, assistant professor, UF/IFAS North Florida REC; Qingren Wang, Extension agent I, UF/IFAS Extension Miami-Dade County; Christian F. Miller, Extension agent I, UF/IFAS Extension Palm Beach County, West Palm Beach; Johan Desaeger, assistant professor, UF/IFAS Gulf Coast REC; Jawwad Qureshi, assistant professor, Entomology and Nematology Department, UF/IFAS Southwest Florida REC; Xavier Martini, assistant professor, UF/IFAS North Florida REC; and Phillip B. Williams, assistant professor, Horticultural Sciences Department, UF/IFAS Southwest Florida REC; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Table 7.1. Planting information for cucurbits.

Planting Dates	Cucumber	Cantaloupe	Pumpkin	Squash		Watermelon
North Florida	Feb–Apr; July–Aug	Feb–Apr	Early July	Feb–Apr; Aug–Sept		Feb–Apr
Central Florida	Jan–Mar; Sept	Jan–Mar	Mid-July	Jan–Apr; Aug–Sept		Jan–Mar
South Florida	Sept–Feb	Dec–Mar	Early August	Aug–Mar		Dec–Mar
Planting Information				Bush	Vining	
Distance between rows (in)	48–60	60–72	60–108	36–48	60–108	60–108
Distance between plants (in)	6–12	20–36	36–60	12–24	36–60	24–48
Seeding depth (in)	0.5–0.75	0.5–1.0	1.5–2.0	1.0–1.5	1.5–2.0	1.5–2.0
Seed per acre (lb)	2–4	1–2	4–5	2–3	1–1.5	1–3
Days to maturity from seed	40–65	85–110	80–100	40–50	85–120	80–100
Days to maturity from transplant	Not recommended		70–90	Not recommended		60–90
Plant populations (acre)	21,780	4,356	2,904	7–14,520	2,904	1,815

## Cultivars

Table 7.2. Cultivars for cantaloupe.

Eastern Shipper			Harper Type		Calabaza (Tropical Pumpkin)
Aphrodite (H)	Athena (H)	Avatar (H)	Caribbean Gold	Timeless Gold	Agriset 9001 (vining type)
Ariel (H)	Atlantis (H)		Infinite Gold		La Estrella (compact plant)
H=hybrid					

Table 7.3. Cultivars for cucumber.

Pickling				Slicing		
Calypso (H) (GY)	Excursion	Jackson Supreme	Supremo (H)	Bristol	Mongoose (H) (GY)	SV4719CS* (H) (GY)
Deli King	Expedition (H)	Maxi Pack (H)	Transamerica (H)	Cobra (H) (GY)	Python (H)	Talladega (H) (GY)
Deli Star	Fancipak (H) (GY)	Maxi Pick (H)	Vlaspik (H)	Dominator (H) (GY)	Speedway (H) (GY)	Thunder (H) (GY)
Eureka (H) (MO)	FMX 5020 (H)	Powerpak (H) (GY)		Impact (H) (GY)	SV3462CS* (H) (GY)	
Excel (H) (GY)	Jackson Classic (H) (GY)	Royal (H) (GY)				
H=hybrid; flower habit: GY=gynoecious, MO=monoecious. * Improved downy mildew resistance.						

Table 7.4. Cultivars for Halloween pumpkin (north Florida only).

Miniature: < 1 lb	Small: 1–5 lb	Medium: 5–10 lb	Large: 10–20 lb		Giant: 25–80 lb
Bumpkin (H)	Hybrid Pam (H)	Autumn Gold (H)	Merlin (H)	Apogee (H)	Magician (H)
Gold Dust (H)	Pick-A-Pie (H)	Goose Bumps II (H)	October (H)	Connecticut Field	Magic Wand (H)
Jill-Be-Little	Small Sugar	Jack of All Trades (H)		Early Giant (H)	Prizewinner (H)
Wee-Be-Little (PVP)		Knucklehead (H)		Early King (H)	
PVP=Plant Variety Protection; H=hybrid					

Table 7.5. Cultivars for squash.

Summer (yellow)			Acorn	Butternut	
Conqueror III (H) (SN)*	Goldprize (H) (SN)	Prelude II (H) (CN) *	Mesa Queen (H)	Atlas (H)	Quantam (H)
Destiny III (H) (CN)*	Gold Star (H) (CN)	Sunglo (H) (CN)	Royal Ace (H)	Avalon (H)	Ultra (H)
Enterprise (H) (SN)	Grand Prize (H) (SN)	XPT 1832 III (H) (SN)*	Table Ace (H)	Canesi (H)	Waltham
Fortune (H) (SN)	Lazor (H) (SN)		TayBelle PM (H)	Early Butternut (H)	Zenith (H)
Gentry (H) (CN)	Lioness (H) (SN)			Polaris (H)	
Summer (zucchini)				Summer (yellow zucchini)	Summer (gray zucchini)
Everglades (H)	Justice III (H)*	Payroll (H)	Spineless King (H)	Golden Delight (H) (SN)	Hurakan (H)
Green Eclipse (H)	Pascola (H)	Respect (H)	Spineless Perfection (H)	Golden Glory (H) (SN)	Lolita (H)
Green Machine (H)	Paycheck (H)	Senator (H)	SV6009YG (H)*	Goldrush (H) (SN)	Luciana
Judgement III (H)*	Payload (H)	Spineless Beauty (H)			
H=hybrid; type: CN=crookneck, SN=straightneck. *Biotech: Specific requirements and restrictions apply to these products.					

Table 7.6. Cultivars for watermelon.

Diploid			Triploid (seedless, large)			
Estrella (H)	Stargazer (H)		ACX 6177 (H)	Crunchy Red (H)	Secretariat(H)	Unbridled (H)
Gold Strike (H) (orange flesh)	Summer Flavor 790 (H)		Admiration (H)	Exclamation (H)	Sugar Fresh (H)	Warrior (nun 1009) (H)
Jamboree (H)	Summer Flavor 800 (H)		Affirmed (H)	Fascination (H)	Sugar Heart (H)	SS 7167 (H)
Mardi Gras (H)	Summer Flavor 8490 (H)		Bold Ruler (H)	Joy Ride (SV 8298) (H)	Summer Breeze (H)	SS 7177 (H)
Sangria (H)	Summer Velvet 2800 (H)		Bottle Rocket (H)	Kingman (H)	Sweet Gem (H)	SS 7197 (H)
Sentinel (H)	Top Gun (H)		Captivation (H)	Maxima (H)	Sweet Polly (H)	SV 0241 (H)
SSX 8585 (H)	Vista (H)		Citation (H)	Melody (H)	Talca (H)	
Starbrite (H)			Crisp N' Sweet (H)	Razorback (H)	Traveler (H)	
			Cronos (H)	Road Trip (H)	Troubadour (H)	
Nonharvestable Pollenizer			Triploid (seedless, mini)			
Accomplice	Polimax	Wildcard Plus	Ana (H)	Leopard (H)	Promesa (H)	Sky View (H)
Ace Plus	SP-6	Wingman	Belmont (H)	Ocelot (H)	Sky Line (H)	Sugar Bite (H)
Patron	SP-7		Extazy (H)	Pixie (H)	Sky Star (H)	

H=hybrid

## Asian Cucurbits

This group includes cucurbit fruits that can be eaten either immature or mature as well as several vegetables with edible tender stems and leaves. All can be grown on raised beds, with or without plastic mulch, and with drip, overhead, or subsurface irrigation. Most of the crops are trellised, primarily to maximize space and sufficient sunlight, to minimize bud drop and fruit rot caused by overshading and exposure to soil moisture and pathogens, and to promote straight fruit. Winter melon is the exception because it is generally too heavy to trellis. Fertilizer recommendations for cucumbers are applicable for fuzzy melon, long gourd, both luffas, Chinese cucumber, bittermelon, and snake gourd. There are two types of bittermelon: Indian and Chinese. The Indian type has smaller fruit with dark green color and rough skin, and the Chinese type has larger fruit with light green color and smooth skin. Chinese cucumber,

though rarely commercially available, has fruit more than 1 foot long with dark green color, very thin and rough skin, and crisp texture, similar to European cucumber. Recommendations for watermelon should be followed for winter melon and chayote. With the exception of chayote, where the entire fruit is planted, these crops are started from seed and grown as transplants prior to being set in the field.

Table 7.7. Planting information for Asian cucurbits.

Planting Dates	Bittermelon	Long Gourd	Angled Luffa	Smooth Luffa
North Florida	Feb–Apr; July–Aug	Feb–Apr; July–Aug	Feb–Apr; July–Aug	Feb–Apr; July–Aug
Central Florida	Jan–Mar; Sept	Jan–Mar; Sept	Jan–Mar; Sept	Jan–Mar; Sept
South Florida	Sept–Feb	Sept–Feb	Sept–Feb	Sept–Feb
Planting Information				
Distance between rows (in)	60–72	60–72	60–72	60–72
Distance between plants (in)	36–60	36–60	36–60	36–60
Days to maturity from seed	80–100			
Plant populations (acre)	2904	2904	2904	2904
Planting Dates	Fuzzy Melon	Snake Gourd	Chayote	Winter Melon
North Florida	Feb–Apr; July–Aug	Feb–Apr; July–Aug	Not recommended	Feb–Apr
Central Florida	Jan–Mar; Sept	Jan–Mar; Sept	Not recommended	Jan–Mar
South Florida	Sept–Feb	Sept–Feb	Sept–Feb	Dec 15–Mar 1
Planting Information				
Distance between rows (in)	60–72	60–72	60–72	72–108
Distance between plants (in)	36–60	36–60	36–60	36–72
Seeding depth (in)	1.5–2.0	1.5–2.0	Whole fruit is used: cover halfway	1.5–2.0
Seed per acre			2904—whole fruit	
Plant populations (acre)	2904	2904	2904	1452–2420

Table 7.8. Cultivars for Asian cucurbits.

Bittermelon (Chinese)*	Bittermelon (Indian)*	Winter Melon	Snake Gourd*	Angled Luffa*
Hong Kong Green	Hybrid India Star	Giant Wax	Buag Ngu	Bonanza
Hybrid Bangkok Large	India Green Long	Hybrid Red Doll	Extra Long Dancer	Buam Liam Hybrid
Japan Green Sprindle	India Green Queen	Hybrid Small Round	Hybrid Snaky	Hybrid Green Glory
Taiwan Large	Palee	Hybrid Wonder Wax	Long EX	Lucky Boy
		Round Tong Gwa	Polo	Summer Long
Smooth Luffa*	Long Gourd*	Fuzzy Melon*	Chayote*	
Dok Hybrid	Bitter Gourd Long	Chiang Shin Joker	The seed is the viviparous fruit itself. There is some debate as to whether varieties remain true. Fruit purchased from grocery stores may not sprout.	
Hybrid Smooth Beauty	Hybrid Asia Short	Mokwa		
Smooth Boy	Hybrid India Long	Mokwa Long		
		Seven Star Long		

\* Should be trellised

Table 7.9. Herbicides approved for managing weeds in cucurbit crops.

Active Ingredient lb a.i./Acre	Trade Name Product/Acre	MOA Code	Crops	Weeds Controlled/Remarks
<b>Labels change frequently. Be sure to read a current product label before applying any chemical.</b>				
<b>***PREEMERGENCE***</b>				
<b>Bensulide</b> 5.0–6.0	(Prefar) 4 E 5–6 qt	8	All cucurbits	Annual broadleaf and grass control. Incorporate or irrigate 1 to 2 in. within 36 hr of application. Nonlabeled crops should not be planted within 120 days of application.
<b>Carfentrazone</b> up to 0.031	(Aim) 1.9 E up to 2 fl oz (Aim) 2.0 EC up to 2 fl oz	14	All cucurbits	Apply as a preplant burndown for emerged broadleaf weeds. Use crop oil concentrate, methylated seed oil, or nonionic surfactant at recommended rates. Maximum rate of 0.096 lb a.i./A per season. No pretransplant interval.
<b>Clomazone</b> 0.15–0.38	(Command) 3 ME 0.4–1 pt	13	Cucumber	Annual broadleaf and grass weeds. Use lower rates on coarse soils. Read label for additional restrictions.
<b>Clomazone</b> 0.15–0.25	(Command) 3 ME 0.4–0.67 pt	13	Muskmelon, watermelon	Annual broadleaf and grass weeds. Use lower rates on coarse soils. Read label for additional restrictions.
<b>Clomazone</b> 0.25–0.5	(Command) 3 ME 0.67–1.33 pt	13	Summer squash	Annual broadleaf and grass weeds. Broadcast for bare-ground production; row middles only if growing on plastic.
<b>Clomazone</b> 0.25–0.75	(Command) 3 ME 0.67–2 pt	13	Winter squash	Annual broadleaf and grass weeds. Broadcast for bare-ground production; row middles only if growing on plastic. Read labels for cultivar restrictions.
<b>Ethalfuralin + Clomazone</b> 0.4–0.6 + 0.13–0.19	(Strategy) 2–3 pt	3 + 13	Cucumbers, melons, summer & winter squash, pumpkin, watermelon	Annual broadleaf and grass control. Must be applied no later than 2 days after seeding. Overhead irrigation or rainfall of 0.5 in. within 5 days. Do not apply under row mulch or over top of plants.
<b>Flumioxazin</b> up to 0.125	(Chateau) 51 WDG up to 4 oz	14	Cucumber, muskmelon, watermelon, pumpkins, summer & winter squash	Broadleaf control. Row middles only. Do not apply after crops are transplanted/seeded. Label is a Third-Party Registration (TPR, Inc.). Use without a signed authorization and waiver of liability is a misuse of the product.
<b>Glyphosate</b>	(various formulations) consult labels	9	All cucurbits	Controls emerged broadleaf and grass weeds. Consult individual labels for restrictions.
<b>Halosulfuron</b> 0.024	(Sandea) 75 DG 0.5 oz	2	Cantaloupe, cucumber, crenshaw, honeydew	Yellow and purple nutsedge and broadleaf control. Apply uniformly with ground equipment in a minimum of 15 gallons of water/A.
<b>Halosulfuron</b> 0.024–0.036	(Sandea) 75 DG 0.5–0.75 oz	2	Watermelon	Yellow and purple nutsedge and broadleaf control. May be applied preemergence to seeded watermelon on bare-ground or preseeding to mulch-cultured watermelon. Transplanting should be no sooner than 7 days after application. Use lighter rates on sandy soils with low organic matter.
<b>Halosulfuron</b> 0.024–0.036	(Sandea) 75 DG 0.5–0.75 oz	2	Pumpkin, winter squash	Yellow and purple nutsedge and broadleaf control. Apply before soil cracking or pretransplant. Transplanting should not be made sooner than 7 days after application. May be applied POST over the top when plants reach the 4–5 true-leaf stage, but before first female flowers appear.
<b>Paraquat</b> 0.63–0.94	(Gramoxone) 2 SL 2–4 pt (Firestorm) 3 SL 1.3–2.7 pt	22	Cucumber, muskmelon, cantaloupe, pumpkin, squash, watermelon	Controls emerged weeds. Apply prior, during, or after planting, but before crop emergence. Use a nonionic surfactant. <b>Consult label for new restrictions.</b>
<b>Pelargonic acid</b>	(Scythe) 4.2 EC 3%–10% v/v	27	All cucurbits	Controls emerged weeds. Apply before emergence of crop. Product is a contact, nonselective, foliar-applied herbicide. There is no residual activity. May be tank-mixed with soil residual herbicides.
<b>Pendimethalin</b> 1	(Prowl H20) 2.1 pt	3	Cantaloupe, citron melon, muskmelon, watermelon	Annual broadleaf and grass weeds. Apply to the row middles only. May be applied as a sequential application with 2.1 pt/A before and 2.1 pt/A after transplanting or seeding with 21 days between applications. Do not exceed 4.2 pt/A per season.

Active Ingredient lb a.i./Acre	Trade Name Product/Acre	MOA Code	Crops	Weeds Controlled/Remarks
<b>S-metolachlor</b> 0.95–1.26	(Brawl, Dual Magnum) 1.0–1.33 pt	15	Pumpkin	Annual broadleaf and grass weeds and nutsedge control. Apply as inter-row or inter-hill application. Leave a 1-foot untreated area over the seeded row (6 in. on either side of the row). Use lower rates on lighter soils. Apply before weeds emerge.
<b>*** POSTEMERGENCE ***</b>				
<b>Carfentrazone</b> up to 0.031	(Aim) 1.9 E up to 2 fl oz (Aim) 2.0 EC up to 2 fl oz	14	All cucurbits	Emerged broadleaf control. Post-direct hooded application to row middles for burndown of emerged broadleaf weeds. Use crop oil concentrate or nonionic surfactant at recommended rates. PHI 0 days.
<b>Clethodim</b> 0.94–0.13 0.07–0.13	(Arrow, Intensity, Select) 2 EC 6–8 oz (Select Max) 1 EC 9–16 fl oz	1	Cucumber, squash, melons, and all commodities in crop group	Annual and perennial grass control. Use a crop oil concentrate at 1% v/v spray volume. Use nonionic surfactant in Select Max. PHI 14 days.
<b>DCPA</b> 4.5–10.5	(Dacthal) 6 F 6–14 pt	3	Muskmelon, cantaloupe, honeydew, watermelon	Annual grasses and certain broadleaf control. Apply only when plants have 4–5 true leaves, are well-established, and growing conditions are favorable for good plant growth. Cultivate prior to application to control emerged weeds.
<b>Ethalfuralin + Clomazone</b>	(Strategy) 2–3 pt	3 + 13	Cucumber, melon, summer & winter squash, pumpkin, watermelon	Annual broadleaf and grass control. After transplanting apply to row middles only. Does not control emerged weeds. PHI 45 days.
<b>Glyphosate</b>	(various formulations) consult labels	9	All cucurbits	Control emerged broadleaf and grass weeds. Apply to row middles only. Use a hooded sprayer to reduce drift. Consult individual labels for restrictions.
<b>Halosulfuron</b> 0.04–0.05	(Sanda) 75 DF 0.5–1 oz	2	Cucumber, cantaloupe, honeydew, crenshaw melon	Apply after the 3–5 true leaf stage (no sooner than 14 days after transplanting) but before the first female flower. Spot spray a POST application if a PRE application was made. Cucumber PHI 30 days. Cantaloupe, honeydew, crenshaw 57 days.
<b>Halosulfuron</b> 0.02–0.05 0.02–0.04	(Sanda) 75 DF 0.5–1.0 oz 0.5–0.75 oz	2	Watermelon, summer squash; Pumpkin, winter squash	Nutsedge and broadleaf weeds. Apply to the row middles only. Include a NIS in the spray solution. PHI 57 days in watermelon. PHI 30 days for summer/winter squash and pumpkin.
<b>Imazosufuron</b> 0.19–0.3	(League) 75 WG 4–6.4 oz	2	Muskmelon, watermelon	Nutsedge and broadleaf weeds. No more than 6.4 oz/A/yr. One application per years. Surfactant is needed if weeds are emerged. Row middles only. PHI 45 days.
<b>Paraquat</b> 0.47–0.93	(Gramoxone) 2 SL 1.88–3.72	22	Cucumber, muskmelon, cantaloupe, pumpkin, squash, watermelon	Controls emerged weeds. Row middles only. Limit of 3 applications per year. Gramoxone SL is a supplemental label consult labels for other paraquat formulations. <b>Consult label for new restrictions.</b>
<b>Pelargonic acid</b>	(Scythe) 4.2 EC 3%–10% v/v	27	Cucumber, gourd, muskmelon, cantaloupe, pumpkin, squash, watermelon	Controls emerged weeds. Row middles only. Use a shielded sprayer directed to the row middles to reduce drift to the crop.
<b>Pendimethalin</b> 1	(Prowl H20) 2.1 pt	3	Cantaloupe, citron melon, muskmelon, watermelon	Annual broadleaf and grass weeds. Apply to the row middles only. Apply 2.1 pt/A after transplanting or seeding and before vines are off the bed and in the spray area. If a preplant application was applied, do not exceed 4.2 pt/A per season. PHI 35 days.
<b>S-metolachlor</b> 0.95–1.26	(Brawl, Dual Magnum) 7.62 EC 1.0–1.33 pt	15	Pumpkin	Annual broadleaf and grass weeds and nutsedge. Apply as inter-row or inter-hill application. Leave a 1-foot untreated area over the plant (6 in. on either side of the row). Use lower rates on lighter soils. Apply before weeds emerge. PHI 30 days.
<b>Sethoxydim</b> 0.19–0.28	(Poast) 1.5 EC 1.0–1.5 pt	1	All cucurbits	Growing grass weeds. Include a crop oil concentrate. Efficacy is decreased if weeds are under stress. Use 1 pt for seedling grasses and 1.5 pt on perennial grasses. PHI 14 days.
<b>Terbacil</b> 0.1–0.2	(Sinbar) 80 WP 2–4 oz	5	Watermelon	Annual broadleaf weeds. Apply to row middles only. Do not allow contact with plant foliage. Do not exceed 4 oz per year. PHI 70 days.
<b>Trifluralin</b>	(Treflan) 4 EC 1.0 pt (Treflan) TR 10 5 lb	3	All cucurbits	Annual broadleaf and grass control. Apply after crop is 3–4 true leaf stage. Row middles only. PHI 30 days except for watermelon 60 days.
<b>*** POSTHARVEST ***</b>				
<b>Diquat</b> 0.25	(Reglone Dessicant) 1.0 pt	22	Cantaloupe	Minimum of 35 gal/A. Include an NIS. Thorough coverage is required.
<b>Diquat</b> 0.38	(Reglone Dessicant) 1.5 pt	22	Cucumber, watermelon, squash (except acorn)	Minimum of 35 gal/A. Include an NIS. Thorough coverage is required.
<b>Diquat</b> 0.5	(Reglone Dessicant) 2.0 pt	22	Squash (acorn)	Minimum of 35 gal/A. Include an NIS. Thorough coverage is required.

Table 7.10. Insecticides labeled for management of arthropod pests of cucurbit crops.

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
Aphids	1B	<b>Dimethoate 4 EC</b> (dimethoate)	<b>Melons:</b> 1 pt <b>Watermelons:</b> 0.5–1 pt	2 pt per year	48	3	Highly toxic to bees. Not for squash or cucumber.
	1B	<b>Malathion 5EC</b> <b>Malathion 8</b> (malathion)	<b>5EC:</b> cucumber: 1.5–2.8 pt summer squash: 2.0–2.8 pt winter squash: 1.6 pt watermelon: 1.5–2.5 pt melon: 1.6 pt <b>8:</b> summer squash: 1.75 pt summer squash: 1.75 pt winter squash: 1.75 pt cucumber: 1.75 pt	<b>5EC:</b> cucumber: 5.6 pt summer squash: 8.4 pt winter squash: 4.8 pt watermelon: 5 pt <b>8:</b> cucumber: 3.5 pt summer squash: 5.25 pt winter squash: 5.25 pt	24	1	<b>5EC:</b> Maximum number of applications per year is 2. <b>8:</b> Squash and cucumbers only.
	3A	<b>*Brigade 2 EC</b> (bifenthrin)	2.6–6.4 fl oz	19.2 fl oz	12	3	Do not make more than 2 applications after bloom. Highly toxic to foraging bees.
	3A	<b>*Athena</b> (bifenthrin)	7–17 fl oz	51 fl oz	12	7	Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
	3A	<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees. Can be a tank mix with 3–4 oz of Belay insecticide.
	4A	<b>Actara</b> (thiamethoxam)	1.5–3.0 oz	11.0 oz	12	0	Apply before pests reach damaging levels. Highly toxic to foraging bees.
	4A	<b>Admire Pro</b> (imidacloprid) (see appropriate labels for other brands)	7–10.5 oz <b>Planthouse:</b> 0.44 fl oz/10,000 plants	10.5 fl oz	12	<b>Soil:</b> 21	<b>Planthouse:</b> One application to transplants. See label for use on mature greenhouse cucumbers.
	4A	<b>Assail 305G</b> (acetamiprid)	2.5–4 oz	20.0 oz	12	0	No more than 5 applications per season. Do not use with other group 4A insecticides. Toxic to foraging bees.
	4A	<b>Belay Insecticide</b> (clothianidin)	<b>Foliar:</b> 3.0–4.0 fl oz <b>Soil:</b> 9.0–12.0 fl oz	12 fl oz	12	<b>Foliar:</b> Do not make application after 4th true leaf on main stem is unfolded. <b>Soil:</b> 21	Do not apply when plants are blooming.
	4A	<b>Platinum 755G</b> (thiamethoxam)	5–11 fl oz 1.66–3.67 oz	11 fl oz 3.67 oz	12	30	For most crops that are not on the label, a 120-day plant-back interval must be observed. Highly toxic to foraging bees.
	4D	<b>Sivanto Prime</b> (flupyradifurone)	<b>Foliar:</b> 7–14 fl oz <b>Soil:</b> 21–28 fl oz	<b>Foliar:</b> 28 fl oz <b>Soil:</b> 28 fl oz per crop season	4	<b>Foliar:</b> 1 <b>Soil:</b> 21	<b>Foliar:</b> Minimum application interval: 7 days <b>Soil:</b> Soil maximum: no more than 3 crop seasons per year.
	9B	<b>PQZ</b> (pyrifluquinazon)	2.4 to 3.2 fl oz	9.6 fl oz	12	7	7 days before reapplication
	9B	<b>Fulfill</b> (pymetrozine)	2.75 oz	5.5 oz	12	0	Minimum of 7 days between applications.
	9D	<b>Sefina</b> (afidopyropen)	3.0 fl oz	28 fl oz	12	0	Minimum of 7 days between applications. Do not make more than 2 consecutive applications of Sefina. Maintain a 10-foot vegetative filter strip of permanent vegetation between the field edge and aquatic habitat.
	28	<b>Exirel</b> (cyantraniliprole)	13.5–20.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazypyr or cyantraniliprole-containing products per crop, either as foliar or soil applications.

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	28	<b>Verimark</b> (cyantraniliprole)	<b>At planting:</b> 10–13.5 fl oz drip <b>Chemigation:</b> 10 fl oz	<b>At planting:</b> 13.5 fl oz drip <b>Chemigation:</b> 20 fl oz (or 10 fl oz if at plant also used)	4	1	Both at plant and drip chemigation. Do not apply more than 2 drip chemigation applications (one if at-plant application used).
	29	<b>Beleaf 50 SG</b> (flonicamid)	2.0–2.8 oz	8.4 oz	12	0	Begin applications before pests reach damaging levels.
	-	<b>Azanguard</b> (azadirachtin)	10–16 fl oz		4	0	Use in combination with 0.25%–1.0% nonphytotoxic crop oil. OMRI-listed <sup>2</sup> .
	-	<b>Azatin O</b> (azadirachtin)	10–16 fl oz	16 fl oz	4	0	Antifeedant, repellent, insect growth regulator. Rate depends on pest—see label.
	-	<b>M-Pede</b> (potassium salts of fatty acids)	1%–2% v/v		12	0	OMRI-listed <sup>2</sup> . Do not apply to stressed plants (high heat or drought conditions).
	-	<b>Grandevo</b> ( <i>Chromobacterium subsugae</i> )	2–3 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
	-	<b>SunSpray 98.8%, Ultra-Saf-T-Side, Others</b> Oil, Insecticidal	1–2 gal/100 gal <b>JMS:</b> 3–6 qt/100 gal (JMS)		4	0	Organic Stylet-Oil and Saf-T-Side are OMRI-listed <sup>2</sup> .
<b>Beetles</b> (including cucumber beetle and flea beetle)	1A	<b>Sevin 4F, XLR</b> (carbaryl)	<b>4F, XLR:</b> 0.5–1.0 qt	<b>4F, XLR:</b> 6 qt	12	3	Do not apply when plants are wet. Use of carbaryl has been reported to flare (increase) mites. Do not apply to plants in bloom.
	1B	<b>Malathion 5EC</b> <b>Malathion 8</b> (malathion)	<b>5EC:</b> cucumber: 1.5–2.8 pt summer squash: 2.0–2.8 pt winter squash: 1.6 pt watermelon: 1.5–2.5 pt melon: 1.6 pt <b>8:</b> summer squash: 1.75 pt winter squash: 1.75 pt cucumber: 1.75 pt	<b>5EC:</b> cucumber: 5.6 pt summer squash: 8.4 pt winter squash: 4.8 pt watermelon: 5 pt <b>8:</b> cucumber: 3.5 pt summer squash: 5.25 pt winter squash: 5.25 pt	24	1	<b>5EC:</b> Maximum number of applications per year is 2. <b>8: Squash and cucumbers only</b>
	3A	<b>*Brigade 2 EC</b> (bifenthrin)	2.6–6.4 fl oz	19.2 fl oz	12	3	Do not make more than 2 applications after bloom. Highly toxic to foraging bees.
	3A	<b>*Baythroid XL</b> (beta-cyfluthrin)	2.4–2.8 fl oz	11.2 fl oz	12	0	Can be used on all cucurbits in Crop Group 9. Highly toxic to foraging bees.
	3A	<b>*Capture LFR</b> (bifenthrin)	6.8–8.5 fl oz, at planting	8.5 fl oz	12	n/a—applied at planting	At planting, banded over open furrow or in-furrow with seed or transplant.
	3A	<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees.
	3A	<b>*Pounce 25 WP</b> (permethrin)	6.4–12.8 oz	76.8 oz, 51.2 for cantaloupe	12	0	Use high rate for aphids and squash bug. Highly toxic to foraging bees.
	3A	<b>*Warrior II</b> (lambda-cyhalothrin)	1.28–1.92 fl oz	11.5 fl oz	24	1	Do not apply to blooming crops.
	4A	<b>Admire Pro</b> (imidacloprid) (see appropriate labels for other brands)	7–10.5 oz	10.5 fl oz	12	21	<b>Not for planthouse</b>
	4A	<b>Actara</b> (thiamethoxam)	1.5–3.0 oz	11.0 oz	12	0	Apply before pests reach damaging levels. Highly toxic to foraging bees.

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	4A	<b>Assail 305G</b> (acetamiprid)	2.5–5.3 oz	26.5 oz	12	0	No more than 5 applications per season. Toxic to foraging bees.
	4A	<b>Belay Insecticide</b> (clothianidin)	<b>Foliar:</b> 3.0–4.0 fl oz <b>Soil:</b> 9.0–12.0 fl oz	12 fl oz	12	<b>Foliar:</b> Do not make application after 4th true leaf on main stem is unfolded. <b>Soil:</b> 21	Do not apply when plants are blooming.
	4A	<b>Scorpion 355L Insecticide</b> (dinotefuran)	<b>Foliar:</b> 2–7 fl oz <b>Soil:</b> 9–10.5 fl oz	<b>Foliar:</b> 10.5 fl oz <b>Soil:</b> 21 fl oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Use only one application method (soil or foliar). Highly toxic to foraging bees for more than 38 hours after application. See label.
	4A	<b>Venom Insecticide</b> (dinotefuran)	<b>Foliar:</b> 1–4 oz	6 oz	12	<b>Foliar:</b> 1	Highly toxic to foraging bees. Use only foliar application for beetles. Note that pests controlled differ depending on application method.
	15	<b>Rimon 0.83EC</b> (novaluron)	9–12 fl oz	36 fl oz	12	1	Do not apply more often than every 14 days.
	28	<b>Exirel</b> (cyantraniliprole)	13.5–20.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazypyr- or cyantraniliprole-containing products per crop, either as foliar or soil applications.
	-	<b>Grandevo</b> ( <i>Chromobacterium subtsugae</i> )	2–3 lb		4	0	
	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	
<b>Caterpillars</b> (including armyworms, cabbage looper, corn earworm, cutworm, melonworm, pickleworm)	1A	<b>Sevin 4F, XLR</b> (carbaryl)	<b>4F, XLR:</b> 0.5–1.0 qt	<b>4F, XLR:</b> 6 qt	12	3	Do not apply when plants are wet. Use of carbaryl has been reported to flare (increase) mites. Do not apply to plants in bloom. Pickleworm and melonworm only.
	1B	<b>Malathion 5EC</b> <b>Malathion 8</b> (malathion)	<b>5EC:</b> cucumber: 1.5–2.8 pt summer squash: 2.0–2.8 pt winter squash: 1.6 pt watermelon: 1.5–2.5 pt melon: 1.6 pt <b>8:</b> summer squash: 1.75 pt winter squash: 1.75 pt cucumber: 1.75 pt	<b>5EC:</b> cucumber: 5.6 pt summer squash: 8.4 pt winter squash: 4.8 pt watermelon: 5 pt <b>8:</b> cucumber: 3.5 pt summer squash: 5.25 pt winter squash: 5.25 pt	24	1	<b>5EC:</b> Maximum number of applications per year is 2. <b>8:</b> Squash and cucumbers only
	3A	<b>*Ambush 25W</b> (permethrin)	6.4–12.8 oz	1.2 lb (0.8 lb for cantaloupe)	12	0	Do not apply more than 8 applications at the highest rate per season. Highly toxic to bees.
	3A	<b>*Asana XL (0.66 EC)</b> (esfenvalerate)	5.8–9.6 fl oz	48 fl oz	12	3	Do not apply more than 5 applications at high rate. For cucumber, melons, pumpkin, summer and winter squash. Highly toxic to foraging bees.
	3A	<b>*Athena</b> (bifenthrin)	7–17 fl oz	51 fl oz	12	7	Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
	3A	<b>*Baythroid XL</b> (beta-cyfluthrin)	0.8–2.8 fl oz	11.2 fl oz	12	0	Can be used on all cucurbits in Crop Group 9. Highly toxic to foraging bees.
	3A	<b>*Brigade 2 EC</b> (bifenthrin)	2.6–6.4 fl oz	19.2 fl oz	12	3	Do not make more than 2 applications after bloom. Highly toxic to foraging bees.
	3A	<b>*Capture LFR</b> (bifenthrin)	3.4–6.8 fl oz, at planting	8.5 fl oz	12	n/a—applied at planting	At planting, banded over open furrow or in-furrow with seed or transplant.
	3A	<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees.
	3A	<b>*Mustang</b> (zeta-cypermethrin)	3.0–4.3 oz	25.8 fl oz	7	12	Do not make applications less than 7 days apart.



Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	3A	<b>*Pounce 25 WP</b> (permethrin)	6.4–12.8 oz 76.8 oz	<b>Cantaloupe:</b> 51.2 oz	12	0	Use high rate for aphids and squash bug. Highly toxic to foraging bees.
	3A	<b>*Warrrior II</b> (lambda-cyhalothrin)	1.28–1.92 fl oz	11.5 fl oz	24	1	Do not apply to blooming crops.
	4A	<b>Assail 30SG</b> (acetamiprid)	2.5–5.3 oz	26.5 oz	12	0	No more than 5 applications per season. Do not use with other group 4A insecticides. Toxic to foraging bees.
	5	<b>Radiant SC</b> (spinetoram)	5–10 fl oz	34 fl oz	4	3; 1 for cucumbers	No more than 6 applications. Toxic to bees for up to 3 hours following application.
	11A	<b>Agree WG</b> ( <i>Bacillus thuringiensis</i> subspecies <i>aizawai</i> )	0.5–2.0 lb		4	0	Apply when larvae are small for best control. OMRI-listed <sup>2</sup> .
	11A	<b>Biobit HP</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	0.5–2.0 lb		4	0	Treat when larvae are young. Good coverage is essential. Can be used in the greenhouse. OMRI-listed <sup>2</sup> .
	11A	<b>Crymax WDG</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	0.5–2.0 lb		4	0	Use high rate for armyworms. Treat when larvae are young. Genetically engineered strain of <i>Bt</i> containing toxins from both <i>Bt kurstaki</i> and <i>Bt aizawai</i> strains. Not for organic production.
	11A	<b>Deliver</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	0.25–1.5 lb		4	0	Use higher rates for armyworms. OMRI-listed <sup>2</sup> .
	11A	<b>DiPel DF</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	0.5–2.0 lb		4	0	Treat when larvae are young. Good coverage is essential. For organic production.
	11A	<b>Javelin WG</b> ( <i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> )	0.12–1.50 lb		4	0	Treat when larvae are young. Thorough coverage is essential. Use higher rates for armyworms. OMRI-listed <sup>2</sup> .
	11A	<b>Xentari DF</b> ( <i>Bacillus thuringiensis</i> subspecies <i>aizawai</i> )	0.5–2.0 lb		4	0	Treat when larvae are young. Thorough coverage is essential. May be used in the greenhouse. Can be used in organic production.
	15	<b>Rimon 0.83EC</b> (novaluron)	12 fl oz	36 fl oz	12	1	Do not apply more often than every 14 days.
	18	<b>Intrepid 2F</b> (methoxyfenozide)	4–10 fl oz	64 fl oz	4	3	Do not make more than 4 applications per season.
	22	<b>Avaunt</b> (indoxacarb)	2.5–6.0 oz	24 oz	12	3	Highly toxic to foraging bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area. The minimum interval between sprays is 5 days.
	28	<b>Coragen</b> (chlorantraniliprole)	<b>Drip:</b> 2.0–7.5 fl oz soil <b>At planting:</b> 3.5–7.5 <b>Foliar:</b> 2.0–7.0	15.4 fl oz (10 fl oz for combined at-plant and drip chemigation soil application)	4	1	May be applied through drip (chemigation), as well as to soil at planting, or as a foliar spray.
	28	<b>Exirel</b> (cyantraniliprole)	7.0–13.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazypyr- or cyantraniliprole-containing products per crop, either as foliar or soil applications.
	28	<b>Verimark</b> (cyantraniliprole)	<b>At planting:</b> 6.75–13.5 fl oz drip <b>Chemigation:</b> 5–10 fl oz	<b>At planting:</b> 13.5 fl oz drip <b>Chemigation:</b> 20 fl oz (or 10 fl oz if at plant also used)	4	1	Both at plant and drip chemigation. Do not apply more than 2 drip chemigation applications (one if an at-plant application is used).
	-	<b>Grandevo</b> ( <i>Chromobacterium subsugae</i> )	2–3 lb		4	0	OMRI-listed <sup>2</sup> . Can be tank-mixed with a contact insecticide for improved control of targeted pests.

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	OMRI-listed <sup>2</sup> . Can be tank-mixed with a contact insecticide for improved control of targeted pests.
<b>Fire ants</b>	7A	<b>Extinguish</b> (S-methoprene)	0.75 lb		4	0	Slow-acting IGR (insect growth regulator). Best applied early spring and fall where crop will be grown. Colonies will be reduced after three weeks and eliminated after 8 to 10 weeks. May be applied by ground equipment or aerially.
	7C	<b>Esteem Ant Bait</b> (pyriproxyfen)	1.5–2.0 lb	2.0 lb	12	1	Apply when ants are actively foraging.
<b>Leafminers</b>	1B	<b>Dimethoate 4 EC</b> (dimethoate)	<b>Melons:</b> 1 pt <b>Watermelons:</b> 0.5–1 pt	2 pt per year	48	3	Highly toxic to bees. <b>Not for squash or cucumber.</b>
	3A	<b>*Ambush 25W</b> (permethrin)	6.4–12.8 oz	1.2 lb (0.8 lb for cantaloupe)	12	0	Do not apply more than 8 applications at the highest rate per season. Highly toxic to bees.
	3A	<b>*Athena</b> (bifenthrin)	13.5–17 fl oz	51 fl oz	12	7	Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
	3A	<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees.
	4A	<b>Scorpion 355L Insecticide</b> (dinotefuran)	<b>Foliar:</b> 2–7 fl oz <b>Soil:</b> 9–10.5 fl oz	<b>Foliar:</b> 10.5 fl oz <b>Soil:</b> 21 fl oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Use only one application method (soil or foliar). Do not use with other Group 4A insecticides. Highly toxic to foraging bees for more than 38 hours after application. See label.
	4A	<b>Belay Insecticide</b> (clothianidin)	<b>Foliar:</b> 3.0–4.0 fl oz <b>Soil:</b> 9.0–12.0 fl oz	12 fl oz	12	<b>Foliar:</b> Do not make application after 4th true leaf on main stem is unfolded <b>Soil:</b> 21	Do not apply when plants are blooming.
	4A	<b>Venom Insecticide</b> (dinotefuran)	<b>Foliar:</b> 1–4 oz <b>Soil:</b> 5–7.5 oz	<b>Foliar:</b> 6 oz <b>Soil:</b> 12 oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Highly toxic to foraging bees. Use only one application method (soil or foliar). Note that pests controlled differ depending on application method.
	5	<b>Radiant SC</b> (spinetoram)	5–10 fl oz	34 fl oz	4	3	No more than 6 applications. Toxic to bees for up to 3 hours following application.
	6	<b>*Agri-Mek SC</b> (abamectin)	1.75–3.50 fl oz	10.25 fl oz	12	7	No more than 2 sequential applications. <b>Must</b> be mixed with an adjuvant—see label for types. Highly toxic to foraging bees.
	15	<b>Rimon 0.83EC</b> (novaluron)	9–12 fl oz	36 fl oz	12	1	Do not apply more often than every 14 days.
	17	<b>Trigard</b> (cyromazine)	2.66 oz	15.96 oz	12	0	Do not make more than six applications.
	28	<b>Coragen</b> (chlorantraniliprole)	<b>Drip:</b> 2.0–7.5 fl oz soil <b>At planting:</b> 3.5–7.5 <b>Foliar:</b> 2.0–7.0	15.4 fl oz (10 fl oz for combined at-plant and drip chemigation soil application)	4	1	May be applied through drip (chemigation), as well as to soil at planting, or as a foliar spray.
	28	<b>Exirel</b> (cyantraniliprole)	13.5–20.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazypyr- or cyantraniliprole-containing products per crop, either as foliar or soil applications.
	28	<b>Verimark</b> (cyantraniliprole)	<b>At planting:</b> 6.75–13.5 fl oz drip <b>Chemigation:</b> 5–10 fl oz	<b>At planting:</b> 13.5 fl oz drip <b>Chemigation:</b> 20 fl oz (or 10 fl oz if at-plant also used)	4	1	Both at plant and drip chemigation. Do not apply more than 2 drip chemigation applications (one if an at-plant application is used).
	-	<b>Azaguard</b> (azadirachtin)	10–16 fl oz		4	0	Use in combination with 0.25%–1.0% nonphytotoxic crop oil. OMRI-listed <sup>2</sup> .

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
Mites	1B	<b>Malathion 8C</b> <b>Malathion 5C</b> (malathion)	<b>5EC:</b> cucumber: 1.5–2.8 pt summer squash: 2.0–2.8 pt winter squash: 1.6 pt watermelon: 1.5–2.5 pt melon: 1.6 pt <b>8:</b> summer squash: 1.75 pt winter squash: 1.75 pt cucumber: 1.75 pt	<b>5EC:</b> cucumber: 5.6 pt summer squash: 8.4 pt winter squash: 4.8 pt watermelon: 5 pt <b>8:</b> cucumber: 3.5 pt summer squash: 5.25 pt winter squash: 5.25 pt	24	1	<b>5EC:</b> Maximum number of applications per year is 2 <b>8: Squash and cucumbers only</b>
	3A	<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees.
	3A	<b>*Athena</b> (bifenthrin)	13.5–17 fl oz	51 fl oz	12	7	Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
	6	<b>*Agri-Mek SC</b> (abamectin)	1.75–3.50 fl oz	10.25 fl oz	12	7	No more than 2 sequential applications. <b>Must</b> be mixed with an adjuvant—see label for types. Highly toxic to foraging bees.
	10B	<b>Zeal</b> (etoxazole)	2.0–3.0 oz	3.0 oz <b>Cucumbers:</b> 6 oz	12	7	Apply when populations are low. One application per season, except two for cucumbers (supplemental label).
	20B	<b>Kanemite 15 SC</b> (acequinocyl)	31 fl oz	62 fl oz	12	1	Do not apply more than twice per season. Do not apply by air. <b>Melons, watermelon, and cucumbers only.</b>
	20D	<b>Acramite-50WS</b> (bifenazate)	0.75–1.0 lb	1.0 lb	12	3	One application per season.
	21A	<b>Portal XLO</b> (fenpyroximate)	1.0–2.0 pt	4.0 pt	12	3 <b>Cucumbers: 1</b>	<b>All melons, watermelon, and cucumbers.</b> Two applications per season. Effective for broad mite control.
	23	<b>Oberon 25C</b> (spiromesifen)	7.0–8.5 fl oz	25.5 fl oz	12	7	No more than 3 applications. See label for plant-back intervals.
	-	<b>Grandevo</b> ( <i>Chromobacterium subsugae</i> )	2–3 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
	-	<b>Azaguard</b> (azadirachtin)	10–16 fl oz		4	0	Use in combination with 0.25%–1.0% nonphytotoxic crop oil. OMRI-listed <sup>2</sup> .
	Squash bug, leaf-footed bug	1A	<b>Sevin 4F, XLR</b> (carbaryl)	<b>4F, XLR:</b> 0.5–1.0 qt	<b>4F, XLR:</b> 6 qt	12	3
3A		<b>*Ambush 25W</b> (permethrin)	6.4–12.8 oz	1.2 lb (0.8 lb for cantaloupe)	12	0	Do not apply more than 8 applications at the highest rate per season. Highly toxic to bees.
3A		<b>*Athena</b> (bifenthrin)	13.5–17 fl oz	51 fl oz	12	7	Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
3A		<b>*Brigade 2 EC</b> (bifenthrin)	2.6–6.4 fl oz	19.2 fl oz	12	3	Do not make more than 2 applications after bloom. Highly toxic to foraging bees.
3A		<b>*Danitol 2.4 EC</b> (fenpropathrin)	10.67–16 fl oz	42.67 fl oz	24	7	Highly toxic to foraging bees.
3A		<b>*Pounce 25 WP</b> (permethrin)	6.4–12.8 oz	76.8 oz <b>Cantaloupe:</b> 51.2 oz	12	0	Use high rate for aphids and squash bug. Highly toxic to foraging bees.
3A		<b>*Warrior II</b> (lambda-cyhalothrin)	1.28–1.92 fl oz	11.5 fl oz	24	1	Do not apply to blooming crops.

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	4A	<b>Assail 30SG</b> (acetamiprid)	5.3 oz	26.5 oz	12	0	No more than 5 applications per season. Do not use with other group 4A insecticides. Application for squash bug are most effective against newly laid eggs and nymphs. Toxic to foraging bees.
	4A	<b>Belay Insecticide</b> (clothianidin)	<b>Foliar:</b> 3.0–4.0 fl oz <b>Soil:</b> 9.0–12.0 fl oz	12 fl oz	12	<b>Foliar:</b> Do not make application after 4th true leaf on main stem is unfolded <b>Soil:</b> 21	Do not apply when plants are blooming.
	4A	<b>Scorpion 35SL Insecticide</b> (dinotefuran)	<b>Foliar:</b> 2–7 fl oz <b>Soil:</b> 9–10.5 fl oz	<b>Foliar:</b> 10.5 fl oz <b>Soil:</b> 21 fl oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Use only one application method (soil or foliar). Do not use with other Group 4A insecticides. Highly toxic to foraging bees for more than 38 hours after application.
	4A	<b>Venom Insecticide</b> (dinotefuran)	<b>Foliar:</b> 1–4 oz <b>Soil:</b> 5–6 oz	<b>Foliar:</b> 6 oz <b>Soil:</b> 12 oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Highly toxic to foraging bees. Use only one application method (soil or foliar). Note that pests controlled differ depending on application method.
	4D	<b>Sivanto Prime</b> (flupyradifurone)	<b>Foliar:</b> 7–14 fl oz <b>Soil:</b> 21–28 fl oz	<b>Foliar:</b> 28 fl oz <b>Soil:</b> 28 fl oz per crop season	4	<b>Foliar:</b> 1 <b>Soil:</b> 21	<b>Foliar:</b> Minimum application interval: 7 days <b>Soil:</b> Soil maximum: no more than 3 crop seasons per year.
	15	<b>Rimon 0.83EC</b> (novaluron)	9–12 fl oz	36 fl oz	12	1	Do not apply more often than every 14 days.
	1B	<b>Dimethoate 4 EC</b> (dimethoate)	<b>Melons:</b> 1 pt <b>Watermelons:</b> 0.5–1 pt	2 pt per year	48	3	Highly toxic to bees. <b>Not for squash or cucumber.</b>
<b>Thrips</b> (check label for species controlled)	4A	<b>Admire Pro</b> (imidacloprid) (see appropriate labels for other brands)	7–10.5 oz	10.5 fl oz	12	<b>Soil:</b> 21	Do not use with other Group 4A insecticides. <b>Not for planthouse.</b> Only for foliage feeding thrips; will not control thrips feeding on flowers.
	4A	<b>Platinum 755G</b> (thiamethoxam)	5–11 fl oz <b>755G:</b> 1.66–3.67 oz	11 fl oz 3.67 oz	12	30	For most crops that are not on the label, a 120-day plant-back interval must be observed. Highly toxic to foraging bees.
	4A	<b>Scorpion 35SL Insecticide</b> (dinotefuran)	<b>Foliar:</b> 2–7 fl oz <b>Soil:</b> 9–10.5 fl oz	<b>Foliar:</b> 10.5 fl oz <b>Soil:</b> 21 fl oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Use only one application method (soil or foliar). Do not use with other Group 4A insecticides. Highly toxic to foraging bees for more than 38 hours after application. See label.
	4A	<b>Venom Insecticide</b> (dinotefuran)	<b>Foliar:</b> 1–4 oz <b>Soil:</b> 5–7.5 oz	<b>Foliar:</b> 6 oz <b>Soil:</b> 12 oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Highly toxic to foraging bees. Use only one application method (soil or foliar). Note that pests controlled differ depending on application method.
	5	<b>Radiant SC</b> (spinetoram)	6–10 fl oz	34 fl oz	4	<b>Cucumbers:</b> 1	No more than 6 applications. Toxic to bees for up to 3 hours following application.
	5	<b>Entrust SC</b> (spinosad)	6–8 fl oz	29 fl oz	1	4	No more than 5 applications per year. OMRI-listed.
	28	<b>Exirel</b> (cyantraniliprole)	13.5–20.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazapyr- or cyantraniliprole-containing products per crop, either as foliar or soil applications.
-	-	<b>Grandevo</b> ( <i>Chromobacterium subtsugae</i> )	2–3 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
-	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
-	-	<b>Azaguard</b> (azadirachtin)	10–16 fl oz		4	0	Use in combination with 0.25%–1.0% nonphytotoxic crop oil. OMRI-listed <sup>3</sup> .

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
Whitefly	3A	<b>*Athena</b> (bifenthrin)	13.5–17 fl oz	51 fl oz	12	7	Only for adults. Do not make more than 2 consecutive applications. Do not apply more than 2 applications after bloom.
	3A	<b>*Brigade 2 EC</b> (bifenthrin)	2.6–6.4 fl oz	19.2 fl oz	12	3	Do not make more than 2 applications after bloom. Highly toxic to foraging bees.
	4A	<b>Actara</b> (thiamethoxam)	1.5–5.5 oz	11.0 oz	12	0	Apply before pests reach damaging levels. Highly toxic to foraging bees.
	4A	<b>Admire Pro</b> (imidacloprid) (see appropriate labels for other brands)	7–10.5 oz <b>Planthouse:</b> 0.44 fl oz/10,000 plants	10.5 fl oz	12	<b>Soil:</b> 21	Do not use with other Group 4A insecticides. <b>Planthouse:</b> One application to transplants. See label for use on mature greenhouse cucumbers.
	4A	<b>Assail 30SG</b> (acetamiprid)	2.5–5.3 oz	26.5 oz	12	0	No more than 5 applications per season. Do not use with other group 4A insecticides. Toxic to foraging bees.
	4A	<b>Platinum 755G</b> (thiamethoxam)	5–11 fl oz <b>755G:</b> 1.66–3.67 oz	11 fl oz 3.67 oz	12	30	For most crops that are not on the label, a 120-day plant-back interval must be observed. Highly toxic to foraging bees.
	4A	<b>Scorpion 355L Insecticide</b> (dinotefuran)	<b>Foliar:</b> 2–7 fl oz <b>Soil:</b> 9–10.5 fl oz	<b>Foliar:</b> 10.5 fl oz <b>Soil:</b> 21 fl oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Use only one application method (soil or foliar). Do not use with other Group 4A insecticides. Highly toxic to foraging bees for more than 38 hours after application. See label.
	4A	<b>Venom Insecticide</b> (dinotefuran)	<b>Foliar:</b> 1–4 oz <b>Soil:</b> 5–7.5 oz	<b>Foliar:</b> 6 oz <b>Soil:</b> 12 oz	12	<b>Foliar:</b> 1 <b>Soil:</b> 21	Highly toxic to foraging bees. Use only one application method (soil or foliar). Note that pests controlled differ depending on application method.
	4D	<b>Sivanto Prime</b> (flupyradifurone)	<b>Foliar:</b> 10.5–14 fl oz <b>Soil:</b> 21–28 fl oz	<b>Foliar:</b> 28 fl oz <b>Soil:</b> 28 fl oz per crop season	4	<b>Foliar:</b> 1 <b>Soil:</b> 21	<b>Foliar:</b> Minimum application interval: 7 days. <b>Soil:</b> Soil maximum: no more than 3 crop seasons per year.
	7C	<b>Knack IGR</b> (pyriproxyfen)	8–10 fl oz	20 fl oz	12	7	Immatures only. Apply when nymphs first appear. Do not apply more than twice per growing season.
	9B	<b>PQZ</b> (pyrifluquinazon)	3.2 fl oz	9.6 fl oz	12	7	7 days before reapplication
	9B	<b>Fulfill</b> (pymetrozine)	2.75 oz	5.5 oz	12	0	Minimum of 7 days between applications.
	9D	<b>Sefina</b> (afidopyrophen)	14 fl oz	28 fl oz	12	0	Minimum of 7 days between applications. Maintain a 10-foot vegetative filter strip of permanent vegetation between the field edge and aquatic habitat.
	15	<b>Rimon 0.83EC</b> (novaluron)	9–12 fl oz	36 fl oz	12	1	Do not apply more often than every 14 days.
	16	<b>Courier 40SC</b> (buprofezin)	9.0–13.6 fl oz	27.2 fl oz	12	7	Immatures only. Insect growth regulator. Do not make more than 2 applications per season per crop or 4 per year.
	23	<b>Oberon 25C</b> (spiromesifen)	7.0–8.5 fl oz	25.5 fl oz	12	7	No more than 3 applications.
	28	<b>Exirel</b> (cyantraniliprole)	13.5–20.5 fl oz	61.6 fl oz	12	1	Do not apply a total of more than 0.4 lb a.i. per acre of cyazypyr- or cyantraniliprole-containing products per crop, either as foliar or soil applications.
	28	<b>Verimark</b> (cyantraniliprole)	6.75–13.5 fl oz	<b>At planting:</b> 13.5 fl oz drip <b>Chemigation:</b> 20 fl oz (or 10 fl oz if at-plant also used)	4	1	Both at plant and drip chemigation. Do not apply more than 2 drip chemigation applications (one if at-plant application is used).
-	<b>Grandevo</b> ( <i>Chromobacterium subsugae</i> )	2–3 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.	

Insect or Mite Pest	MOA Code <sup>1</sup>	Trade Name (Active Ingredient) *Restricted	Rate (Product/Acre)	Max Rate Product per Season	REI (Hours)	Days to Harvest	Remarks <sup>2</sup>
	-	<b>M-Pede</b> (potassium salts of fatty acids)	1%–2% v/v		12	0	OMRI-listed <sup>2</sup> . Do not apply to stressed plants (high heat or drought conditions).
	-	<b>Venerate</b> ( <i>Burkholderia rinojensis</i> )	2–4 lb		4	0	OMRI-listed. Can be tank-mixed with a contact insecticide for improved control of targeted pests.
	1B	<b>*Diazinon 50 W, *AG500</b> (diazinon)	<b>AG500:</b> 2–4 qt <b>50W:</b> 4–8 lb	4 qt 8 lb	72	preplant	<b>Melons and watermelons only.</b> Not for squash or cucumbers. One application per year.
	3A	<b>*Capture LFR</b> (bifenthrin)	3.4–8.5 fl oz, at planting	8.5 fl oz	12	n/a—applied at planting	At planting, banded over open furrow or in-furrow with seed or transplant.

<sup>1</sup> Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 9.4, March 2020. Number codes (1 through 29) are used to distinguish the main insecticide mode of action groups, with additional letters for certain subgroups within each main group. All insecticides within the same group (with the same number) indicate the same active ingredient or similar mode of action. This information must be considered for the insecticide resistance management decisions. - = unknown, or a mode of action that has not been classified yet.

<sup>2</sup> Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned. OMRI-listed: Listed by the Organic Materials Review Institute for use in organic production.

\* Restricted use insecticide.

**Table 11. Cucurbit fungicides ordered by disease and then FRAC group according to their mode of action.**

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to Harvest	Days to Reentry	Remarks <sup>2</sup>
			Appl.	Season			
<b>Please refer to Chapter 19 for information on biopesticides, including materials labeled for certified organic production.</b>							
<b>Alternaria leaf spot/blight</b>	M1	(copper compounds) <b>Many brands available:</b> AmeriCop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Instill, Kalmor, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, Kocide HCU, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75WG, Nu-Cop 3L, Nu-Cop 50DF, Nu-Cop 50 WP, Nu-Cop HB, Nu-Cop 30 HB, Nu-Cop XLR, Phyton 27 AG, Previsto	See label	See label	1	Varies from 4 hr to 2 days	Check label for specific guidelines.
	M1 & M3	<b>ManKocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75 WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for specific guidelines.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo Zn, EQUUS 720 SST, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Praiz, Rialto 720F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot and other environmental conditions as indicated on the label.
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	P4	<b>Vacciplant</b> (laminarin)	22 fl oz	See label	0	4 hr	Initiate as preventative application.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	3	<b>Mettle 125 ME</b> (tetraconazole)	8 fl oz	24 fl oz/year	1	0.5	Make no more than 2 sequential applications before rotating to another fungicide with alternate mode of action.
	3 & 9	<b>Inspire Super</b> (cyprodinil + difenoconazole)	20 fl oz	80 fl oz/year	7	0.5	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber.
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	3.25 pt	See label	0	2	Limit is 4 appl./crop
	7	<b>Endura</b> (boscalid)	6.5 oz	26 oz/year	0	0.5	
	7	<b>Fontelis</b> (penthiopyrad)	16 fl oz	67 fl oz/year	1	0.5	Maximum rate is for year. Do not make more than 2 appl. sequentially with any FRAC 7 containing fungicide. See label for guidelines.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	
	7 & 3	<b>Luna Experience</b> (fluopyram + tebuconazole)	17 fl oz	34 fl oz/year	7	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 3 fungicide.
	7 & 11	<b>Luna Sensation</b> (fluopyram + trifloxystrobin)	7.6 fl oz	27.1 fl oz/year	0	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 11 fungicide.
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	See label.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	See label.
	9	<b>Vango WG</b> (cyprodinil)	7 oz	28 oz/year	1	0.5	Alternate with fungicide in another FRAC group after 2 applications.
	12	<b>Emblem</b> (fludioxonil)	7 fl oz	28 fl oz/year	1	0.5	
	7 & 12	<b>Miravis Prime</b> (pydiflumetofen + fludioxonil)	11.4 fl oz	22.8 fl oz/year	1	0.5	Do not make more than two applications of Miravis Prime or other Group 7 and 12 before alternation with a fungicide that is not in Group 7 or 12.
	9 & 12	<b>Alterity 62.5 Switch 62.5WG</b> (cyprodinil + fludioxonil)	14 oz	56 oz/year	1	0.5	Do not make more than two consecutive applications.
	11	<b>Reason 500 SC</b> (fenamidone)	5.5 fl oz	22 fl oz/year	14	0.5	Do not make more than one application of Reason 500 SC Fungicide before alternating with a fungicide from a different resistance management group.
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 2 SC, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxzone, AZteroid FC, Azteroid FC 3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11	<b>Cabrio EG</b> (pyraclostrobin)	16 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	<b>Evito 480 SC Tepera</b> (fluoastrobilin)	See label	See label	See label	See label	
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV</b> <b>Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all QoI fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-QoI-containing fungicide.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers.
	12	<b>Emblem Spirato</b> (fludioxonil)	7 fl oz	28 fl oz/year	1	0.5	
	19	<b>Affirm WDG PH-D</b> (polyoxin D zinc salt)	6.2 oz	See label	0	4 hr	
	22 & M3	<b>Gavel 75DF</b> (zoxamide + mancozeb)	2 lb	16 lb/year	5	2	Limit is 8 appl./crop. Some cantaloupe are sensitive. See label.
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	27 & 11	<b>Tanos</b> (cymoxanil + famoxadone)	8 oz	See label	3	0.5	Limit is 4 appl./crop. Must tank-mix with a contact fungicide.
	29	<b>Omega 500 F</b> (fluzinam)	24 fl oz	See label	7	0.5	See label for crop-specific guidelines.
	33	<b>K-Phite Fungi-Phite Reliant</b> (phosphorous acid and salts)	See label	See label	0	4 hr	Check label for guidelines on foliar and aerial application, as well as preventative and control programs. Do not apply at less than 3 days interval. Check label to see whether it is suppression of disease only.
<b>Angular leaf spot</b>	M1	(copper compounds) <b>Many brands available:</b> AmeriCop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION+++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Instill, Kalmor, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75WG, Nu-Cop 50 WP, Nu-Cop 3L, Nu-Cop HB, Nu-Cop 30 HB, Nu-Cop 50DF, Nu-Cop XLR, Phyton 27 AG, Previsto	See label	See label	1	Varies from 4 hr to 2 days	Check label for crop-specific guidelines.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. See label for details.
	P1	<b>Actigard 50WG</b> (acibenzolar-S-methyl)	1 oz	8 oz/year	0	0.5	Apply preventatively prior disease development. Suppression of disease.



Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
Anthracnose	M1	(copper compounds) <b>Many brands available:</b> AmeriCop 40 DF, Badge SC, Badge X2, Basic Copper 53, C-O-C-S WDG, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION+, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Instill, Kaimor, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75WG, Nordox 50WP, Nu-Cop 3L, Nu-Cop 30 HB, Nu-Cop 50DF, Nu-Cop HB, Nu-Cop XLR, Phytos 27 AG, Previsto, Stretch	See label	See label	1	Varies from 4 hr to 2 days	Check label for crop-specific guidelines.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. See label for details.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for crop-specific guidelines.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Praiz, Rialto 720 F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	Limit is 4 appl./crop.
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	3.25 pt	See label	0	2	Follow resistance management guidelines on label. Check label for additional guidelines.
	1	(thiophanate-methyl) <b>Many brands available:</b> 3336 EG, 3336 F, 3336 WP, Cercobin, Incognito 4.5F, Incognito 85 WDG, T-Methyl 4.5 AG, T-Methyl 4.5 F, T-Methyl 70 WP, Talaris 4.5F, Talaris 70 WSB, Thiophanate methyl 85 WDG, TM 4.5 F, Topsin 4.5 FL, Topsin M WSB	See label	See label	1	1	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber. See label for guidelines.
	3 & 9	<b>Inspire Super</b> (cyprodinil + difenoconazole)	20 fl oz	80 fl oz/year	7	0.5	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 3	<b>Luna Experience</b> (flupyryram + tebuconazole)	17 fl oz	34 fl oz/year	7	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 3 fungicide.
	7 & 11	<b>Luna Sensation</b> (flupyryram + trifloxystrobin)	7.6 fl oz	27.1 fl oz/year	0	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 11 fungicide.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	See label.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	See label.
	11	<b>Cabrio EG</b> (pyraclostrobin)	16 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	(azoxystrobin) <b>Many brands available:</b> Acadia 2 SC, A-Zox 25SC, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxystar, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplant production. See label for details.
	11	<b>Evito 480 SC</b> <b>Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV</b> <b>Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all Qol fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-Qol-containing fungicide.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers.
	19	<b>Oso 5% SC</b> (polyoxin D zinc salt)	13.0 fl oz	See label	0	4 hr	
	27 & 11	<b>Tanos</b> (cymoxanil + famoxadone)	8 oz	See label	3	0.5	Limit is 4 appl./crop. Must tank-mix with a contact fungicide. See label.
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
<b>Bacterial fruit blotch</b>	M1	(copper compounds) <b>Many brands available:</b> Americop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, Kocide HCU, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75 WG, Nu-Cop 50 WP, Nu-Cop 3L, Nu-Cop 30HB, Nu-Cop 50 DF, Nu-Cop HB, Nu-Cop, Previsto, Stretch	See label	See label	1	Varies from 4 hr to 2 days	Suppression only. Check label for crop-specific guidelines.
	P1	<b>Actigard 50WG</b> (acibenzolar-S-methyl)	1 oz	8 oz/year	0	0.5	Apply preventatively prior disease development. Suppression of disease.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. See label for details.
	27 & 11	<b>Tanos</b> (cymoxanil + famoxadone)	10 oz	See label	3	0.5	Limit is 4 appl./crop. Must tank-mix with a contact fungicide. Bacterial fruit blotch disease suppression only.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
Belly rot	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	1	(thiophanate-methyl) <b>Many brands available:</b> 3336 EG, 3336 F, 3336 WP, Cercobin, T-Methyl 4.5 AG, T-Methyl 4.5 F, T-Methyl 70 WP, Incognito 4.5F, Incognito 85WDG, Miramar, Talaris 70 WSB, Thiophanate methyl 85 WDG, TM 4.5 F, Topsin 4.5 FL, Topsin M WSB	See label	See label	1	1	Follow resistance management guidelines on label.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 3	<b>Luna Experience</b> (fluopyram + tebuconazole)	17 fl oz	34 fl oz/year	7	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 3 fungicide.
	7 & 11	<b>Luna Sensation</b> (fluopyram + trifloxystrobin)	7.6 fl oz	27.1 fl oz/year	0	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 11 fungicide.
	11	<b>Evito 480 SC</b> <b>Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 2 SC, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxzone, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV</b> <b>Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all Qol fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-Qol-containing fungicide.
	19	<b>PH-D</b> (polyoxin D zinc salt)	6.2 oz	See label	0	4 hr	
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for crop-specific guidelines.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Praiz, Rialto 720 F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	3.25 pt	See label	0	2	Limit is 4 appl./crop.
	3	<b>Mettle 125 ME</b> (tetraconazole)	8 fl oz	24 fl oz/year	1	0.5	Make no more than 2 sequential applications before rotating to another fungicide with alternate mode of action.
	3 & 9	<b>Inspire Super</b> (cyprodinil + difenoconazole)	20 fl oz	80 fl oz/year	7	0.5	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	See label.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	See label.
	7 & 12	<b>Miravis Prime</b> (pydiflometofen + fludioxonil)	11.4 fl oz	22.8 fl oz/year	1	0.5	Do not make more than two applications of Miravis Prime or other Group 7 and 12 before alternation with a fungicide that is not in Group 7 or 12.
	11	<b>Cabrio EG</b> (pyraclostrobin)	16 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	<b>Many brands available:</b> A-Zox 25SC, Acadia 25C, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxyzone, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop & alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11	<b>Evito 480 SC</b> <b>Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV</b> <b>Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all QoI fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-QoI-containing fungicide.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	22 & M3	<b>Gavel 75DF</b> (zoxamide + mancozeb)	2 lb	16 lb/year	5	2	Limit is 8 appl./crop. Some cantaloupe are sensitive. See label.
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	Check label for crop-specific guidelines.
<b>Downy mildew</b>	M1	(copper compounds) <b>Many brands available:</b> Americop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Kalmor, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, Kocide HCU, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75 WG, Nu-Cop 50 WP, Nu-Cop 3L, Nu-Cop 30HB, Nu-Cop 50 DF, Nu-Cop HB, Nu-Cop, Previsto, Stretch	See label	See label	1	Varies from 4 hr to 2 days	Check label for crop-specific guidelines.
	M1 & M2	<b>Top Cop w Sulfur</b> (copper + sulfur)	1 qt	See label	-	1	Do not apply if air temperatures greater than 90°F are expected.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for crop-specific guidelines.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Praiz, Rialto 720 F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	P1	<b>Actigard 50WG</b> (acibenzolar-S-methyl)	1 oz	8 oz/year	0	0.5	Apply preventatively prior disease development. Suppression of disease.
	P4	<b>Vacciplant</b> (laminarin)	22 fl oz	See label	0	4 hr	Initiate as preventative application.
	U15 & 40	<b>Orondis Ultra</b> (oxathiapiprolin + mandipropamid)	8 fl oz	32 fl oz/year	0	4 hr	Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. Check label for additional details.
	4 & M1	<b>Ridomil Gold Copper</b> (mefenoxam + copper hydroxide)	2 lb	8 lb	5	2	Limit is 4 appl./crop.
	4 & M3	<b>Ridomil Gold MZ WG</b> (mancozeb + mefenoxam)	2.5 lb	10 lb	5	2	Limit is 4 appl./crop. Rate is for downy mildew.
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	2.5 pt	See label	0	2	Limit is 4 appl./crop.
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	3 & 33	<b>Viathon</b> (tebuconazole + potassium phosphite)	4 pt	12 pt	7	0.5	Disease suppression only. See label.
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	See label.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	See label
	11	<b>Cabrio EG</b> (pyraclostrobin)	12 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	<b>Evito 480 SC</b> <b>Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11	<b>Flint Extra</b> <b>Flint</b> (trifloxystrobin)	See label	See label	See label	0.5	(Disease suppression) Limit is 4 appl./crop and alternate chemistry.
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 25C, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxzone, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11	<b>Reason 500SC</b> (fenamidone)	5.5 fl oz	22 fl oz	14	0.5	See label.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV</b> <b>Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all QoI fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-QoI-containing fungicide.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers.
	19	<b>Oso 5% SC</b> (polyoxin D zinc salt)	13.0 fl oz	See label	0	4 hr	
	21	<b>Ranman</b> <b>Ranman 400 SC</b> (cyazofamid)	2.75 fl oz	16.5 fl oz	0	0.5	Limit is 6 appl./crop. Follow resistance management guidelines.
	22	<b>Elumin</b> (ethaboxam)	8 fl oz	16 fl oz/year	1	0.5	Application interval should not be less than 14 days.
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	
	22 & M3	<b>Gavel 75DF</b> (zoxamide + mancozeb)	2 lb	16 lb/year	5	2	Limit is 8 appl./crop. Some cantaloupe are sensitive. See label.
	27	<b>Curzate 60DF</b> (cymoxanil)	5 oz	30 oz	3	0.5	Use only with a labeled rate of protectant fungicide.
	27 & 11	<b>Tanos</b> (cymoxanil + famoxadone)	8 oz	See label	3	0.5	Limit is 4 appl./crop. Must tank-mix with a contact fungicide.
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	28	<b>Previcur Flex Promess</b> (propamocarb hydrochloride)	1.2 pt	6 pt	2	0.5	See label for instructions.
	29	<b>Omega 500 F</b> (fluzinam)	24 fl oz	See label	7	0.5	See label for crop-specific guidelines.
	33	<b>Aliette 80WDG, Alude, Fosphite, Fungi-Phite, Prophyt, K-Phite 7 LP, Linebacker WDG, Reliant</b> (fosetyl-Al, phosphorous acid and salts)	See label	See label	See label	See label	See label for specific product guidelines.
	40	<b>Forum</b> (dimethomorph)	6 oz	30 oz	See label	0.5	Limit is 5 appl./crop. Apply with another fungicide that has a different mode of action and alternate.
	40	<b>Orondis Ultra B Revus</b> (mandipropamid)	8 fl oz	32 fl oz	0	4 hr	See label. For suppression only.
	43	<b>Presidio</b> (flupicolide)	4 fl oz	8 fl oz/year	2	0.5	Tank-mix with another fungicide product with a different FRAC code.
	45 & 40	<b>Zampro</b> (ametoctradin + dimethomorph)	14 fl oz	42 fl oz/year	0	0.5	No more than 2 sequential applications. Addition of a spreading/penetrating adjuvant is recommended.
	U15	<b>Orondis Opti A</b> <b>Orondis Ultra A</b>	4.8 fl oz	19.2 fl oz/year	0	4 hr	Cucurbit vegetables, crop group 9. See label.
<b>Gummy stem blight</b>	M1	(copper compounds) <b>Many brands available:</b> AmeriCop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Kalmor, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, Kocide HCU, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75WG, Nu-Cop 50 WP, Nu-Cop 3L, Nu-Cop 30 HB, Nu-Cop 50DF, Nu-Cop HB, Nu-Cop XLR, Previsto	See label	See label	1	Varies from 4 hr to 2 days	Check label for crop-specific guidelines.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for crop-specific guidelines.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Praiz, Rialto 720F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to Harvest	Min. Days to Reentry	Remarks <sup>2</sup>
			Appl.	Season			
	1	(thiophanate-methyl) <b>Many brands available:</b> 3336 EG, 3336 F, 3336 WP, Cercobin, T-Methyl 4.5 AG, T-Methyl 4.5 F, T-Methyl 70 WP, Incognito 4.5F, Incognito 85WDG, Miramar, Talaris 70 WSB, Thiophanate methyl 85 WDG, TM 4.5 F, Topsin 4.5 FL, Topsin M WSB	See label	See label	1	1	Follow resistance management guidelines on label.
	3	<b>Topguard</b> (flutriafol)	14 fl oz	56 fl oz/year	0	0.5	Check label.
	3	(tebuconazole) <b>Many brands available:</b> Buzz Ultra DF, Incognito 4.5F, Monsoon, Onset 3.6L, Orius 3.6F, Teb 3.6 SC, Tebu-Crop 3.6F, Tebu 3.6F, Tebuconazole 3.6 F, Tebustar 3.6L, Tebuzol 3.6F, Toledo, Vibe	See label	See label	1	1	Follow resistance management guidelines indicated on the label. Check label for additional guidelines.
	3	<b>Proline</b> (prothioconazole)	5.7 fl oz	17.1 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year.
	3	<b>Rhyme</b> (flutriafol)	7 fl oz	28 fl oz/year	0	0.5	Do not apply more than 2 sequential applications before rotating with a different FRAC group fungicide.
	3	<b>Mettle 125 ME</b> (tetraconazole)	8 fl oz	24 fl oz/year	1	0.5	Make no more than 2 sequential applications before rotating to another fungicide with alternate mode of action.
	3 & M5	<b>Muscle ADV</b> (tebuconazole + chlorothalonil)	2.1 pt	6.4 pt	7	0.5	Watermelon, squash, pumpkin, and melon
	3 & 33	<b>Viathon</b> (tebuconazole + potassium phosphite)	4 pt	12 pt	7	0.5	
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	2.5 pt	See label	0	2	Limit is 4 appl./crop.
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	7	<b>Endura</b> (boscalid)	6.5 oz	26 oz/year	0	0.5	Do not make more than one application before alternating to another labeled fungicide with a different mode of action for at least one application.
	7	<b>Fontelis</b> (penthiopyrad)	16 fl oz	67 fl oz/year	1	0.5	Do not make more than 2 applications sequentially.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 3	<b>Luna Experience</b> (fluopyram + tebuconazole)	17 fl oz	34 fl oz/year	7	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 3 fungicide.
	7 & 3	<b>Propulse</b> (fluopyram + prothioconazole)	13.6 fl oz	34.2 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year. See label for harvesting guidelines.
	7 & 11	<b>Luna Sensation</b> (fluopyram + trifloxystrobin)	7.6 fl oz	27.1 fl oz/year	0	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 11 fungicide.
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	Limit is 4 appl./crop and alternate chemistry.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	Do not make more than one application before alternating to a non-FRAC 7, non-FRAC 11 group fungicide. In areas where gummy stem blight resistance to FRAC 7 or 11 fungicides have been reported, tank-mix with chlorothalonil at full rate.
	3 & 9	<b>Inspire Super</b> (cyprodinil + difenoconazole)	20 fl oz	80 fl oz/year	7	0.5	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber.



Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
Phytophthora blight/fruit and root rot	9 & 12	<b>Altarity 62.5 Switch 62.5WG</b> (cyprodinil + fludioxonil)	14 oz	56 oz/year	1	0.5	Do not make more than two consecutive applications before switching to fungicide with a different mode of action.
	11	<b>Evito 480 SC Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	See label for tank-mixing restrictions.
	11	<b>Cabrio EG</b> (pyraclostrobin)	16 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 25C, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxzone, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11	<b>Narvos 50 WDG Sovran</b> (kresoxim-methyl)	4.8 oz	19.2 oz	0	0.5	Follow resistance management guidelines on label.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-QoI-containing fungicide
	11 & M5	<b>Arius ADV Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all QoI fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers
	12	<b>Emblem</b> (fludioxonil)	7 fl oz	28 fl oz/year	1	0.5	
	7 & 12	<b>Miravis Prime</b> (pydiflumetofen + fludioxonil)	11.4 fl oz	22.8 fl oz/year	1	0.5	Do not make more than two applications of Miravis Prime or other Group 7 and 12 before alternation with a fungicide that is not in Group 7 or 12.
	19	<b>Afirm WDG Oso 5% SC PH-D</b> (polyoxin D zinc salt)	See label	See label	0	4 hr	
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	29	<b>Omega 500 F</b> (fluazinam)	24 fl oz	See label	7	0.5	See label for crop-specific guidelines.
	4	<b>Orodis Gold B Ridomil Gold SL</b> (mefenoxam)	2 pt	See label	5	2	See label for application-specific guidelines (soil-injected or soil-incorporated, soil-directed and other foliar). Suppression only.
	4 & M1	<b>Ridomil Gold Copper</b> (mefenoxam + copper hydroxide)	2 lb	8 lb	5	2	Limit is 4 appl./crop.
	21	<b>Ranman 400 SC</b> (cyazofamid)	2.75 fl oz	16.5 fl oz	0	0.5	Limit is 6 appl./crop. Follow resistance management guidelines.
	22	<b>Elumin</b> (ethaboxam)	8 fl oz	16 fl oz/year	2	0.5	Application interval should not be less than 14 days.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	22 & M3	<b>Gavel 75DF</b> (zoxamide + mancozeb)	2 lb	16 lb/year	5	2	Limit is 8 appl./crop. Some cantaloupe are sensitive. See label.
	27 & 11	<b>Tanos</b> (cymoxanil + famoxadone)	10 oz	See label	3	0.5	Limit is 4 appl./crop. Must tank-mix with a contact fungicide. Phytophthora blight foliar and fruit phase only (disease suppression).
	29	<b>Omega 500 F</b> (fluzinam)	24 fl oz	See label	7	0.5	See label for crop-specific guidelines.
	33	<b>Alude, Aliette 80WDG, Fosphite, ProPhyt, K-Phite 7 LP, Linebacker WDG, Reliant</b> (fosetyl-Al, phosphorous acid and salts)	See label	See label	See label	See label	See label for specific product guidelines.
	40	<b>Forum</b> (dimethomorph)	6 oz	30 oz	See label	0.5	Limit is 5 appl./crop. Apply with another fungicide that has a different mode of action and alternate.
	40	<b>Orondis Ultra B Revus</b> (mandipropamid)	8 fl oz	32 fl oz	0	4 hr	Phytophthora blight suppression only.
	43	<b>Presidio</b> (fluopicolide)	4 fl oz	8 fl oz/year	2	0.5	Tank-mix with another fungicide product with a different FRAC code.
	45 & 40	<b>Zampro</b> (ametoctradin + dimethomorph)	14 fl oz	42 fl oz/year	0	0.5	No more than 2 sequential applications. Addition of a spreading/penetrating adjuvant is recommended.
	U15	<b>Orondis Gold 200</b> <b>Orondis Opti A</b> <b>Orondis Ultra A</b> (oxathiapiprolin)	See label	See label	0	4 hr	Cucurbit vegetables, crop group 9. See label.
	U15 & 40	<b>Orondis Ultra</b> (oxathiapiprolin + mandipropamid)	8 fl oz	32 fl oz/year	0	4 hr	Make no more than 2 sequential applications before rotating to a fungicide with a different mode of action. Check label for additional details.
	M1	(copper compounds) <b>Many brands available:</b> AmeriCop 40 DF, Badge SC, Badge X2, Basic Copper 53, Camelot-O, Champ DP Dry Prill, Champ Formula 2 FL, Champ WG, ChampION++, Copper-Count-N, CS 2005, Cueva, Cuprofix Ultra 40 Dispers, Cuproxat, ET-F, Kentan DF, Kocide 2000, Kocide 2000-O, Kocide 3000, Kocide 3000-O, Kocide HCU, KOP-5, Kop-Hydroxide, MasterCop, Nordox 75WG, Nu-Cop 50 WP, Nu-Cop 3L, Nu-Cop 30 HB, Nu-Cop 50DF, Nu-Cop HB, Nu-Cop XLR, Previsto	See label	See label	1	Varies from 4 hr to 2 days	Check label for crop-specific guidelines.
	M1 & M2	<b>Top Cop w Sulfur</b> (copper + sulfur)	1 qt	See label	-	1	Do not apply if air temperatures greater than 90°F are expected.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M2	(sulfur) <b>Many brands available:</b> Acoidal, Cosavet DF, Cosavet DF Edge, Crusade DF, Defend DF, Micronized Wettable Sulfur, Micro Sulf, Suffa, Sulfur 6L, Sulfur 90W, Thiolux	See label	See label	See label	1	See individual label. Do not use when temperatures are greater than 90°F or on sulfur-sensitive varieties. Check label for use.
	M2 + 3	<b>Unicorn</b> (sulfur + tebuconazole)	3.75 lb	15 lb	14	0.5	Suppression only. Only available for use on some cucurbits. See label.
M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Praiz, Rialto 720F	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.	

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	P1	<b>Actigard 50WG</b> (acibenzolar-S-methyl)	1 oz	8 oz	0	0.5	Apply preventatively prior disease development. Suppression of disease.
	1	(thiophanate-methyl) <b>Many brands available:</b> 3336 EG, 3336 F, 3336 WP, Cercobin, T-Methyl 4.5 AG, T-Methyl 4.5 F, T-Methyl 70 WP, Incognito 4.5F, Incognito 85WDG, Talaris 70 WSB, Thiophanate methyl 85 WDG, TM 4.5 F, Topsin 4.5 FL, Topsin M WSB	See label	See label	1	1	Follow resistance management guidelines on label.
	3	<b>Rally 40WSP</b> (myclobutanil)	5 oz	See label	0	1	
	3	(tebuconazole) <b>Many brands available:</b> Buzz Ultra DF, Incognito 4.5F, Monsoon, Onset 3.6L, Orius 3.6F, Teb 3.6 SC, Tebu-Crop 3.6F, Tebu 3.6F, Tebuconazole 3.6 F, Tebustar 3.6L, Tebuzol 3.6F, Toledo, Vibe	See label	See label	1	1	Follow resistance management guidelines indicated on the label. Check label for additional guidelines.
	3	<b>Procure 480SC</b> <b>Trionic 4 SC</b> (triflumizole)	8 fl oz	24 fl oz	0	0.5	Follow resistance management guidelines on label.
	3	<b>Proline</b> (prothioconazole)	5.7 fl oz	17.1 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year.
	3	<b>Mettle 125 ME</b> (tetraconazole)	8 fl oz	24 fl oz/year	1	0.5	Make no more than 2 sequential applications before rotating to another fungicide with alternate mode of action.
	3 & M5	<b>Muscle ADV</b> (tebuconazole + chlorothalonil)	2.1 pt	6.4 pt	7	0.5	Watermelon, squash, pumpkin, and melon
	3	<b>Topguard</b> (flutriafol)	14 fl oz	56 fl oz/year	0	0.5	Check label.
	3	<b>Rhyme</b> (flutriafol)	7 fl oz	28 fl oz/year	0	0.5	Do not apply more than 2 sequential applications before rotating with a different FRAC group fungicide.
	3 & 9	<b>Inspire Super</b> (cyprodinil + difenoconazole)	20 fl oz	80 fl oz/year	7	0.5	Do not make more than two consecutive applications. See label for greenhouse use only for cucumber.
	3 & 33	<b>Viathon</b> (tebuconazole + potassium phosphite)	4 pt	12 pt	7	0.5	
	7	<b>Fontelis</b> (penthiopyrad)	16 fl oz	67 fl oz/year	1	0.5	Maximum rate is for year. Do not make more than 2 applications sequentially.
	7	<b>Endura</b> (boscalid)	6.5 oz	26 oz/year	0	0.5	Do not make more than one application before alternating to another labeled fungicide with a different mode of action for at least one application.
	7	<b>Velum Prime</b> (fluopyram)	6.84 fl oz	13.7 fl oz/year	0	0.5	

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 3	<b>Luna Experience</b> (fluopyram + tebuconazole)	17 fl oz	34 fl oz/year	7	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 3 fungicide.
	7 & 3	<b>Propulse</b> (fluopyram + prothioconazole)	13.6 fl oz	34.2 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year. See label for harvesting guidelines.
	7 & 11	<b>Luna Sensation</b> (fluopyram + trifloxystrobin)	7.6 fl oz	27.1 fl oz/year	0	0.5	No more than 2 sequential applications before rotating to a non-FRAC 7, non-FRAC 11 fungicide.
	7 & 11	<b>Pristine</b> (boscalid + pyraclostrobin)	18.5 oz	74 oz/year	0	0.5	Limit is 4 appl./crop and alternate chemistry.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	Do not make more than one application before alternating to a non-FRAC 7, non-FRAC 11 group fungicide. In areas where gummy stem blight resistance to FRAC 7 or 11 fungicides have been reported, tank-mix with chlorothalonil at full rate.
	9 & 12	<b>Alterity 62.5 Switch 62.5WG</b> (cyprodinil + fludioxonil)	14 oz	56 oz/year	1	0.5	Do not make more than two consecutive applications.
	11	<b>Evito 480 SC Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11	<b>Flint Extra Flint</b> (trifloxystrobin)	See label	See label	See label	0.5	(Disease suppression) Limit is 4 appl./crop and alternate chemistry.
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 25C, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxyzone, AZteroid FC, Azteroid FC 3.3, Heritage, Heritage SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions. Heritage and Heritage SC fungicides are only for transplants production. See label for details.
	11	<b>Narvos 50 WDG Sovran</b> (kresoxim-methyl)	4.8 oz	19.2 oz	0	0.5	Follow resistance management guidelines on label.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	11 & M5	<b>Arius ADV Quadris Opti</b> (azoxystrobin + chlorothalonil)	3.2 pt	See label	1	0.5	Limit is 4 appl./crop for all QoI fungicides. Do not make more than one consecutive application. Do not apply to mature watermelons under dry, hot, or other environmental conditions listed on label.
	11 & 3	<b>Quadris Top</b> (azoxystrobin + difenoconazole)	14 fl oz	56 fl oz/year	1	0.5	Must be rotated with a non-QoI-containing fungicides.
	11 & 7	<b>Mural</b> (azoxystrobin + benzovindiflupyr)	See label	See label	0	0.5	Only for vegetable plants grown for resale to consumers.
	12	<b>Emblem</b> (fludioxonil)	7 fl oz	28 fl oz/year	1	0.5	
	7 & 12	<b>Miravis Prime</b> (pydiflometofen + fludioxonil)	11.4 fl oz	22.8 fl oz/year	1	0.5	Do not make more than two applications of Miravis Prime or other Group 7 and 12 before alternation with a fungicide that is not in Group 7 or 12.
	13	<b>Quintec</b> (quinoxifen)	6 fl oz	See label	3	0.5	Do not make more than 4 applications. Do not make more than 2 consecutive applications. Not labeled for all cucurbits; labeled for various melons, cantaloupe, winter squash, gourds, pumpkins, and watermelon.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	19	<b>Affirm WDG Oso 5% SC</b> (polyoxin D zinc salt)	13 fl oz	See label	0	4 hr	
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	3 & 33	<b>Viathon</b> (tebuconazole + potassium phosphite)	3 pt	12 pt	7	0.5	
	U6	<b>Torino</b> (cyflufenamid)	3.4 oz	6.8 oz/year	0	4 hr	
	U8	<b>Vivando</b> (metrafenone)	15.4 fl oz	46.2 fl oz/year	0	0.5	See label for instructions.
	50	<b>Prolio 300 SC</b> (pyriofenone)	5 fl oz	16 fl oz/year	0		Do not make more than 2 sequential applications before rotating to another fungicide with a different mode of action.
<b>Pythium spp.</b>	4	<b>Apron XL, Orondis Gold B, Ridomil Gold SL, Ultra Flourish</b> (mefenoxam)	See label	See label	See label	See label	
	4	(metalaxyl) <b>Many brands available:</b> Acquire, Allegiance FL, Metalaxyl 318 FS, Metalaxyl 265 ST, Metalaxyl 4.0 ST, Metastar 2E, Metastar 4S, MetXL, Sebring 318 FS, Sebring 480 FS	See label	See label	See label	See label	
	4 & 11	<b>Uniform</b> (mefenoxam & azoxystrobin)	See label	See label	See label	0	Apply 0.34 fl oz/1000 ft of row as in-furrow spray in a minimum of 5 gallons of water per acre at planting. Make only one application per season.
	28	<b>Previcur Flex Promess</b> (propamocarb hydrochloride)	1.2 pt	6 pt	2	0.5	Use a tank-mix partner. See label for directions using a contact fungicide and Pythium suppression.
<b>Target spot</b>	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, EQUUS 720 SST, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Orondis Opti B, Praiz	See label	See label	See label	0	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	1	(thiophanate-methyl) <b>Many brands available:</b> 3336 EG, 3336 F, 3336 WP, Cercobin, T-Methyl 4.5 AG, T-Methyl 4.5 F, T-Methyl 70 WP, Incognito 4.5F, Incognito 85WDG, Talaris 70WSB, Thiophanate methyl 85 WDG, TM 4.5 F, Topsin 4.5 FL, Topsin M WSB	See label	See label	1	1	Follow resistance management guidelines on label.
	3	<b>Mettle 125 ME</b> (tetraconazole)	8 fl oz	24 fl oz/year	1	0.5	Make no more than 2 sequential applications before rotating to another fungicide with alternate mode of action.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	3	<b>Topguard</b> (flutriafol)	14 fl oz	56 fl oz/year	0	0.5	Check label.
	3	<b>Rhyme</b> (flutriafol)	7 fl oz	28 fl oz/year	0	0.5	Do not apply more than 2 sequential applications before rotating with a different FRAC group fungicide.
	11	<b>Evito 480 SC</b> <b>Tepera</b> (fluoxastrobin)	See label	See label	See label	See label	
	11	<b>Cabrio EG</b> (pyraclostrobin)	16 oz	64 oz	0	0.5	See label for tank-mixing restrictions.
	11	(azoxystrobin) <b>Many brands available:</b> A-Zox 25SC, Acadia 25C, Aframe, Azoxy 25C, Azoxy 25C Prime, AzoxyStar, Azoxystar, AZteroid FC, Azteroid FC 3.3, Equation, Equation SC, Mazolin, Quadris, Satori, Tetraban, Trevo	See label	See label	1	4 hr	Limit is 4 appl./crop and alternate chemistry. See label for tank-mixing restrictions.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 11	<b>Merivon</b> (fluxapyroxad + pyraclostrobin)	5.5 fl oz	16.5 fl oz/year	0	0.5	See label.
	11 & 3	<b>Topguard EQ</b> (azoxystrobin + flutriafol)	8 fl oz	See label	1	0.5	See label for crop-specific use.
	19	<b>Affirm WDG</b> <b>Oso 5% SC</b> <b>PH-D</b> (polyoxin D zinc salt)	See label	See label	0	4 hr	
	22 & M5	<b>Zing</b> (zoxamide + chlorothalonil)	36 fl oz	See label	0	0.5	
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
<b>Scab</b>	M1	<b>Cueva</b> (copper octanoate)	See label	See label	0	4	See label for crops listed.
	M3	(mancozeb) <b>Many brands available:</b> Dithane DF Rainshield, Dithane F45 Rainshield, Dithane M45, Fortuna 75WDG, Koverall, Manzate Max, Manzate Pro-Stick, Penncozeb 75DF, Penncozeb 80WP, Protect DF, Roper DF Rainshield	See label	See label	5	1	Check label for crop-specific guidelines.
	M1 & M3	<b>Mankocide</b> (copper hydroxide + mancozeb)	3 lb	24 lb	5	2	Some cantaloupe varieties are sensitive. Check label for details.
	M5	(chlorothalonil) <b>Many brands available:</b> Bravo Ultrex, Bravo Weather Stik, Bravo ZN, Chlorothalonil 720, Chlorothalonil 720 SC, Echo 720, Echo 90 DF, Echo ZN, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN, Oranil 6L, Orondis Opti B, Rialto 720 F, Praiz	See label	See label	0	1	Check label for crop-specific guidelines. Follow label recommendations on watermelon after fruit set. Do not apply to mature watermelons under dry, hot, or other environmental conditions indicated in the label.
	M5 & U15	<b>Orondis Opti</b> (chlorothalonil + oxathiapiprolin)	2.5 pt	10 pt/year	0	0.5	Check label for additional guidelines. Make no more than 2 sequential applications before rotating to a fungicide with different mode of action.
	M5 & 27	<b>Ariston</b> (chlorothalonil + cymoxanil)	3 pt	17.5 pt/year	3	0.5	
	4 & M5	<b>Ridomil Gold Bravo SC</b> (mefenoxam + chlorothalonil)	3.25 pt	See label	0	2	Limit is 4 appl./crop.

Pertinent Diseases or Pathogens	Fungicide Group <sup>1</sup>	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks <sup>2</sup>
			Appl.	Season	Harvest	Reentry	
	M3 & 11	<b>Dexter Max</b> (mancozeb + azoxystrobin)	3.2 lb	12.8 lb/year	5	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11.
	M3, 11 & 3	<b>Dexter Xcel</b> (mancozeb + azoxystrobin + tebuconazole)	72 fl oz	288 fl oz/year	7	1	See guidelines of adjuvants and pesticides to be avoided in tank mixing. Do not apply more than one application of Dexter Max or other group 11 fungicides before alternation with fungicides not in group 11. Some cantaloupe varieties are sensitive. Check label.
	P1	<b>Actigard 50WG</b> (acibenzolar-S-methyl)	1 oz	8 oz	0	0.5	Apply preventatively prior to disease development. Suppression of disease.
	7 & 3	<b>Aprovia Top</b> (benzovindiflupyr + difenoconazole)	13.5 fl oz	53.6 fl oz/year	0	0.5	See label for guidelines.
	7 & 12	<b>Miravis Prime</b> (pydiflumetofen + fludioxonil)	11.4 fl oz	22.8 fl oz/year	1	0.5	Do not make more than two applications of Miravis Prime or other Group 7 and 12 before alternation with a fungicide that is not in Group 7 or 12.
	19	<b>Affirm WDG Oso 5% SC PH-D</b> (polyoxin D zinc salt)	See label	See label	0	4 hr	
<b>Fusarium wilt (watermelon)</b>	3	<b>Proline</b> (prothioconazole)	5.7 fl oz	17.1 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year.
	7 & 3	<b>Propulse</b> (fluopyram + prothioconazole)	13.6 fl oz	34.2 fl oz/year	7	0.5	Limit is 1 soil application and 2 foliar applications per year. See label for harvesting guidelines.

<sup>1</sup> FRAC code (fungicide group): Number (1 through 50) and letter U is used to distinguish the fungicide mode of action groups. All fungicides within the same group (with the same number or letter) indicate the same active ingredient or similar mode of action. This information must be considered for the fungicide resistance management decisions. U = unknown, or a mode of action that has not been classified yet. Source: FRAC Code List 2020: <http://www.frac.info/> FRAC = Fungicide Resistance Action Committee.

<sup>2</sup> Information provided in this table applies only to Florida. Be sure to read a current product label before applying any chemical. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned.

Table 7.12. Nonfumigant nematicides for cucurbit crops in Florida.

Product	Application Directions
Mocap 15G (a.i. ethoprop) (only labeled on cucumber)	Apply 13 pounds per acre (7-foot row spacing) or 2.1 pounds per 1,000 linear feet of row in a band 12 to 15 inches wide on the row at planting. Prepare bed and knock off top to provide a level area for treatment. Mix with the top 2 to 4 inches of soil with mechanical equipment right after application. DO NOT use as a seed furrow treatment or allow granules to contact the seed. Only 1 application per season.
Vydate L (a.i. oxamyl)	Apply ½ to 1 gal/acre at planting, preferably via drip application. Make additional applications on a 10-to-14-day interval. Do not apply more than 2 gallons per acre per season. Minimum retreatment interval is 7 days unless a longer interval is stated. If an at-plant application of more than ½ gal/acre is made, do not make more than 2 additional applications.
Nimitz (a.i. fluensulfone)	All applications must be incorporated either physically or via drip or overhead irrigation. Make preplant applications at a rate of 3.5 to 7 pints (56.0 to 80.0 fl oz) per acre, a minimum of seven days before planting. Do not plant any unlisted crops into treated land for 365 days after application of the product. Do not apply more than one application per crop and no more than 112 fl oz of product per acre per year (365 days). Provides control only for nematodes. Growers applying Nimitz must consult the product label to observe the plant-back (recropping) intervals for a variety of leafy vegetables and brassica crops, onions, bananas, sugarcane, and other crops.
Velum (a.i. fluopyram)	Apply max 6.84 fl oz/acre using only chemigation into root zone through low-pressure drip, trickle, microsprinkler, or equivalent equipment. Observe minimum 5-day interval between soil applications. Do not apply more than 13.7 fl oz of Velum (0.446 lb fluopyram) per acre per year, regardless of formulation (Velum and/or Luna) or method of application (soil or foliar). For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after Velum should be a product from a different FRAC group.
Mocap and Vydate are insecticide/nematicides; Velum is a fungicide/nematicide; Nimitz is a true nematicide. Unlike fumigants, these products are not volatile and will move through the soil via water; depending on the water solubility, these products will have different recommendations as far as how to best apply them (see specific label recommendations); when nematode pressure is high, they may not be as consistently effective against root nematodes as the fumigants.	

Table 7.13. Fumigant nematicides for cucurbit crops in Florida.

Nematicide	Broadcast Application <sup>1</sup>		In-the-Row Applications
	Gallons or lb per Acre	fl oz/1000 ft/ chisel-spaced 12" apart	
Telone II <sup>2,3</sup>	9 to 12 gal	26 to 35	For any row spacing, application rates given may be concentrated in the row but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-17 <sup>2,3</sup>	10.8 to 17.1 gal	31.8 to 50.2	For any row spacing, application rates given may be concentrated in the row but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-35 <sup>2,3</sup>	13 to 20.5 gal	38 to 60	For any row spacing, application rates given may be concentrated in the row but shall never exceed labeled maximum for broadcast applications. Consult the product label for additional detail.
Pic-Clor 60	19 to 31.5 gal	57 to 90	For any row spacing, application rates should never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Vapam HL	75 gal	-	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration, and flow-modifying directions and procedures.
KPam HL	60 gal	-	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration, and flow-modifying directions.
Dimethyl Disulfide (DMDS)	51.3 gal	-	Compared to broadcast application, apply proportionately less for in-the-row applications based on the ratio of bed width to row spacing. Consult the product label for additional detail and rate-modifying recommendation.
Allyl Isothiocyanate (AITC) Dominus	40 gal	-	For drip or in-row fumigation and crop termination, consult product label for overall rates, drip concentration, and other flow-modifying directions.

<sup>1</sup> Gallons/acre and fl oz/1000 feet provided only for mineral soils. Higher rates may be possible for heavier-textured (loam, silt, clay) or highly organic soils.

<sup>2</sup> All of the fumigants mentioned are for retail sale and use only by state-certified applicators or persons under their direct supervision. New supplemental labeling for the Telone products must be in the hands of the user at the time of application. See new label details for additional use restrictions based on soil characteristics, buffer zones, requirements for Fumigant Management Plans (FMP) and personal protective equipment (PPE), mandatory good agricultural practices (GAPs), product and applicator training certification, and other rate-modifying recommendations with use of highly retentive mulch films.

<sup>3</sup> Higher application rates are possible in the presence of cyst-forming nematodes.

Rates are believed to be correct for products named, and similar products of other brand names, when applied to mineral soils. Higher rates are required for muck (organic) soils. However, the grower has the final responsibility to see that each product is used legally; read the label of the product to be sure that you are using it properly.