

Facts about Fats and Oils¹

Tiffany N. Stodtko and Wendy J. Dahl²

Fats and oils are important for good health. Although fats and oils are not considered a food group, it is recommended that we consume them in small amounts. Fats provide your body with energy and essential fatty acids and enable your body to absorb fat-soluble vitamins (A, D, E, K). Oils are needed in the diet in small amounts because they are a major source of vitamin E, which has antioxidant properties. Linolenic acid, an omega-3 fatty acid, and linoleic acid, an omega-6 fatty acid, are essential for health.

Types of Fats

There are many different types of dietary fats and oils. These include saturated fats, monounsaturated fats, polyunsaturated fats, and trans fats. All fats and oils contain a mixture of saturated and unsaturated fats, but in different amounts. In general, solid fats contain a greater proportion of saturated fats than do liquid oils.

Saturated fats are found naturally in foods such as beef, lamb, pork, and cheese. Lard (pork fat), butter, and cream also contain higher amounts of saturated fat. The current Dietary Guidelines for Americans includes a recommendation for limiting intake of saturated fats to less than 10% of your total energy intake (USDA 2015) to decrease the risk of cardiovascular disease (e.g., a heart attack). This should be achieved by replacing saturated fats with polyunsaturated and monounsaturated fats. However, the association of saturated fat with heart disease is controversial because the type of saturated fat is also an important consideration (Chowdhury et al. 2014; Dawczynski et al. 2015; Praagman

et al. 2016). Saturated fat from dairy sources may not increase the risk of heart disease (Praagman et al. 2016). A recent study also suggests that total saturated fat intake may not be associated with an increased risk of death from cardiovascular disease, coronary heart disease, or stroke (de Souza et al. 2015).

Monounsaturated fats are found in high levels in olive oil and in oils from high-oleic-acid varieties of canola, soybean, sunflower, and safflower. Beef tallow, lard, peanut oil, and palm oil contain moderate levels of monounsaturated fats. Soybean, canola, and corn oils as well as cocoa butter also contain monounsaturated fats. Although higher intakes of monounsaturated fats are encouraged for heart health, recent evidence suggests that the source of the monounsaturated fat is also important (Schwingshackl and Hoffmann 2014). Olive oil may provide a greater benefit because a higher intake of olive oil is associated with a reduced risk of cardiovascular events (e.g., a heart attack) and stroke (Schwingshackl and Hoffmann 2014).

Polyunsaturated fats are found in higher amounts in safflower, soybean, sunflower, corn, and cottonseed oils. Walnuts, sunflower seeds, and fish are other good sources of polyunsaturated fats. Polyunsaturated fats include omega-3 fatty acids and omega-6 fatty acids. Fatty fish such as salmon, tuna, and sardines are good sources of long-chain, omega-3 fatty acids. Although it is recommended that you consume polyunsaturated fats instead of saturated fats (USDA 2015), most vegetable oils may not provide a strong heart health benefit (Chowdhury et al. 2014). However,

1. This document is FSHN16-3, one of a series of the Food Science and Human Nutrition Department, UF/IFAS Extension. Original publication date June 2016. Revised February 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Tiffany N. Stodtko, former graduate student; and Wendy J. Dahl, associate professor; Food Science and Human Nutrition Department, UF/IFAS Extension, Gainesville, FL 32611.

eating fatty fish, which provide long-chain, omega-3 fatty acids, reduces the risk of heart disease (Kromhout 2012) and stroke (Chowdhury et al. 2012).

Trans fats are found naturally in small amounts within certain animal fats. Most trans fat in our diet has been from industry-produced, partially hydrogenated vegetable oils. Hydrogenation is the process of converting a vegetable oil into a more saturated, more solid fat. Since the 1950s, trans fats were widely used in the manufacturing of baked goods such as biscuits, doughnuts, pastries, cakes, cookies, and pie crusts. Fried foods, snack foods like popcorn and crackers, and stick margarines and vegetable shortenings have also contained trans fats. Trans fats raise blood cholesterol levels and increase the risk of heart disease (Chowdhury et al. 2014; Mozaffarian, Aro, and Willett 2009). There is no safe level of intake of trans fat. Partially hydrogenated oils containing trans fats are no longer Generally Recognized as Safe (GRAS) and will soon be eliminated from the food supply (FDA 2018). Naturally occurring trans fats found in low levels in dairy products do not seem to raise cholesterol to the same levels as the trans fats in partially hydrogenated oils (Gayet-Boyer et al. 2014).

Animal and Plant Sources of Fat

Fats and oils can be categorized by whether they come from an animal or a plant (vegetable) source. Animal fats, such as butter and lard, are primary sources of saturated fats and tend to be solid at room temperature. Fish oils are an exception as they are liquid at room temperature. Vegetable fats and oils usually contain more unsaturated fats than do animal fats. They tend to be liquid at room temperature unless they have been hydrogenated. As hydrogenated fats are derived from vegetable oils, these fats are also grouped under plant sources. Examples include vegetable shortenings and stick margarines.

Daily Recommended Amount

The USDA recommends that you consume small amounts of oils, about 5 teaspoons per day for most adults (USDA 2015). The USDA MyPlate website provides a table of recommended oil allowances and how much oil and fat is present in common foods (USDA n.d.). See <http://www.choosemyplate.gov/oils>. Although essential for health, oils are a rich source of energy. Solid fats and oils contain about 40 calories per teaspoon, making it important to limit the amount consumed to avoid unwanted weight gain.

Cooking with Oils

Oils are often used in cooking. Some oils are more sensitive to high temperatures than others. The temperature at which an oil begins to break down is called the smoke point. Cooking at a temperature that is higher than the smoke point of the oil can result in undesirable flavors caused by breakdown products (e.g., aldehydes, ketones). Peanut and sesame oils have high smoke points and therefore are good choices for frying (Bockisch 1998). Flaxseed and walnut oil have low smoke points and are better suited for foods that do not require heating such as salad dressings (Bockisch 1998). The exposure of oils to high temperatures, and the reuse of frying oils in particular, produce substances that may have adverse effects on health (Dobarganes and Marquez-Ruiz 2015).

Unrefined oils have been purified to a lesser degree than refined oils. Unrefined oils have a more noticeable color, aroma, and flavor. These oils also have a shorter shelf life than refined oils. In general, unrefined oils are suited for foods such as salad dressings or for lower-temperature cooking, whereas most common refined oils are suited for higher-temperature cooking or frying (Bockisch 1998).

In general, solid fats are more stable than oils due to their higher saturated fat content and lower polyunsaturated fat content. Saturated fats are less likely to become rancid and produce undesirable odors and flavors. Storing oils properly is important because oils may become rancid. Oils should be stored in a cool, dark place. Refrigeration after opening is recommended. Make sure to regularly check your oil and discard it if it has developed an “off” odor, usually described as “painty” or “fishy”.

Types of Oils

Canola oil is derived from the seeds of the canola plant. Canola oil is considered a healthy oil due to its low content of saturated fat and its high content of monounsaturated fat (oleic acid). Canola oil has a mild, neutral flavor and higher smoke point, making it a flexible option for both cooking and baking. It can be used to sauté and stir-fry foods. Canola oil with a very high level of monounsaturated fat (oleic acid) is available to the food industry for use as a frying oil.

Coconut oil is extracted from mature coconuts. Although known as an oil, at room temperature, it is a solid fat due to its very high saturated fat content. It has a sweet, nutty flavor and is used in stews, curry and fish dishes, and baked goods. It can be used as a substitute for butter for someone

following a vegan diet. Coconut oil has a low to medium smoke point (Bockisch 1998).

Flaxseed oil is extracted from flaxseed (linseed). Flaxseed oil contains a high amount of polyunsaturated fat, including linolenic acid, an essential fatty acid (Mridula, Barnwal, and Singh 2015). Flaxseed oil is not suitable for frying due to its low smoke point and its lack of oxidative and thermal stability (Boskisch 1998). Using flaxseed oil in salad dressings or for drizzling over pasta or rice dishes is recommended.

Olive oil is extracted from ripe olives by pressing. Most olive oil is sold in virgin (unrefined) form. Refined olive oil is used mainly as a food ingredient, e.g., margarines. Extra virgin olive oil is the highest quality olive oil. Olive oil is high in monounsaturated fat (oleic acid) and may help to reduce the risk of heart disease and stroke (Schwingshackl and Hoffmann 2014). It is a healthier option than butter or margarine, can be used in many foods such as spreads and marinades, and is great as a dip for bread or as a salad dressing. Olive oil has a lower smoke point, so it is best used at lower temperatures (Bockisch 1998).

Peanut oil is derived from peanuts and has a nutty flavor. It is high in monounsaturated fat (oleic acid) and contains vitamin E. Due to its high smoke point, peanut oil is used for deep-frying (Bockisch 1998). It is also used in stir-fries and in many Asian dishes. The majority of peanut oil consumed in the United States has been refined, that is, the protein (which is responsible for the allergic reaction) has been removed (Crevel, Kerkhoff, and Koning 2000).

Safflower oil is extracted from the seed of the safflower plant. Safflower oil is high in polyunsaturated fat (linoleic acid) and is suitable for use as a salad dressing and cooking oil. Safflower oil possessing a very high level of monounsaturated fat (oleic acid) is available to the food industry for use as a frying oil.

Sesame oil is derived from sesame seed. Sesame oil contains both monounsaturated and polyunsaturated fats. It has a strong, nutty flavor and a high smoke point (Bockisch 1998). It can be used in dressings, sauces, stir-fries, and for searing meats. Sesame oil is commonly used in Asian dishes.

Soybean oil is derived from soybean seed and is high in linoleic acid (an omega-6 fatty acid). Soybean oil has a low smoke point (Bockisch 1998), thus it is not recommended for high temperature frying. Soybean oil possessing a very

high level of monounsaturated fat (oleic acid) is available to the food industry for use as a frying oil.

Sunflower oil is made from sunflower seed. Sunflower oil is high in polyunsaturated fat and is suitable for use as a salad dressing and cooking oil. Sunflower oil, which is very high in monounsaturated fat (oleic acid), is available to the food industry for use as a frying oil.

Walnut oil is derived from walnuts by pressing and can be consumed in its unrefined form (Martinez et al. 2010). Walnut oil contains a high amount of linolenic acid, an omega-3 fatty acid. Walnut oil has a deep, nutty flavor, and is best used in salad dressings and for drizzling rather than as a cooking oil due to its low smoke point. Walnut oil becomes rancid quickly, so it needs to be stored in the refrigerator.

In summary, consume oils and fats in moderation. Choosing an oil high in monounsaturated fat, such as olive oil, is a positive step toward health.

References

- Bockisch, Michael. *Fats and Oils Handbook*. Champaign: AOCS Press, 1998.
- Chowdhury, Rajiv, Sarah Stevens, Donal Gorman, An Pan, Samantha Warnakula, Susmita Chowdhury, Heather Ward, Laura Johnson, Francesca Crowe, Frank B. Hu, and Oscar H. Franco. 2012. "Association between fish consumption, long chain omega 3 fatty acids, and risk of cerebrovascular disease: systematic review and meta-analysis." *British Medical Journal* 345:e6698. doi: 10.1136/bmj.e6698.
- Chowdhury, Rajiv, Samantha Warnakula, Setor Kunutsor, Francesca Crowe, Heather A. Ward, Laura Johnson, Oscar H. Franco, Adam S. Butterworth, Nita G. Forouhi, Simon G. Thompson, Kay T. Khaw, Dariush Mozaffarian, John Danesh, and Emanuele Di Angelantonio. 2014. "Association of dietary, circulating, and supplement fatty acids with coronary risk: a systematic review and meta-analysis." *Annals of Internal Medicine* 160(6):398–406. doi: 10.7326/m13-1788.
- Crevel, René, Mía. A. Kerkhoff, and M. M. Koning. 2000. "Allergenicity of refined vegetable oils." *Food and Chemical Toxicology* 38(4):385–93.
- Dawczynski, Christine, Marcus E. Kleber, W. Marz, Gerhard Jahreis, and Stefan Lorkowski. 2015. "Saturated fatty acids are not off the hook." *Nutrition, Metabolism &*

Cardiovascular Diseases 25(12):1071–8. doi: 10.1016/j.numecd.2015.09.010.

De Souza, Russell J., Andrew Mente, A. Maroleanu, Adriana I. Cozma, Vanessa Ha, Teruko Kishibe, Elizabeth Uleryk, Patrick Budyłowski, Holger Schunemann, Joseph Beyene, and Sonia Anand. 2015. “Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies.” *British Medical Journal* 351:h3978. doi: 10.1136/bmj.h3978.

Dobarganes, Carmen, and Gloria Marquez-Ruiz. 2015. “Possible adverse effects of frying with vegetable oils.” *British Journal of Nutrition* 113 Suppl 2:S49–57. doi: 10.1017/s0007114514002347.

Gayet-Boyer, Constance, Fanny Tenenhaus-Aziza, Caroline Prunet, Corinne Marmonier, Corinne Malpuech-Brugere, Benoit Lamarche, and Jean Michael Chardigny. 2014. “Is there a linear relationship between the dose of ruminant trans-fatty acids and cardiovascular risk markers in healthy subjects: results from a systematic review and meta-regression of randomised clinical trials.” *British Journal of Nutrition* 112(12):1914–22. doi: 10.1017/s0007114514002578.

Kromhout, Daan. 2012. “Omega-3 fatty acids and coronary heart disease. The final verdict?” *Current Opinion in Lipidology* 23(6):554–9. doi: 10.1097/MOL.0b013e328359515f.

Martinez, Marcela L., Diana O. Labuckas, Alicia L. Lamarque, and Damián M. Maestri. 2010. “Walnut (*Juglans regia* L.): genetic resources, chemistry, by-products.” *Journal of the Science of Food and Agriculture* 90(12):1959–67. doi: 10.1002/jsfa.4059.

Mozaffarian, D., A. Aro, and Walter C. Willett. 2009. “Health effects of trans-fatty acids: experimental and observational evidence.” *European Journal of Clinical Nutrition* 63 Suppl 2:S5–21. doi: 10.1038/sj.ejcn.1602973.

Mridula, D., Pradyuman Barnwal, and Kaushalendra Singh. 2015. “Screw pressing performance of whole and dehulled flaxseed and some physico-chemical characteristics of flaxseed oil.” *Journal of Food Science and Technology* 52(3):1498–506. doi: 10.1007/s13197-013-1132-6.

Praagman, Jaïke, Joline W. J. Beulens, Marjan Alssema, Peter L. Zock, Anne J. Wanders, Ivonne Sluijs, Yvonne T. van der Schouw. 2016. “The association between dietary saturated fatty acids and ischemic heart disease depends on the type and source of fatty acid in the European

Prospective Investigation into Cancer and Nutrition-Netherlands cohort.” *The American Journal of Clinical Nutrition* 103(2):356–65. doi: 10.3945/ajcn.115.122671.

Schwingshackl, Lukas, and Georg Hoffmann. 2014. “Mono-unsaturated fatty acids, olive oil and health status: a systematic review and meta-analysis of cohort studies.” *Lipids in Health and Disease* 13:154. doi: 10.1186/1476-511x-13-154.

United States Department of Agriculture (USDA). n.d. “How Much is My Allowance for Oils?” Accessed February 11, 2020. <http://www.choosemyplate.gov/oils>

United States Department of Agriculture (USDA). 2015. “Dietary Guidelines for Americans 2015-2020.” Accessed February 11, 2020. <http://health.gov/dietaryguidelines/2015/guidelines/>

United States Food and Drug Administration (FDA). 2018. “Final Determination Regarding Partially Hydrogenated Oils (Removing Trans Fat).” Accessed February 11, 2020 <https://www.fda.gov/food/food-additives-petitions/final-determination-regarding-partially-hydrogenated-oils-removing-trans-fat>