

Rainfast Time for Postemergence Herbicides Commonly Used in the Everglades Agricultural Area: Quick Reference Sheet¹

D.C. Odero²

If rainfall occurs soon after herbicide application, it can affect the effectiveness of postemergence herbicides. Certain postemergence herbicides differ in their time requirements for rain-free periods (rainfast) following application. The exact time required to protect herbicide activity and efficacy will vary with weed species, environmental conditions, and other factors, including the type of adjuvant added to the spray mix. It is important to always READ THE LABEL instructions on rainfastness before applying postemergence herbicides during periods when rainfall is frequent. For more information on rainfastness and pesticide use, see the EDIS publication PI238 – *Rainfastness of Pesticides* (<http://edis.ifas.ufl.edu/pi238>).

The table below (Table 1) shows the required time interval between herbicide application and rain-free period for postemergence herbicides commonly used in sugarcane and vegetable production in the Everglades Agricultural Area and the surrounding region.

1. This document is SS-AGR-351, one of a series of the Agronomy Department, UF/IFAS Extension. Original publication date August 2011. Revised September 2018. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. D.C. Odero, associate professor, Agronomy Department, UF/IFAS Everglades Research and Education Center, Belle Glade, FL.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition. Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Table 1. Rainfast time for postemergence herbicides commonly used in the Everglades Agricultural Area.

Herbicide	Common name	Hours until rainfast
Aim	Carfentrazone	1
Armezon	Topramezone	1
Assure II	Quizalofop	1
Asulam (various)	Asulam	48
Asulox	Asulam	48
Atrazine (various)	Atrazine	4
Banvel	Dicamba	4
Basagran	Bentazon	4
Cadet	Fluthiacet	4
Callisto	Mesotrione	1
Clethodim (various)	Clethodim	1
Clincher SF	Cyhalofop	2
Dicamba (various)	Dicamba	4
Envoke	Trifloxysulfuron	3
Evik	Ametryn	4
Fusilide DX	Fluazifop	1
Glyphosate (several)	Glyphosate	1–4
Goal/Goal Tender	Oxyfluorfen	1
Gramoxone Inteon	Paraquat	0.25–0.5
Grasp SC	Penoxsulam	1
Grasp Xtra	Penoxsulam + triclopyr	1
Londax	Bensulfuron	4
Metribuzin (various)	Metribuzin	6
Poast	Sethoxydim	1
Propanil (various)	Propanil	4
Pursuit	Imazethapyr	1
Reflex	Fomesafen	1
Regiment	Bispyribac-sodium	8
Roundup Original Max	Glyphosate	1–2
Roundup PowerMax	Glyphosate	0.5
Roundup WeatherMax	Glyphosate	0.5
Sandea	Halosulfuron	4
Select/Select Max	Clethodim	1
Sharpen	Saflufenacil	1
Touchdown CT2	Glyphosate	0.5–1
Touchdown HiTech	Glyphosate	0.5–1
Touchdown Total	Glyphosate	0.5–1
Ultra Blazer	Acifluorfen	4
Yukon	Halosulfuron + Dicamba	4
2,4-D amine (various)	2,4-D amine	6–8
2,4-D ester (various)	2,4-D ester	1–3