

# 2025–2026 Florida Citrus Production Guide: Melanose<sup>1</sup>

Megan M. Dewdney<sup>2</sup>

Management of melanose, caused by the fungus *Diaporthe citri*, is often necessary in mature groves where fruit is intended for the fresh market, particularly if recently killed twigs and wood are present because of freezes or other causes. Melanose management is not usually needed in groves less than 10 years old or in those where fruit is intended for processing. As a caveat, more dead wood is present in canopies of young trees because of huanglongbing (HLB) than historically was the case. Growers should monitor young groves for dead wood and start applications before melanose becomes economically damaging. Grapefruit is especially susceptible to melanose, but all citrus may be damaged by the disease.

Spores produced by *D. citri* in the asexual stage are the primary inoculum of this fungus. The sexual stage is relatively unimportant for disease management. The asexual spores are produced in flask-shaped structures embedded in dead twigs in the tree canopy, especially those twigs that have died within the last several months. In addition to producing spores on melanose-affected twigs after they die, *D. citri* is able to colonize dead twigs that were originally healthy and can produce inoculum on them as well. Spores are produced during wet periods and dispersed to young, susceptible fruit, leaves, and twigs by rain splash. No spores are produced from symptomatic

leaves, fruit, or living twigs. At 75°F–80°F (23.8°C–26.7°C), 10–12 hours of leaf wetness are required for infection, and at lower temperatures, up to 18–24 hours may be needed. These leaf wetness periods are not uncommon in Florida.

Grapefruit is susceptible to melanose infection from fruit set until it reaches 2.5–3.0 inches (6.4 cm–7.6 cm) in diameter, normally in late June or early July. Fungicides are effective for only short periods when applied to rapidly expanding fruit or leaves. Because April is usually a low-rainfall month and fruit is small and growing rapidly, the first spray for melanose control is not usually applied until mid- to late April. One or two applications are sufficient for control on oranges and most tangerines unless the trees have abundant dead wood, as in a year after a freeze. For fresh-market grapefruit, the first application should be made when the fruit reaches a diameter of 0.25 inch to 0.5 inch (0.64 cm–1.3 cm) (about mid- to late April depending on local conditions). With average-quality copper products, usually about 2 lb/acre of metallic copper are needed for each 3-week period. If using a calendar application schedule, applications should continue until the fruit become resistant. Rates can be reduced if applications are made more frequently or increased if applications are made less often. For melanose control on large trees, no more than 8 lb–12 lb metallic copper are needed per year even if

1. This document is PP-145, one of a series of the Department of Plant Pathology, UF/IFAS Extension. Original publication date December 1999. Revised annually. Most recent revision August 2025. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. © 2025 UF/IFAS. This publication is licensed under [CC BY-NC-ND 4.0](#).

2. Megan M. Dewdney, associate professor and Extension specialist, Department of Plant Pathology, UF/IFAS Citrus Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.

copper is also used for the control of scab, canker, or greasy spot.

Copper residues are reduced with fruit expansion and as a result of rainfall. The [Citrus Copper Application Scheduler](#) is a model that estimates whether residues remaining on fruit are sufficient to control the disease. It is based on fruit growth models, the rate and time of the last application, and rainfall since the last application. It has proven helpful for optimizing the timing of applications for melanose control. Further information about the model is available in EDIS publication [PP289](#), “A Web-Based Tool for Timing Copper Applications in Florida Citrus”.

An early-June application of copper to manage late melanose damage will serve as the first greasy-spot spray. However, applications of copper in hot, dry weather may cause fruit spotting or darken existing blemishes. If copper fungicides are applied from May to September, they should be applied when temperatures are moderate (<94°F) at rates no more than 2 lb of metallic copper per acre, without petroleum oil, and using spray volumes of at least 125 gal/acre.

The strobilurin-containing fungicides Amistar Top, Gem, Headline, Pristine, and Quadris are also effective for melanose control and can be used at any time for disease control. Copper fungicides are more economical and are most important for melanose control. However, because copper fungicides applied in hot weather can damage fruit, use of strobilurin-containing fungicides at that time will avoid phytotoxicity and control greasy spot as well as melanose. Strobilurins appear to have lower residual activity for melanose control than do copper fungicides. Thus, applications may have to be made at shorter intervals, especially when rainfall is high.

Fungi may develop resistance to strobilurin fungicides. These materials (FRAC 11) should never be used more than twice in a row, and no more than two strobilurin-containing applications should be used for melanose control.

## Web Addresses for Links

Citrus Copper Application Scheduler: <http://agroclimate.org/tools/citrus-copper-application-scheduler/>

EDIS publication PP289, “A Web-Based Tool for Timing Copper Applications in Florida Citrus”: <https://doi.org/10.32473/edis-pp289-2012>

## Recommended Chemical Controls

READ THE LABEL.

See Table 1.

Rates for pesticides are given as the maximum amount required to treat mature citrus trees unless otherwise noted. To treat smaller trees with commercial application equipment including handguns, mix the per-acre rate for mature trees in 125 gal of water. Calibrate and arrange nozzles to deliver thorough distribution, and treat as many acres as this volume of spray allows.

Table 1. Recommended chemical controls for melanose.

Pesticide	FRAC MOA <sup>1</sup>	Mature Trees Rate/Acre <sup>2</sup>
copper fungicide	M 01	Use label rate.
Amistar Top <sup>4</sup>	11/3	15.4 fl oz. Do not apply more than 61.5 fl oz (0.2 lb azoxystrobin and 0.13 lb difenoconazole)/acre/season for all uses. Do not apply more than 0.5 lb a.i./acre/season of difenoconazole. Do not apply more than 1.5 lb a.i./acre/season of azoxystrobin. Do not apply more than 3 times. Minimum retreatment interval is 7 days.
Gem 500 SC <sup>3</sup>	11	1.9–3.8 fl oz. Do not apply more than 15.2 fl oz/acre/season for all uses. Do not apply within 7 days of harvest.
Headline SC <sup>3</sup>	11	12–15 fl oz. Do not apply more than 54 fl oz (0.88 lb a.i.)/acre/season for all uses.
Pristine <sup>4</sup>	11/7	16–18.5 oz. Do not apply more than 74 oz/acre/season for all uses, which is equivalent to 1.17 lb a.i./acre/season of boscalid and 0.592 lb a.i./acre/season of pyraclostrobin. Up to 0.88 lb a.i./acre/season of pyraclostrobin can be used. Minimum retreatment interval is 10 days.
Quadris <sup>3</sup>	11	12.0–15.5 fl oz. Do not apply more than 90 fl oz (1.5 lb a.i.)/acre/season for all uses. Minimum retreatment interval is 7 days.

<sup>1</sup> Mode of action class for citrus pesticides from the Fungicide Resistance Action Committee (FRAC) 2024. Refer to chapter 4, “[Pesticide Resistance and Resistance Management](#)”, for more details.

<sup>2</sup> Lower rates can be used on smaller trees. Do not use less than the minimum label rate.

<sup>3</sup> Do not use more than 4 applications of strobilurin-containing fungicides/season. Do not make more than 2 sequential applications of strobilurin-containing fungicides.

<sup>4</sup> Do not make more than 4 applications of Pristine or Amistar Top/season. Do not make more than 2 sequential applications of Pristine or Amistar Top before alternating to a non-strobilurin, non-SDHI, non-DMI fungicide.