

Chapter 12. Onion, Leek, and Chive Production¹

Johan Desaegeer, Peter J. Dittmar, Pamela D. Roberts, Xavier Martini, and Lincoln Zotarelli²

Botany and Planting

Onion—*Allium cepa*, Alliaceae (Amaryillidaceae)

Chive—*Allium schoenoprasum*

Bunching onion—*Allium fistulosum*

Leek—*Allium ampeloprasum*

Table 1. Planting dates for onion and alliums.

Planting Region	Dates	
	Seeded	Transplanted
North Florida	mid-Sept–mid-Nov	Nov–Jan
Central Florida	Oct	Dec–Jan
South Florida	Oct	Dec–Jan

Table 2. Planting information for onion and alliums.

Planting Parameters	Measurements	
	Seeded	Transplanted
Distance between rows (in)	14–18	14–18
Distance between plants (in)	3–4	4–6
Seeding depth (in)	0.25–0.5	–
Seed per acre (lb)	3–4	1
Days to maturity	100–130	100–130
Plant populations (acre)	149,343	112,123

1. This document is HS730, one of a series of the Department of Horticultural Sciences, UF/IFAS Extension. Original publication date June 2015. Revised annually. Most recent revision August 2025. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. © 2025 UF/IFAS. This publication is licensed under CC BY-NC-ND 4.0.

2. Johan Desaegeer, associate professor, nematology, Department of Entomology and Nematology, UF/IFAS Gulf Coast Research and Education Center; Peter J. Dittmar, associate professor, Department of Horticultural Sciences; Pamela D. Roberts, professor, Department of Plant Pathology, UF/IFAS Southwest Florida Research and Education Center; Xavier Martini, associate professor, insect behavior, Department of Entomology and Nematology, UF/IFAS North Florida Research and Education Center; Lincoln Zotarelli, professor, Department of Horticultural Sciences; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

Use pesticides safely. Read and follow directions on the manufacturer’s label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county’s UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.

Cultivars

Table 3. Onion and leek cultivars.

Crop	Cultivars
Onion	Dawn (H), Granex 33 (H), Pirate (H), Plethora (H), Prowler (H), Red Hunter, Savannah Sweet (H), Sweet Caroline (H), Sweet Harvest (H), Sweet Success (H), Vidurra (H)
Bunching Onion	Perfecto Blanco, Tokyo Long White, White Portugal
Leeks	Chives, Jolant, King Richard, Lancelot, Megathon, Staro, Tivi, Tornado, Verina
H=hybrid	

The following tables list registered pesticides that should be integrated with other pest management methods. Additional information on integrated management methods can be requested from UF/IFAS Extension horticulture or agriculture Extension agents. A list of local UF/IFAS Extension offices is available at <https://sfyl.ifas.ufl.edu/find-your-local-office/>.

Table 4. Herbicides approved for managing weeds in onion, garlic, and shallot. **Labels change frequently. Be sure to read a current product label before applying any chemical.** Contact: Peter J. Dittmar, UF/IFAS Department of Horticultural Sciences, Gainesville.

	Active Ingredient lb a.i./Acre	Trade Name Product/Acre	MOA Code	Crops	Weeds Controlled/Remarks
PREEMERGENCE	Bensulide 5.0–6.0	(Prefar) 4 E 5–6 qt	8	Dry bulb vegetables: onion, garlic, shallot	Annual broadleaf and grass control. Incorporate or irrigate 1–2 inches deep within 36 hr of application. Consult label for rotational restrictions.
	Carfentrazone up to 0.031	(Aim) 2 EC up to 2 fl oz	14	Onion, garlic, leek, chive, shallot	Apply as a preplant burndown for emerged broadleaf weeds. Use crop oil concentrate, methylated seed oil, or nonionic surfactant at recommended rates. No more than 0.096 lb a.i./A per season. No pretransplant interval.
	Flumioxazin up to 0.06	(Chateau) EZ up to 2 fl oz	14	Garlic, onion (dry bulb)	Broadleaf control. Apply to transplanted onions between the 2- and 6-leaf stage. Apply to direct seeded onions between the 3- and 6-leaf stage.
	Glyphosate	(various formulations) consult label	9	Onion, garlic, leek, chive, shallot	Control of emerged broadleaf and grass weeds. Consult individual labels for restrictions.
	Oxyfluorfen 2.0	(Goal 2XL) 2 EC 2.0 pt (GoalTender) 4 EC 1.0 pt	14	Onion (dry bulb), garlic (dry bulb)	Certain broadleaf weeds. Transplanted only. Apply within 2 days of transplanting. Necrotic lesions, twisting, or stunting of plants can occur if applications are made during cool, wet weather or prior to the full development of the true leaves.
	Paraquat 0.5–1.0	(Gramoxone) 2 SL 2–4 pt	22	Onion (dry bulb)	Controls emerged weeds. Apply prior, during, or after planting, but before crop emergence. No more than 2 applications per season. Use a nonionic surfactant.
	Paraquat 0.63–1.0	(Gramoxone) 2 SL 2.5–4.0 pt (Firestorm) 3 SL 1.7–2.7 pt	22	Onion (seeded), garlic	Emerged broadleaf and grass weeds. Apply as a preplant burndown. No more than 1 lb a.i./A per season.
	Pelargonic acid	(Scythe) 4.2 EC 3%–10% v/v	27	Onion, garlic, leek, chive, shallot	Emerged broadleaf and grass weeds. Apply before emergence of crop. Product is a contact, nonselective, foliar-applied herbicide. There is no residual activity.

	Active Ingredient lb a.i./Acre	Trade Name Product/Acre	MOA Code	Crops	Weeds Controlled/Remarks
	Pendimethalin Mineral 0.5–0.75 Muck 1.0–2.0	(Prowl) 3.3 EC Mineral: 1.2–1.8 pt Muck: 2.4–4.8 pt (Prowl H20) 3.8 Mineral: 1.0–1.5 pt Muck: 2.0–4.0 pt	3	Onion (dry bulb), shallot (dry bulb), garlic	Mineral soils: Apply when crop has 2–9 true leaves. Muck soils: May be applied as sequential application: preemergence through loop stage, early postemergence (2–6 true leaf stage), late postemergence (6–9 true leaf stage). No more than 5.9 lb a.i./A per growing season. PHI 45 days.
	Pyraflufen ethyl 0.001–0.003	(ET Herbicide/ Defoliant) 0.5–2.0 fl oz	14	Bulb vegetables	Emerged broadleaf and grass weeds. Apply as a preplant burndown treatment.
POSTEMERGENCE	Bromoxynil 0.38–0.5	(Buctril) 2 EC 1.5–2.0 pt	6	Garlic	Broadleaf weeds. Apply after emergence but before 12 inches tall. Weeds are most susceptible up to the 4-leaf stage or 2 inches height or 1 inch wide. PHI 112 days.
	Carfentrazone up to 0.031	(Aim) 2 EC up to 2 fl oz	14	Onion (dry bulb), garlic, leek, chive, shallot	Emerged broadleaf control. Post-direct hooded application to row middles for burndown of emerged weeds. Use crop oil concentrate or nonionic surfactant at recommended rates. PHI 0 days.
	Clethodim 0.09–0.25 0.07–0.25	(Arrow) 2 EC 6–16 fl oz (Select Max) 1 EC 9–32 fl oz	1	Onion (dry bulb), garlic, shallot (dry bulb)	Annual and perennial grass control. Some labels include green onion. Consult label for use of a crop oil concentrate or nonionic surfactant. PHI 45 days.
	Clethodim 0.09–0.13	(Arrow) 2 EC 6–8 fl oz (Select Max) 1 EC 12–16 fl oz	1	Chive, leek	Emerged annual and perennial grasses. Consult labels rate of COC or NIS. No more than 0.5 lb a.i./A per season. Select Max is not registered in leek. PHI 14 days.
	Dimethenamid-p up to 1.0	(Outlook) 6 EC up to 21 fl oz	15	Onion (dry bulb), garlic, leek, shallot (dry bulb, green)	Broadleaf and grass weeds. Apply after the 2-true-leaf stage. Apply as a single application or split application. Split application of 10–14 fl oz, followed by 7–10 fl oz/A with 14 days between applications. No more than 21 fl oz/A per growing season. PHI 30 days.
	Fluazifop 0.188	(Fusilade DX) 12 fl oz	1	Onion (dry bulb), garlic	Annual and perennial grass weeds. No more than 48 fl oz/A per season. Include a COC or NIS. PHI 45 days.
	Oxyfluorfen 0.5	(Goal 2 XL) 2 EC 0.5 pt direct seeded (GoalTender) 4 EC 0.25 pt direct seeded	14	Onion (dry bulb), garlic (dry bulb)	Certain broadleaf weeds. Direct seeded only. Apply after the crop has 2 true leaves. Multiple treatments can be applied. No more than 0.5 lb a.i./A. Necrotic lesions, twisting, or stunting of plants can occur if applications are made during cool, wet weather or prior to development of true leaves. PHI 60 days.
	Sethoxydim 0.19–0.28	(Poast) 1.5 EC 1.0–1.5 pt	1	Onion (dry bulb, green), garlic, leek, shallot	Grass weeds. Include a surfactant. No more than 4.5 pt/A per season. PHI 30 days.
	Treflan 0.35–0.5	(Treflan) 4 EC 0.75–1.0 pt (Treflan TR-10) 3.75–5.0 lb	3	Onion (dry bulb)	Broadleaf and grass weeds. Apply at layby to the soil surface between onion rows. Remove weeds that are present. Do not apply in muck soils. PHI 60 days.

Table 5. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots. **Labels change frequently. Be sure to read a current product label before applying any chemical. Please refer to chapter 19 for information on biopesticides, including materials labeled for certified organic production.** Contact: Xavier Martini, UF/IFAS North Florida Research and Education Center—Quincy.

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
Aphids	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2.5 pt 8F: 1–1.56 pt	24	3	Leeks, shallots, garlic. No more than 3 applications per year for garlic and 2 for all others.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, and garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 applications per season. OMRI-listed. ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 oz/A per season.
Beet armyworm	1A	*Lannate LV; *SP (methomyl)	LV: 1.5 pt SP: 0.5 lb	48	7	Garlic only.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 applications per season. OMRI-listed. ²
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 fl oz/A per season.
	28	Coragen (chlorantraniliprole)	3.5–5.0 fl oz	4	1	No more than 4 foliar applications or 15.4 fl oz/A per crop. For entire bulb vegetable group.
Caterpillars (includes cutworms and armyworms)	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb and garlic only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	Garlic: 6.4–12.8 oz	12	1	Garlic.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	1.92–3.20 fl oz	12	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season. Onion and garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 applications per season. OMRI-listed. ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 fl oz/A per season.
	5	Entrust SC (spinosad)	3–8 fl oz	4	1	No more than 5 applications per year (29 fl oz). For leeks, garlic, and shallots. OMRI-listed. ²
	5	Radiant SC (spinetoram)	5–10 fl oz	4	1	Use with an adjuvant.
	11A	Agree WG (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	1.0–2.0 lb	4	0	Apply when larvae are small for best control.
	11A	Biobit HP (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5–2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. Can be used in the greenhouse. OMRI-listed. ²
	11A	Crymax WDG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5–2.0 lb	4	0	Use high rate for armyworms. Treat when larvae are young.
	11A	Deliver (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25–1.5 lb	4	0	Use higher rates for armyworms. OMRI-listed. ²
	11A	DiPel DF (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25–2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. OMRI-listed. ²
	11A	Javelin WG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.12–1.5 lb	4	0	Treat when larvae are young. Thorough coverage is essential. OMRI-listed. ²
	11A	Xentari DF (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	0.5–2.0 lb	4	0	Treat when larvae are young. Thorough coverage is essential. May be used in the greenhouse. OMRI-listed. ²
	18	Intrepid 2F (methoxyfenozide)	4–12 fl oz	4	1	Slow-acting IGR (insect growth regulator) that only effects larvae.
Fire ants	6	Clinch (abamectin)	Broadcast application: 1 lb Per ant mound: 5–7 tablespoons	4	0	No more than 4 lb/A per year. Only apply when ants are active and no rain is forecast for the next 6 hr.
	7A	Extinguish (S-methoprene)	1–1.5 lb	4	0	Slow-acting IGR. Best applied early spring and fall where crop will be grown. Colonies will be reduced after 3 weeks and eliminated after 8–10 weeks. May be applied by ground equipment or aerially.
	7C	Esteem Ant Bait (pyriproxyfen)	1.5–2.0 lb	12	1	Dry bulb only.
Flea beetles	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Garlic. (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3–8 fl oz	4	1	No more than 5 applications per year (29 fl oz). For leeks, garlic, and shallots. OMRI-listed. ²

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
Liriomyza Leafminers	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Label is registered only for garlic. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 applications per season. OMRI-listed. ²
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 oz	12	30	No more than 38 fl oz/A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Garlic. (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3–8 fl oz	4	1	No more than 5 applications per year (29 fl oz). For leeks, garlic, and shallots. OMRI-listed. ²
	5	Radiant SC (spinetoram)	5–10 fl oz	4	1	Use with an adjuvant.
	6	*Agri-Mek SC (abamectin)	1.75–3.5 fl oz	12	30	No more than 10.25 fl oz/A per season. Must be used with a nonionic activator-type wetting, spreading, and/or penetrating adjuvant, not a binder-sticker-type adjuvant.
	6 + 28	*Minecto Pro (abamectin + cyantraniliprole)	7–10 fl oz	24	30	No more than 20.0 fl oz/A.
	17	Trigard (cyromazine)	2.66 oz	12	7	No more than 6 applications per crop.
	28	Exirel (cyantraniliprole)	13.5–20.5 fl oz	12	1	No more than 0.4 lb a.i./A of cyantraniliprole-containing products per season. Toxic to bees. Do not allow drift to blooming crops or weeds.
Onion maggot (<i>Delia antiqua</i>)	1B	*Diazinon AG500 *Diazinon 50W (diazinon)	AG500: 2–4 qt 50W: 4–8 lb	72	preplant	No more than 1 soil application per year. For leeks, garlic, and shallots.
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2.5 pt 8F: 1–1.56 pt	Leeks, shallots, garlic: 24	3	Leeks, shallots, and garlic. No more than 3 applications per year for garlic and 2 for all others.
	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Label is registered only for garlic. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, and garlic.

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
	3A	*Pounce 25 WP (permethrin)	Onions: 6.4–19.2 oz Garlic: 6.4–12.8 oz	12	1	Garlic.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	1.92–3.20 fl oz	12	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season. Onion and garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 times per season. OMRI-listed. ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 fl oz/A per season.
Plant bugs	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	2.56–3.84 fl oz	12	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season. Onion and garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 applications per season. OMRI-listed. ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.
Stink bugs	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Garlic only. Label is registered only for garlic. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	Karate (lambda-cyhalothrin)	2.56–3.84 fl oz	24	14	No more than 30.72 fl oz/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	Onions: 6.4–19.2 oz Garlic: 6.4–12.8 oz	12	1	Garlic.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	2.56–3.84 fl oz	12	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season. Onion and garlic.
	3A	Pyganic Crop Protection EC 5.0 (pyrethrins)	4.5–15.61 oz	12	0	No more than 10 application per season. OMRI-listed. ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 fl oz/A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Garlic. (Crop Groups 3-07A and 3-07B)

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
Thrips (check label for species controlled)	1A	*Lannate LV; *SP (methomyl)	LV: 1.5–3.0 pt SP: 0.5–1.0 lb	48	7	–
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2.5 pt 8F: 1–1.56 pt	Leeks, shallots, garlic: 24	3	Leeks, shallots, and garlic. No more than 3 applications per year for garlic and 2 for all others.
	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Garlic only. Label is registered only for garlic. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Garlic. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	Onions: 6.4–19.2 oz Garlic: 6.4–12.8 oz	12	1	Bulb onions and garlic.
	3A	*Proaxis Insecticide (gamma-cyhalothrin)	2.56–3.84 fl oz	12	14	No more than 0.12 lb a.i./A (1.92 pt/A) per season. Onion and garlic.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (garlic). No more than 15.36 fl oz/A per season.
	3A + 6	*Gladiator (zeta-cypermethrin + avermectin B1)	19 fl oz	12	30	No more than 38 fl oz/A per season.
	4A	Admire Pro (imidacloprid)	14.0 fl oz	12	21	No more than 14 fl oz/A per season.
	4A	Assail 30SG (acetamiprid)	5.0–8.0 oz	12	7	No more than 4 applications or 32 oz/A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Garlic. (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3–8 fl oz	4	1	No more than 5 applications per year (29 fl oz). For leeks, garlic, and shallots. OMRI-listed. ²
	5	Radiant SC (spinetoram)	5–10 fl oz	4	1	Use with an adjuvant.
	6	*Agri-Mek SC (abamectin)	1.75–3.5 fl oz	12	30	Must be used with a nonionic activator-type wetting, spreading, and/or penetrating adjuvant, not a binder-sticker-type adjuvant. No more than 10.25 fl oz/A per season.
	6 + 28	*Minecto Pro (abamectin + cyantraniliprole)	10 fl oz	12	30	No more than 20.0 fl oz/A.
	7C	Knack IGR (pyriproxyfen)	8 fl oz	12	3	No more than 2 applications, at least 14 days apart. No activity against adult insects.
	28	Exirel (cyantraniliprole)	13.5–20.5 fl oz	12	1	Label contains pollinator protection warnings.
	28	Verimark (cyantraniliprole)	Drip: 6.75–10 fl oz In-furrow at planting: 10–13.5 fl oz	4	1	–

Insect or Mite Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
	U	Requiem EC (extract of <i>Chenopodium ambrosioides</i>)	1.5–3.0 qt	4	0	Begin as soon as thrips are seen.
Wireworms	1B	*Diazinon AG500 *Diazinon 50W (diazinon)	AG500: 2–4 qt 50W: 4–8 lb	72	preplant	No more than 1 soil application per year. For leeks, garlic, and shallots.

¹Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 11.1 January 2024. Number codes are used to distinguish the main insecticide mode of action groups, with additional letters for certain subgroups within each main group. All insecticides within the same group (with the same number) indicate the same active ingredient or a similar mode of action. This information must be considered for the insecticide resistance management decisions. U = unknown, or a mode of action that has not been classified yet.

²Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned. OMRI-listed: Listed by the Organic Materials Review Institute for use in organic production.

*Restricted use insecticide.

Table 6. Insecticides registered for managing arthropod pests of onions. **Labels change frequently. Be sure to read a current product label before applying any chemical. Please refer to chapter 19 for information on biopesticides, including materials labeled for certified organic production.** Contact: Xavier Martini, UF/IFAS North Florida Research and Education Center—Quincy.

Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
Aphids	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2.5 pt 8F: 1.5 pt	12	3	Onions, bulb and green. No more than 2 applications per year. Suppression only.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. No more than 15.36 fl oz/A per season. Suppression only.
Caterpillars (includes cutworms and armyworms)	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb and garlic only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4–19.2 oz	12	1	Bulb onions. No more than 1 lb a.i./A per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions), not green onions. No more than 15.36 fl oz/A per season.
	5	Entrust SC (spinosad)	3–6 fl oz	4	1	No more than 5 applications per year (29 fl oz/A). For onions, bulb and green. OMRI-listed. ²
	5	Radiant SC (spinetoram)	5–10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz/A per year.
	11A	Agree WG (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	1.0–2.0 lb	4	0	Apply when larvae are small for best control.
	11A	Biobit HP (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5–2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. Can be used in the greenhouse. OMRI-listed. ²
	11A	Crymax WDG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5–2.0 lb	4	0	Use high rate for armyworms. Treat when larvae are young.
	11A	Deliver (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25–1.5 lb	4	0	Use higher rates for armyworms. OMRI-listed. ²
	11A	DiPel DF (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25–2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. OMRI-listed. ²
	11A	Javelin WG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.12–1.5 lb	4	0	Treat when larvae are young. Thorough coverage is essential. OMRI-listed. ²
	11A	Xentari DF (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	0.5–2.0 lb	4	0	Treat when larvae are young. Thorough coverage is essential. May be used in the greenhouse. OMRI-listed. ²
	18	Intrepid 2F (methoxyfenozide)	4–8 fl oz	4	1	Green onion subgroup only. No more than 6 applications per year. No more than 64 fl oz/A per year.
	28	Coragen (chlorantraniliprole)	3.5–7.5 fl oz	4	1	Beet armyworm only. No more than 4 foliar applications or 15.4 fl oz/A per crop.

Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
Fire ants	7A	Extinguish (S-methoprene)	1–1.5 lb	4	0	Slow-acting IGR. Best applied early spring and fall where crop will be grown. Colonies will be reduced after 3 weeks and eliminated after 8–10 weeks. May be applied by ground equipment or aerially.
	7C	Esteem Ant Bait (pyriproxyfen)	1.5–2.0 lb	12	1	Dry bulb only.
Flea beetles	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz	12	Foliar: 1	Bulb and green onions (Crop Groups 3-07A and 3-07B). No more than 10.5 fl oz/A per season.
	5	Entrust SC (spinosad)	3–6 fl oz	4	1	No more than 5 applications per year (29 fl oz). For onions, bulb and green. OMRI-listed. ²
Leafminers	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	Onions: 6.4–19.2 oz	12	1	Bulb onions. No more than 1 lb a.i./A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 5.25–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Bulb and green onions (Crop Groups 3-07A and 3-07B). No more than 10.5 fl oz/A per season.
	5	Entrust SC (spinosad)	3–6 fl oz	4	1	No more than 5 applications per year (29 fl oz/A). For onions, bulb and green. OMRI-listed. ²
	5	Radiant SC (spinetoram)	6–10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz/A per year.
	6	*Agri-Mek SC (abamectin)	1.75–3.5 fl oz	12	30	Must be used with a nonionic activator-type wetting, spreading, and/or penetrating adjuvant, not a binder-sticker-type adjuvant. No more than 10.25 fl oz/A per season.
	17	Trigard (cyromazine)	2.66 oz	12	7	No more than 6 applications per crop.
	28	Exirel (cyantraniliprole)	13.5–20.5 fl oz	12	1	No more than 0.4 lb a.i./A of cyantraniliprole-containing products per season. Toxic to bees. Do not allow drift to blooming crops or weeds.
Onion maggot	1B	*Diazinon AG500 *Diazinon 50W (diazinon)	AG500: 2–4 qt 50W: 4–8 lb	72	preplant	No more than 1 soil application per year. For onions, bulb and green.
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2.5 pt 8F: 1.5 pt	12	3	Onions, bulb and green. No more than 2 applications per year.
	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.28 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.

Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
	3A	*Pounce 25 WP (permethrin)	6.4–19.2 oz	12	1	Bulb onions. No more than 1 lb a.i./A per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. No more than 15.36 fl oz/A per season.
	U	Neemix 4.5 EC (azadirachtin)	4–16 fl oz	12	0	OMRI-listed. ²
Plant bugs	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. No more than 15.36 fl oz/A per season.
Stink bugs	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	2.4–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4–19.2 oz	12	1	Bulb onions. No more than 1 lb a.i./A per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions), not green onions. No more than 15.36 fl oz/A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 3.5–7.0 fl oz	12	1	Bulb and green onions (Crop Groups 3-07A and 3-07B). No more than 10.5 fl oz/A per season.
	U	Aza-Direct (azadirachtin)	1–2 pt, up to 3.5 if needed	4	0	Antifeedant, repellent, IGR. OMRI-listed. ²
Thrips (check label for species controlled)	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5–2 pt 8F: 1.56 pt	12	3	Onions, bulb and green. No more than 2 applications per year.
	3A	*Ambush 25W (permethrin)	6.4–19.2 oz	12	1	Dry bulb only. No more than 2 lb a.i./A per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77–1.54 fl oz	24	14	Bulb onions. No more than 0.77 pt/A per season.
	3A	*Mustang (zeta-cypermethrin)	3.2–4.3 oz	12	7	No more than 21.5 oz/A per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	9.6–19.2 oz	12	1	Bulb onions. No more than 1 lb a.i./A per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96–1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. No more than 15.36 fl oz/A per season.
	4A	Admire Pro (imidacloprid)	14.0 fl oz	12	21	No more than 14 fl oz/A per season.
	4A	Assail 30SG (acetamiprid)	5.0–8.0 oz	12	7	No more than 4 applications or 32 oz/A per season.
	4A	Scorpion 35SL (dinotefuran)	Foliar: 5.25–7.0 fl oz Soil: 8.75–10.5 fl oz	12	Foliar: 1 Soil: N/A	Bulb and green onions (Crop Groups 3-07A and 3-07B). No more than 10.5 fl oz/A per season.
	5	Entrust SC (spinosad)	3–8 fl oz	4	1	No more than 5 applications per year (29 fl oz). For onions, bulb and green. OMRI-listed. ²

Pest	MOA Code ¹	Trade Name (Active Ingredient)	Rate Product/Acre	REI (Hours)	Days to Harvest	Remarks ²
	5	Radiant SC (spinetoram)	6–10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz/A per year.
	6	*Agri-Mek SC (abamectin)	1.75–3.5 fl oz	12	30	Must be used with a nonionic activator-type wetting, spreading, and/or penetrating adjuvant, not a binder-sticker-type adjuvant. No more than 10.25 fl oz/A per season.
	7C	Knack IGR (pyriproxyfen)	8 fl oz	12	3	No more than 2 applications. Reapplications at least 14 days apart. No activity against adult insects. Onion (dry bulb only).
	23	Movento (spirotetramat)	5 fl oz	24	3	No more than 10 fl oz per season.
	28	Exirel (cyantraniliprole)	13.5–20.5 fl oz	12	1	No more than 0.4 lb a.i./A of cyazypyr- or cyantraniliprole-containing products per season. Toxic to bees. Rotate with other effective modes of action. Begin making applications when populations are low. If populations are above threshold, use an effective knockdown product prior to applying Exirel.
	28	Verimark (cyantraniliprole)	Drip chemigation: 6.75–10 fl oz	4	1	No more than 2 drip chemigation applications. Begin drip applications when populations first appear. Rotate with other effective modes of action.
	U	Requiem EC (extract of <i>Chenopodium ambrosioides</i>)	1.5–3.0 qt	4	0	Begin as soon as thrips are seen.
Wireworms	1B	*Diazinon AG500 *Diazinon 50W (diazinon)	AG500: 3–4 qt 50W: 4–8 lb	72	preplant	No more than 1 soil application per year. For onions, bulb and green.

¹Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 11.1 January 2024. Number codes are used to distinguish the main insecticide mode of action groups, with additional letters for certain subgroups within each main group. All insecticides within the same group (with the same number) indicate the same active ingredient or a similar mode of action. This information must be considered for the insecticide resistance management decisions. U = unknown, or a mode of action that has not been classified yet.

²Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned. OMRI-listed: Listed by the Organic Materials Review Institute for use in organic production.

*Restricted use insecticide.

Table 7. Onion fungicides ordered by disease and then FRAC group according to their mode of action. **Labels change frequently. Be sure to read a current product label before applying any chemical. Please refer to chapter 19 for information on biopesticides, including materials labeled for certified organic production.** Contact: Pamela Roberts, UF/IFAS Southwest Florida Research and Education Center.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
Bacterial blight (<i>Xanthomonas axonopodis</i> pv. <i>allii</i>)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Champ DP, Champ F2 FL, Champ WG, Cuprofix Ultra 40D, Cuproxat, Cueva, Kentan DF, Kocide 2000, Kocide 3000, Kop-hydroxide, Master Cop, Nu-Cop 50WP, Nu-Cop 50DF, Nu-Cop HB, Previsto	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0	Varies by product from 4 hr to 2 days	Frequent use of copper fungicides may cause foliar burn.
	M1 + M3	Mankocide (copper hydroxide + mancozeb)	2.25 lb	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	P01	Actigard 50WG (acibenzolar-S-methyl)	1 oz	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days of transplanting. Suppression only.
	27 + 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	See label. Suppression only.
Black mold (<i>Aspergillus niger</i>)	9 + 12	Switch 62.5WG Alterity 62.5WG Palladium (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last application.
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	See label	—	—	See label. Suppression only.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	Suppression only.
	9 + 12	Alerity 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz per year	7	0.5	Suppression only.
Botrytis leaf blight (<i>Botrytis squamosa</i>)	M1 + M3	ManKocide (copper hydroxide + mancozeb)	2.25 lb	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Dry bulb: 7 Green: 14	1	–
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	–
	3	(propiconazole) Many brands available: Amtide Propiconazole 41.8% EC, Bumper 41.8 EC, Bumper ES, Fitness, Propicure, Propimax EC, Propistar EC, Shar-Shield PPZ, Tide Propiconazole EC, Tilt	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0.5	Suppression only.
	3 + 7	Luna Experience (fluopyram + tebuconazole)	12.8 fl oz	25.6 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	4 + M5	Ridomil Gold Bravo SC (mefenoxam + chlorothalonil)	See label	–	Dry bulb: 7 Green: 14/21	2	–
	7	Endura (boscalid)	6.8 oz	41 oz	7	0.5	Alternate applications with a different fungicide group.
	7	Fontelis (penthioopyrad)	24 fl oz	72 fl oz	3	0.5	See label.
	7 + 3	Aprovia Top (difenoconazole + benzovindiflupyr)	10.5 fl oz	54.0 fl oz	7	0.5	Adjuvant recommended. No more than 4 applications per year.
	7 + 9	Luna Tranquility (fluopyram + pyrimethanil)	27 fl oz	54.7 fl oz	7	0.5	No more than 2 applications per year.
	7 + 11	Merivon Xemium (fluxapyroxad + pyraclostrobin)	11 fl oz	33 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	No more than 2 consecutive applications.
	9	Vangard WG Vango WG (cyprodinil)	10 oz	28 oz	7	0.5	No more than 2 consecutive applications.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.
	9 + 12	Switch 62.5WG Alterity 62.5WG Palladium (cyprodinil + fludioxonil)	14 oz	56 oz per year	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last applications.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. No more than 5 applications per crop. Suppression only.
	11	Quadris Aframe Many products: see label (azoxystrobin)	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	4 hr	Alternate applications with a different fungicide group.
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	SEE INDIVIDUAL LABEL	14	0.5	Alternate applications with different fungicide groups. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant.
	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz per year	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
	11 + 3	Quilt Xcel (azoxystrobin + propiconazole)	Dry bulb: 21 fl oz Green: 26 fl oz	56 fl oz	Dry bulb: 14 Green: 0	0.5	No more than 1 application before alternating to a non-group 11 fungicide.
	11 + 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	No more than 2 consecutive applications. No more than 6 applications per crop.
	14	Botran 5F (DCNA dicloran)	2–3.2 lb	See label	14	See label	See label.
	22 + M5	Zing! (zoxamide + chlorothalonil)	30 fl oz	7.5 qt	7	0.5	No more than 8 applications per year.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	–
Botrytis neck rot (<i>Botrytis allii</i>)	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Dry bulb: 7 Green: 14	1	–
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	–
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	–
	7 + 3	Aprovia Top (difenoconazole + benzovindiflupyr)	10.5 fl oz	54.0 fl oz	7	0.5	Adjuvant recommended. No more than 4 applications per year.
	7 + 12	Miravis Prime (pydiflumetofen + fludioxonil)	11.4 fl/oz	34.2 fl oz	7	12 hr	–
	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	No more than 2 consecutive applications.
	9	Vangard WG Vango WG (cyprodinil)	10 oz	28 oz	7	0.5	No more than 2 consecutive applications. Suppression only.
	9 + 12	Switch 62.5WG Alterity 62.5WG Palladium (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last application. Suppression only.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	SEE INDIVIDUAL LABEL	14	0.5	Alternate applications with different fungicide groups. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant. Suppression only.
	11 + 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	No more than 2 consecutive applications. No more than 6 applications per crop.
	14	Botran 5F (DCNA dicloran)	3.5 lb	SEE INDIVIDUAL LABEL	14	SEE INDIVIDUAL LABEL	–
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	–
	49 + M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	No more than 2 sequential applications, and do not exceed more than 33% of all fungicide applications with Orondis.
Cercospora leaf spot (<i>Cercospora duggiae</i>)	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	–
Cladosporium leaf blotch (<i>Cladosporium allii</i>)	3 + 7 + 11	Trivapro (propiconazole + benzovindiflupyr + azoxystrobin)	27.7 fl oz	55.4 fl oz per year	Dry bulb: 7 Green: 14	0.5	Alternate applications with a different fungicide group not in 7 or 11.
	7 + 3	Aprovia Top (difenoconazole + benzovindiflupyr)	10.5 fl oz	54.0 fl oz	7	0.5	Adjuvant recommended. No more than 4 applications per year.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	–
	7 + 12	Miravis Prime (pydiflumetofen + fludioxonil)	11.4 fl/oz	34.2 fl oz	7	12 hr	–

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	See label	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.
	11	Quadris Aframe Many products available (azoxystrobin)	See label	See label	See label	4 hr	Alternate applications with a different fungicide group.
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant.
	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
	11 + 3	Quilt Xcel (azoxystrobin + propiconazole)	Dry bulb: 21 fl oz Green: 26 fl oz	56 fl oz	Dry bulb: 14 Green: 0	0.5	No more than 1 application before alternating to a non-group 11 fungicide.
Damping-off	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.
	2	(iprodione) Many brands available: Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	–
	4	Ridomil Gold Subdue Maxx (mefenoxam)	–	–	–	–	See individual label.
	4 + 11	Uniform (mefenoxam, azoxystrobin)	0.34 fl oz/1000 ft of row	0.34 fl oz/1000 ft of row	–	–	See label.
	12	Cannonball WG (fludioxonil)	7 oz	32 oz	7	0.5	–
	12	Maxim 4FS (fludioxonil)	0.16 fl oz/100 lb seed	–	–	–	Seed treatment only.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
Downy mildew (<i>Peronospora destructor</i>)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Basic Copper 53, Champ DP, Champ F2 FL, Champ WG, C-O-C DF, C-O-C-S WDG, C-O-C WP, Copper Count N, Cuprofix Ultra 40D, Cueva, Kentan DF, Kocide 2000, Kocide 3000, Kocide DF, Nordox, Nordox 75WP, Nu Cop 3L, Nu Cop 50DF, Nu Cop HB	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0	Varies from 4 hr to 2 days.	Frequent use of copper fungicides may cause foliar burn.
	M1 + M2	Top Cop with sulfur (basic copper sulfate + sulfur)	2–3 qt	SEE INDIVIDUAL LABEL	0	1	Do not use during hot weather or within 4 weeks of an oil application.
	M1 + M3	Mankocide (copper hydroxide + mancozeb)	2.25 lb	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate ZN	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	1	Suppression only.
	4 + M1	Ridomil Gold/Copper (mefenoxam + copper hydroxide)	2 lb	–	Green: 7 Dry bulb: 10	2	No more than 0.4 lb a.i./A of metalaxyl or mefenoxam.
	4 + M3	Ridomil Gold MZ WG (mefenoxam + mancozeb)	2.5 lb	–	Dry bulb: 7	2	–
	4 + M5	Ridomil Gold Bravo SC (mefenoxam + chlorothalonil)	See label	–	Dry bulb: 7 Green: 14/21	2	–
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. No more than 5 applications per crop.
	11	Quadris Aframe (azoxystrobin)	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	4 hr	Alternate applications with a different fungicide group.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	11	Reason 500SC (fenamidone)	5.5 fl oz	22 fl oz	7	0.5	Alternate applications with a different fungicide group.
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	SEE INDIVIDUAL LABEL	14	0.5	Alternate applications with a different fungicide group. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant.
	11 + 3	Quilt Xcel (azoxystrobin + propiconazole)	Dry bulb: 21 fl oz Green: 26 fl oz	56 fl oz	Dry bulb: 14 Green: 0	0.5	No more than 1 application before alternating to a non-group 11 fungicide.
	11 + 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	No more than 2 consecutive applications. No more than 6 applications per crop. Suppression only.
	21	Actigard 50WG (acibenzolar-S-methyl)	1 oz	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting. Suppression only.
	21	Ranman (cyazofamid)	3 fl oz	16.5 fl oz	0	0.5	Alternate with fungicides of dissimilar modes of action. No more than 3 sequential applications.
	22 + M5	Zing! (zoxamide + chlorothalonil)	30 fl oz	7.5 qt	7	0.5	No more than 8 applications per year.
	27 + 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	–
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	–
	33	Aliette WDG Linebacker WDG (fosetyl-Al)	3 lb	7 applications	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Do not tank mix with copper fungicides, adjuvants, or foliar fertilizers.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	33	(mono- and dipotassium salts of phosphorous acid) Many brands available: Alude, Confine Extra, Fosiphite, Fungi-Phite, K-Phite 7LP, Oxiphos, Phiticide, Prophyt, Phostrol, Rampart, Reveille	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0	4 hr	See label for details.
	40	Forum (dimethomorph)	6 oz	30 oz	0	0.5	Must be applied in a tank mix with another fungicide active against downy mildew. No more than 2 sequential applications.
	40	Revus (mandipropamid)	8 fl oz	Dry: 32 fl oz Green: 24 fl oz	7	4 hr	No more than 2 consecutive applications before switching to a non-group 40 fungicide. A silicon-based adjuvant must be added at recommended rates.
	40 + 45	Zapro (dimethomorph + ametoctradin)	14 fl oz	42 fl oz	0	0.5	See label.
	43	Presidio (fluopicolide)	4 fl oz	–	2	0.5	No more than 4 applications of Presidio per season.
	49 + M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	No more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
	49 + 40	Orondis Ultra (oxathiapiprolin + mandipropamid)	8 fl oz	Dry: 32 fl oz Green: 24 fl oz	7	4 hr	No more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
Iris Yellow Spot (Iris yellow spot virus)	21	Actigard 50WG (acibenzolar-S-methyl)	1 oz	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting. Suppression only.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
Powdery mildew	M2	(sulfur) Many brands available: Kumulus DF, Microfine Sulfur, Micro Sulf, Microthiol Disperss, Sulfur 90W, Yellow Jacket Wettable Sulfur	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0	1	Do not use during hot weather or within 4 weeks of an oil application.
	3 + 7 + 11	Trivapro (propiconazole + benzovindiflupyr + azoxystrobin)	27.7 fl oz	55.4 fl oz per year	Dry bulb: 7 Green: 14	0.5	Alternate applications with a different fungicide group not in 7 or 11.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	–
	7 + 3	Aprovia Top (difenoconazole + benzovindiflupyr)	10.5 fl oz	54.0 fl oz	7	0.5	Adjuvant recommended. No more than 4 applications per year.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	–
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. No more than 5 applications per crop.
	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
Purple blotch (<i>Alternaria porri</i>)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Basic Copper 53, Champ DP, Champ F2 FL, Champ WG, C-O-C DF, C-O-C-S WDG, C-O-C WP, Copper Count N, Cueva, Cuprofix Ultra 40D, Cuproxat, Kentan DF, Kocide 2000, Kocide 3000, Kocide DF, Kop-hydroxide, Nordox, Nordox 75WP, Nu Cop 3L, Nu Cop 50DF, Nu Cop HB	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0	Varies from 4 hr to 2 days.	Frequent use of copper fungicides may cause foliar burn.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	M1 + M2	Top Cop with sulfur (basic copper sulfate + sulfur)	3 qt	See label	0	1	Do not use during hot weather or within 4 weeks of an oil application.
	M1 + M3	Mankocide (copper hydroxide + mancozeb)	2.25 lb	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate Zn	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	1	–
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	–
	3	(propiconazole) Many brands available: Amtide Propiconazole 41.8% EC, Bumper 41.8 EC, Bumper ES, Fitness, Propimax EC, Propistar EC, Shar-Shield PPZ, Tide Propionazole EC, Tilt	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	0.5	–
	3	(tebuconazole) Many brands available: Monsoon, Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	0.5	–
	3	Cevya (Mefentrifluconazole)	5 fl oz	15 fl oz	7	0.5	–
	3 + M5	Muscle ADV (tebuconazole + chlorothalonil)	1.6 pt	3.2 pt	7	0.5	Apply in a protective fungicide schedule.
	3 + 7	Luna Experience (fluopyram + tebuconazole)	12.8 fl oz	25.6 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	–

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	3 + 7 + 11	Trivapro (propiconazole + benzovindiflupyr + azoxystrobin)	27.7 fl oz	55.4 fl oz per year	Dry bulb: 7 Green: 14	0.5	Alternate applications with a different fungicide group not in 7 or 11.
	4 + M5	Ridomil Gold Bravo SC (mefenoxam + chlorothalonil)	SEE INDIVIDUAL LABEL	–	Dry bulb: 7 Green: 14/21	2	–
	7	Endura (boscalid)	6.8 oz	41 oz	7	0.5	Alternate applications with a different fungicide group.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	See label.
	7 + 3	Aprovia Top (difenoconazole + benzovindiflupyr)	10.5 fl oz	54.0 fl oz	7	0.5	Adjuvant recommended. No more than 4 applications per year.
	7 + 9	Luna Tranquility (fluopyram + pyrimethanil)	27 fl oz	54.7 fl oz	7	0.5	No more than 2 applications per year.
	7 + 11	Merivon Xemium (fluxapyroxad + pyraclostrobin)	11 fl oz	33 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	7 + 12	Miravis Prime (pydiflumetofen + fludioxonil)	11.4 fl oz	34.2 fl oz	7	0.5	–
	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	No more than 2 consecutive applications.
	9	Vanguard WG Vango WG (cyprodinil)	10 oz	28 oz	7	0.5	No more than 2 consecutive applications.
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.
	9 + 12	Switch 62.5WG Alterity 62.5WG Palladium (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last application.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. No more than 5 applications per crop.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	11	Quadris (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11	Reason 500SC (fenamidone)	5.5 fl oz	22 fl oz	7	0.5	Alternate applications with a different fungicide group.
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	SEE INDIVIDUAL LABEL	14	0.5	Alternate applications with different fungicide groups. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant.
	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
	11 + 3	Quilt Xcel (azoxystrobin + propiconazole)	Dry bulb: 21 fl oz Green: 26 fl oz	56 fl oz	Dry bulb: 14 Green: 0	0.5	No more than 1 application before alternating to a non-group 11 fungicide.
	11 + 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	No more than 2 consecutive applications. No more than 6 applications per crop.
	22 + M5	Zing! (zoxamide + chlorothalonil)	30 fl oz	7.5 qt	7	0.5	No more than 8 applications per year.
	27 + 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	–
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	–
	33	Aliette WDG Linebacker WDG (fosetyl-Al)	3 lb	7 appl.	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Do not tank mix with copper fungicides, adjuvants, or foliar fertilizers. Suppression only.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	49 + M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	No more than 2 sequential applications, and do not exceed more than 33% of all fungicide applications with Orondis.
Pythium damping-off	4	(mefenoxam) Many brands available: Apron XL, Ridomil Gold EC, Ridomil Gold SL, Ultra Flourish	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	N/A	2	–
	4	(metalaxyl) Many brands available: Allegiance, Aquire, MetaStar 2E, Sebring 2.65 ST, Sebring 318 FS, Sebring 480 FS	–	–	–	–	–
	11 + 4	Uniform (azoxystrobin + mefenoxam)	SEE INDIVIDUAL LABEL	–	–	–	–
Rhizoctonia damping-off	11	Dynasty (azoxystrobin)	0.38 fl oz/100 lb seed	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	4 hr	Seed treatment only.
	11	Quadris (azoxystrobin)	0.8 fl oz/1000 row feet	SEE INDIVIDUAL LABEL	0	4 hr	–
	11 + 4	Uniform (azoxystrobin + mefenoxam)	SEE INDIVIDUAL LABEL	–	–	–	–
Rust (<i>Puccinia allii</i>)	3	(tebuconazole) Many brands available: Monsoon, Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	0.5	–
	3 + M5	Muscle ADV (tebuconazole + chlorothalonil)	1.6 pt	3.2 pt	7	0.5	Apply in a protective fungicide schedule.
	3 + 7 + 11	Trivapro (propiconazole + benzovindiflupyr + azoxystrobin)	27.7 fl oz	55.4 fl oz per year	Dry bulb: 7 Green: 14	0.5	Alternate applications with a different fungicide group not in 7 or 11.
	7 + 3	Luna Flex (fluopyram + difenoconazole)	13.6 fl oz	27.2 fl oz	7	0.5	–
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. No more than 5 applications per crop.
	11	Quadris (azoxystrobin)	SEE INDIVIDUAL LABEL	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11 + 3	Quilt Xcel (azoxystrobin + propiconazole)	Dry bulb: 21 fl oz Green: 26 fl oz	56 fl oz	Dry bulb: 14 Green: 0	0.5	No more than 1 application before alternating to a non-group 11 fungicide.
Seedling blight	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	Do not apply to exposed bulbs.
	11	Dynasty (azoxystrobin)	0.38 fl oz/100 lb seed	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	4 hr	–
	12	Maxim 4FS Spirato 480 FS (fludioxonil)	0.16 fl oz/100 lb seed	–	–	0.5	–
	14	Botran 5F (DCNA dicloran)	3.5 lb	SEE INDIVIDUAL LABEL	14	SEE INDIVIDUAL LABEL	–
Stemphylium blight (<i>Stemphylium vesicarium</i>)	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	1	–
	3 + 7 + 11	Trivapro (propiconazole + benzovindiflupyr + azoxystrobin)	27.7 fl oz	55.4 fl oz per year	Dry bulb: 7 Green: 14	0.5	Alternate applications with a different fungicide group not in 7 or 11.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	–
	7 + 3	Luna Flex (fluopyram + difenoconazole)	13.6 fl oz	27.2 fl oz	7	0.5	–
	9 + 3	Inspire Super Vango ESQ (cyprodinil + difenoconazole)	20 fl oz	SEE INDIVIDUAL LABEL	Green: 14 Dry bulb: 7	0.5	No more than 2 applications before alternating with a different fungicide group.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	9 + 12	Switch 62.5WG Alterity 62.5WG Palladium (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last application.
	11 + 3	Quadris Top Acadia (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	No more than 1 application before alternating to another fungicide with a non-Qol (Group 11) mode of action.
	11 + 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	No more than 2 consecutive applications. No more than 6 applications per crop.
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	1	Begin sprays at first development and continue on 7- to 14-day schedule.
White rot (<i>Sclerotium cepivorum</i>)	1	(thiophanate-methyl) Many brands available: Incognito 4.5 F, NuFarm T-Methyl 4.5 F, NuFarm T-Methyl 70 WSB, Thiophanate-methyl 85 WDG, Topsin 4.5 FL, Topsin M 70 WDG, Topsin M 70 WP, Topsin M WSB	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	—	3	—
	3	(tebuconazole) Many brands available: Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL	SEE INDIVIDUAL LABEL	7	0.5	—
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	7	0.5	Spray into empty furrow at time of planting.
	7 + 3	Luna Flex (fluopyram + difenocoazole)	13.6 fl oz	27.2 fl oz	7	0.5	—
	9 + 12	Switch 62.5WG Alterity 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	No more than 2 consecutive applications. Do not plant crops not on the label for 30 days after last application.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (Active Ingredients)	Max Rate/Acre		Min. Days to		Remarks ²
			Appl.	Season	Harvest	Reentry	
	11 + M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	SEE INDIVIDUAL LABEL	14	0.5	Alternate applications with different fungicide groups. No more than 3 applications per crop. Do not tank-mix with COC, MSO, or silicon adjuvant.
	12	Cannonball WG (fludioxonil)	7 oz	32 oz	7	0.5	–
	12	Maxim 4FS (fludioxonil)	0.16 fl oz/100 lb seed	–	–	0.5	–
	14	Botran 5F (DCNA dicloran)	3.5 lb	SEE INDIVIDUAL LABEL	14	0.5	–

¹FRAC code (fungicide group): Number (1–54) and letters are used to distinguish the fungicide mode of action groups. All fungicides within the same group (with the same number or letter) indicate the same active ingredient or a similar mode of action. This information must be considered for the fungicide resistance management decisions. Source: FRAC Code List 2025; <http://www.frac.info/> (FRAC = Fungicide Resistance Action Committee).

²Information provided in this table applies only to Florida. Be sure to read a current product label before applying any chemical. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by UF/IFAS Extension nor discrimination against similar products or services not mentioned. OMRI-listed: Listed by the Organic Materials Review Institute for use in organic production.

Table 8. Nonfumigant nematicides for onion and alliums.

Product	Application Directions
Velum (a.i. fluopyram)	Apply 6.5–6.8 fl oz/A either (1) preplant banded or broadcast spray directed to the soil and incorporated into the planting bed or (2) via chemigation into root zone through low-pressure drip, trickle, microsprinkler, or equivalent irrigation. Reapplications at least 14 days apart. No more than 13.7 fl oz/A of Velum (0.446 lb/A fluopyram) per year, regardless of formulation or method of application (soil or foliar). Do not apply within 30 days of harvest.
Biological nematicides	See chapter 19.

Table 9. Fumigant nematicides for onion and alliums.

Nematicide	Broadcast Application ¹		In-the-Row Applications
	Gallons per Acre	fl oz/1000 ft/chisel spaced 12" apart	
Telone II ^{2,3}	9–12	26–35	For any row spacing, application rates given may be concentrated in the row but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-17 ^{2,3}	10.8–17.1	31.8–50.2	For any row spacing, application rates given may be concentrated in the row but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-35 ^{2,3}	13–20.5	38–60	For any row spacing, application rates given may be concentrated in the row but shall never exceed labeled maximum for broadcast applications. Consult the product label for additional detail.
Pic-Clor 60 ^{2,3}	19–31.5	57–90	For any row spacing, application rates should never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Vapam HL	75	–	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration, and flow-modifying directions.
KPam HL	60	–	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration, and flow-modifying directions.
Allyl Isothiocyanate (AITC) Dominus	40	–	For drip or in-row fumigation and crop termination, consult product label for overall rates, drip concentration, and flow-modifying directions.

¹Gallons/acre and fl oz/1000 feet provided only for mineral soils. Higher rates may be possible for heavier-textured (loam, silt, clay) or highly organic soils.

²All of the fumigants mentioned are for retail sale and use only by state-certified applicators or persons under their direct supervision. New supplemental labeling for the Telone products must be in the hands of the user at the time of application. See new label details for additional use restrictions based on soil characteristics, buffer zones, requirements for Fumigant Management Plans (FMP) and Personal Protective Equipment (PPE), mandatory good agricultural practices (GAPs), product and applicator training certification, and other use and rate-modifying recommendations and restrictions.

³Higher application rates are possible in the presence of cyst-forming nematodes.

Rates are believed to be correct for products named, and similar products of other brand names, when applied to mineral soils. Higher rates are required for muck (organic) soils. However, the **grower** has the final responsibility to see that each product is used legally; **read the label** of the product to be sure that you are using it properly.