

Florida Foliage Plant Cultivar: ‘Diamond Bay’ *Aglaonema*¹

R. J. Henny, J. Chen, T. A. Mellich, and M. S. Brennan²

Origin

Aglaonema are indigenous to southeast Asia and comprise 21 species. They are members of the plant family *Araceae*, and are commonly called “Chinese evergreen.” Historically, new *Aglaonema* cultivars were introduced directly from the wild and were largely variants of *Aglaonema crispum*, *A. commutatum*, or *A. modestum*. Control of *Aglaonema* flowering (Henny; 1983) and development of pollination techniques (Henny 1985) led to the production of many new cultivars in the past twenty years by both public and private breeders.

In 1992, *Aglaonema* ‘Silver Bay’ (Henny et al. 1992) an interspecific hybrid, was developed and released by the Foliage Plant Breeding Program of the UF/IFAS Mid-Florida Research and Education Center (MREC)–Apopka. ‘Silver Bay’ remains a very popular cultivar because of its beautiful foliage, good growth habit and excellent performance in interior conditions.

Commercial tissue culture propagation was used to speed the release of *Aglaonema* ‘Silver Bay’ hybrids to foliage producers. *Aglaonema* ‘Diamond Bay’ is a mutant found among a population of tissue-cultured *Aglaonema* ‘Silver Bay’ plants.

The original *Aglaonema* ‘Diamond Bay’ mutant was noticed because it lacked one of the foliar variegation patterns normally present in ‘Silver Bay’ leaves and petioles. The ‘Diamond Bay’ selection was increased by stem cuttings at UF/IFAS MREC and observed for stability. *Aglaonema* ‘Diamond Bay’ selection remained stable while maintaining its new foliar pattern and other desirable traits of the parent. Therefore this cultivar has been selected for release.

Description

Mature *Aglaonema* ‘Diamond Bay’ leaves are lance-shaped 10–12 cm (4–5 in) wide and 28–30 cm (11–12 in) long. Leaf margins are smooth (entire) and the leaf tissue (lamina) on either side of the midrib tends to be of slightly unequal widths, resulting in slight curving of the blade towards the narrower side. Leaves exhibit a central grey-green area that extends out from each side of the midrib to cover slightly more than half of the total leaf surface (Figure 1). Leaf margins are green and meet the lighter grey-green colored central area along irregular borders. The undersides of the leaves are uniform green.

The outer petiole wings are yellow-green and blend into the petiole center, which is a lighter yellow-green. As it nears the leaf blade, the entire petiole color becomes uniform. The stem color blends from areas that are a darker yellow-green to lighter areas in the same color range. These colors

1. This document is ENH1073, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date October 2007. Reviewed November 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. R. J. Henny, professor; J. Chen, professor; Department of Environmental Horticulture; T. A. Mellich, biologist; and M. Brennan, biologist; UF/IFAS Mid-Florida Research and Education Center, Apopka, FL 32703.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

are not normally visible due to the clasping habit of the petiole, which surrounds the stem.



Figure 1. Mature *Aglaonema* 'Diamond Bay' plants grown for 9 months in 6 inch (1.6 L) containers.

Credits: UF/IFAS

Performance

Growth characteristics of *Aglaonema* 'Diamond Bay' were determined using 50 newly rooted stem cuttings with 4–5 leaves per cutting. Plants were potted into 1.6-L (6-inch) plastic pots containing VerGro Container Mix A (Verlite Co., Tampa, Fla. 33610). Plants were grown in a greenhouse under 70% shade, a temperature range of 59°F to 93°F (15°C to 34°C) and on natural photoperiod. Ten plants were grown in a completely randomized design for 9 months from October 2000 until July 2001 at each of five fertilizer levels equivalent to a total of 1.4, 1.8, 2.2, 2.7, and 3.2 g N per pot. Fertilizer levels were derived from a 3:1:2 ratio of N:P:K applied as a liquid at 100 ml per pot per week.

At the termination of the growth tests, canopy height and width, length and width of largest leaf, number of basal shoots and a visual quality rating where 1 = poor; 3 = acceptable (saleable) and 5 = excellent quality were recorded.

Aglaonema 'Diamond Bay' reached marketable size in 9 months. Plants were well-branched, averaging approximately 3 basal shoots per plant. There were no significant differences in canopy height or width, leaf size or basal shoot number due to different fertilizer levels (Table 1). Visual quality showed a significant response to fertilizer level, but all plants were rated as excellent or near excellent in quality. There was a slight loss of quality at the lowest and at the highest nutritional levels.



Figure 2. *Aglaonema* 'Silver Bay', the parent of *Aglaonema* 'Diamond Bay' performs well in commercial installations.

Credits: UF/IFAS

These important tropical ornamental foliage plants can readily adapt to low light and low relative humidity levels encountered under interior conditions. *Aglaonema* 'Diamond Bay' does not require acclimatization prior to installation in interior environments.

Availability

Aglaonema 'Diamond Bay' is intended for commercial producers growing finished plants in 6-inch or 8-inch (1.6 or 3.9 L) containers. A patent application for this cultivar has been submitted to the United States Patent and Trademark Office and plant patent rights have been assigned to the Florida Foundation Seed Producers. Stock plants have been released to University of Florida licensed cooperating Florida tissue culture labs for propagation and distribution. Information about participating labs may be obtained by writing the Florida Foundation Seed Producers, Inc., PO Box 309, Greenwood, FL 32443. Plants for research purposes may be obtained directly from the author.

Literature Cited

- Henny, R.J. 1983. "Flowering of *Aglaonema commutatum* 'Treubii' following treatment with gibberellic acid." *HortScience* 18:374.
- Henny, R.J. 1985. "In vivo pollen germination of *Aglaonema* affected by relative humidity." *HortScience* 20:142–143.
- Henny, R.J., R.T. Poole and C.A. Conover. 1992. "'Silver Bay' *Aglaonema*." *HortScience* 27:1238.

Table 1. Final canopy height and width, length and width of largest leaf, number of basal shoots and plant quality of *Aglaonema* 'Diamond Bay' grown from cuttings in 6-inch (1.6-L) pots with different rates of N for 9 months from October 2000 until July 2001.

N rate (g per pot)^x	Canopy height (cm)	Canopy width (cm)	Leaf length (cm)	Leaf width (cm)	No. basal shoots	Plant Quality^y
1.4	33.5	55.7	29.6	11.4	3.0	4.6
1.8	33.1	54.9	29.7	11.6	3.4	5.0
2.2	35.4	54.7	29.3	11.4	3.4	5.0
2.7	33.1	53.4	29.6	11.3	2.6	4.9
3.2	32.3	54.0	30.1	11.9	3.0	4.8
Significance ^z	NS	NS	NS	NS	NS	S
^x Total amount of N applied during the 9-month production period. ^y Visual rating where 1 = poor, 3 = saleable and 5 = excellent quality. ^z NS = no significant differences; S = significant differences between treatments.						