

New Florida Foliage Plant Cultivar: *Aglaonema* 'Stripes'¹

R. J. Henny, J. Chen, and T. A. Mellich²

Origin

Aglaonema are members of the plant family Araceae, and are commonly called Chinese evergreens. They are indigenous to Southeast Asia and include 21 species. *Aglaonema* species and cultivars are valuable ornamental foliage plants because of their tolerance of environmental conditions in commercial and residential interiorscapes.

Historically, new *Aglaonema* cultivars were introduced by collecting specimens from the wild. These were largely open-pollinated variants of *Aglaonema crispum*, *A. commutatum*, or *A. modestum*. Over the past twenty years, both public and private breeders have produced many new hybridized varieties due to control of *Aglaonema* flowering (Henny 1983) and development of pollination techniques (Henny 1985).

Aglaonema X 'Stripes' resulted from a cross of *Aglaonema* X 'Manila' and *A. nitidum* 'Curtisii'. *Aglaonema* X 'Stripes' combines the foliar variegation patterns of both 'Manila' and 'Curtisii' and is intermediate in size between the two parents. The foliar variegation of 'Stripes' consists of strong bands of silvery white coloration along the lateral veins, overlaying a fainter, more diffuse pattern in the same area (Figure 1). Leaves average about 2.5 times as long as wide and may reach 35 cm in length.



Figure 1. Mature *Aglaonema* X 'Stripes' growing in 2-gallon pots in a Florida foliage plant nursery.

Performance

Growth characteristics of *Aglaonema* X 'Stripes' were determined in three replicated greenhouse trials over a two-year period. Tip cuttings 6 inches (15 cm) long were rooted directly in 6-inch (1.6L) pots filled with a growing medium of 3:2 Canadian peat: Perlite by volume. The soil was amended with 0.9 kg•m⁻³ of Micromax (Sierra Chemical Company, Milpitas, CA) and 4.1 kg•m⁻³ dolomite. Three growth trials were conducted. In all three tests, plants were grown under two shade levels adjusted to 125 and 250 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$ maximum light intensities and were supplied

1. This document is ENH1101, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date June 2008. Reviewed April 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu>.

2. R. J. Henny, professor; J. Chen, associate professor; and T. A. Mellich, biological scientist; Environmental Horticulture Department, UF/IFAS Mid-Florida Research and Education Center, Apopka, FL 32703.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

three fertility levels consisting of 840, 1680, 2525 kg N ha/year, equivalent (2.2, 4.4, or 6.6 g 19N- 6P₂O₅ – 12 K₂O/ pot every three months). Plants were grown in greenhouses under natural photoperiods and a controlled temperature range of 15°C–34°C (63°F–97°F).

In these trials, *Aglaonema* X ‘Stripes’ reached marketable size in 7–10 months, depending on season (Tables 1 and 2). Light level had little influence except that plants grown under 125 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$ were darker green (Table 1). The best fertilizer rate was between 1680 and 2525 mg N/ha/year or 7.1 - 14.2g (19N-3P-10K)/150-mm pot every three months. Overall, plant quality and color grade was better as fertilizer levels increased. Basal shoot production was not affected by fertilizer.

Plants from Trial II were subjected to two weeks of simulated shipping (no light, no water, 15°C) and then held in a growth room at 14.4 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$ light intensity, 12 hours per day for six weeks and hand watered as necessary to simulate an interior environment. The leaves of *Aglaonema* X ‘Stripes’ became darker in the green field, which contrasted the silvery white variegation pattern more sharply. Thus overall plant quality improved during these shipping and interior tests. Data from only two trials is presented here. For more information, see Henny et al. (1988).

Recommendations

Aglaonema X ‘Stripes’ is intended for commercial foliage producers growing 6-inch (1.6L) or 8-inch (3.9L) containers. Best growth can be expected when plants are grown with a light intensity of 125 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$ (equivalent to 80% shade).

Availability

Aglaonema X ‘Stripes’ has been trademarked by the Florida Foundation Seed Producers, Inc. and has been released to Florida tissue culture labs for propagation and distribution. Inquiries regarding participating labs may be obtained by writing the Florida Foundation Seed Producers, Inc., PO Box 309, Greenwood, FL 32443.

References

Henny R. J. 1983. Flowering of *Aglaonema commutatum* ‘Treubii’ following treatment with gibberellic acid. HortScience 18:374.

Henny R. J. 1985. In vivo pollen germination of *Aglaonema* affected by relative humidity. HortScience 20:142–143.

Henny R. J., R. T. Poole, and C. A. Conover. 1988. ‘Stripes’ *Aglaonema*. HortScience 23(95):920–921.

Table 1. The effect of light intensity and fertilizer rate on growth and quality of *Aglaonema* x *Stripes* after growing in a greenhouse for seven months from June until January.

Light Intensity ($\mu\text{mol s}^{-1} \text{m}^{-2}$)	Canopy Height (cm)	Leaf Length (cm)	Leaf Width (cm)	No. Basal Shoots	Visual Color Grade ^z	Visual Plant Quality ^y
125	56	33	12	3.3	4.5	4.0
250	56	33	12	4.4	3.8	4.1
Significance ^x	NS	NS	NS	L**	L**	NS
Fertilizer Rate						
840	54	32	12	3.4	3.5	3.6
1680	57	33	12	4.1	4.4	4.4
2525	57	34	13	4.1	4.7	4.2
Significance ^x	L*	L*	L*	NS	L**	L*

^zVisual color grade where 1 = poor, 3 = good, light green, and 5 = excellent, dark green and yellow contrast.

^yVisual quality where 1 = poor quality, not salable; 3 = good quality salable, and 5 = excellent quality.

^xSignificance where NS, * and ** equal not significant and significant at 5% and 1% level, respectively; L = linear and Q = quadratic.

Table 2. Evaluation of light intensity and fertilizer rate on growth and quality of *Aglaonema* x *Stripes*. Plants were greenhouse grown for 10 months from December–October.

Light Intensity ($\mu\text{mol s}^{-1} \text{m}^{-2}$)	Canopy Height (cm)	Leaf Length (cm)	Leaf Width (cm)	No. Basal Shoots	Visual Color Grade	Visual Plant Quality
125	51	32	13	3.4	3.8	3.9
250	50	31	13	3.7	3.2	3.7
Significance	NS	NS	NS	NS	L**	NS
Fertilizer Rate						
840	50	31	12	3.1	3.2	3.3
1680	51	32	13	3.5	3.6	3.6
2525	51	32	13	4.1	3.6	3.8
Significance	NS	NS	L**	NS	L**Q**	L**

^zVisual color grade where 1 = poor, 3 = good, light green, and 5 = excellent, dark green and yellow contrast.

^yVisual quality where 1 = poor quality, not salable; 3 = good quality salable, and 5 = excellent quality.

^xSignificance where NS, * and ** equal not significant and significant at 5% and 1% level, respectively; L = linear and Q = quadratic