

Florida Foliage Plant Cultivar: *Dieffenbachia* ‘Sparkles’¹

R. J. Henny, J. Chen, and T. A. Mellich²

Origin

Dieffenbachia are important commercial foliage plants consistently ranking in the top 10 in annual sales. Growers encounter few problems during production, finishing, or shipping since the crop tolerates a wide range of cultural conditions. The attractive variegated leaves appeal to buyers, and the plants easily adapt to interior home or office environments. Several generations of *Dieffenbachia* breeding have sought to improve leaf color, leaf patterns, branching habit, and interior performance. *Dieffenbachia* ‘Sparkles’ is the result of such breeding efforts and has been released from the Foliage Plant Breeding Program at the UF/IFAS Mid-Florida Research and Education Center (MREC) in Apopka.

The dominant feature of *Dieffenbachia* ‘Sparkles’ is its bright leaf color (Figure 1). There is high contrast between the light green and dark green variegation patterning, which is highlighted by a white midrib. The plant branches freely, producing up to 15 or more side shoots per main stem. Short spacing between the leaves along the stem creates a very compact and full appearance.

Description

The petioles of *Dieffenbachia* ‘Sparkles’ are green-white in color and have long, wide wings at the base, giving the plant an appearance of having a ribbed stem. Leaf size averages 20–22 cm (7–8 in.) long and 9–10 cm (4–5 in.) wide. Leaves have a predominantly light green background center color, with random splotches of deep green and irregular blotches

of white occurring parallel to the primary veins. This results in an attractive paisley, mottled appearance. The midrib of each leaf appears as a straight, solid, nearly white stripe due to a lack of chlorophyll. Leaves have continuous, deep green margins, and the edges along the leaf lack undulations, resulting in erect, smooth leaves not commonly found in *Dieffenbachia* varieties. *Dieffenbachia* ‘Sparkles’ plants branch freely, producing multiple side shoots per main stem. Each stem has a high density of leaves, and the short spacing between the leaves along each stem causes *Dieffenbachia* ‘Sparkles’ to be very compact. ‘Sparkles’ retains its lower leaves along the stem even with advancing age.



Figure 1. A mature *Dieffenbachia* ‘Sparkles’ grown in a 6 in. diameter container.

Credits: UF/IFAS

1. This document is ENH1173, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date September 2010. Reviewed November 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. R. J. Henny, professor; J. Chen, professor; Department of Environmental Horticulture; and T.A. Mellich, biologist, UF/IFAS Mid-Florida Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Growth characteristics and performance of *Dieffenbachia* 'Sparkles' were determined from three greenhouse production trials. Trials were initiated using uniform liners obtained from a commercial tissue-culture laboratory. Thirty 10-week-old liners were potted into 6 in. diameter (1.6 L) plastic pots filled with Vergro Container Mix A. Plants were treated with low, medium, or high rates of Osmocote® 19:6:12 (nitrogen: phosphorus: potassium). Osmocote® was applied at 2.5, 4.2, or 5.9 g per pot (5.5 g = approximately 1 tsp.). Fertilizer was applied when the experiment started and again after 12 weeks. There were 10 replications per fertilizer level. Tests were conducted in a greenhouse shaded to exclude 70% of sunlight, under natural photoperiod, and at a temperature range of 15°C–34°C (59°F–93°F). Plants were grown for 24 weeks, at which time plant height, leaf size, branching, and quality were measured. Data were analyzed using the ANOVA procedures of the SAS program.

Dieffenbachia 'Sparkles' performed well over a range of fertilization rates (Table 1). Plant height, leaf size, and number of basal shoots were not significantly different among the low, medium, and high fertilizer rates. The lack of significant statistical differences in growth rates between the fertilizer levels indicates that 'Sparkles' yields full and compact plants for commercial growers over a wide range of conditions (Figure 2). All plants in this growth test performed well and received an "excellent" rating for visual quality, where 1 = poor, 3 = acceptable (saleable), and 5 = excellent.



Figure 2. *Dieffenbachia* 'Sparkles' performs well in commercial production.

Credits: UF/IFAS

After greenhouse testing, *Dieffenbachia* 'Sparkles' plants were placed in a growth chamber to evaluate their performance in a consumer home or office environment. Plants received light from cool white fluorescent bulbs for 12 hours per day. Plants were evaluated after three months. Quality ratings were excellent. All plants maintained compact growth habit, leaf retention, and good color definition. This cultivar performs exceptionally well in interior environments.

Availability

Dieffenbachia 'Sparkles' is protected under U.S. Plant Patent No. 9050. Commercial foliage plant growers can obtain licensed stock for *Dieffenbachia* 'Sparkles' through Florida tissue-culture laboratories. Names of cooperating laboratories may be obtained from the Florida Foundation Seed Producers, PO Box 309, Greenwood, FL 32443. The dense branching and compact growth habit of *Dieffenbachia* 'Sparkles' make it suitable for production in 4 in. diameter (0.5 L) pots or 6 in. diameter (1.6 L) pots.

Additional Information

Henny, R. J., C. A. Conover, and R. T. Poole. 1987. "Triumph' *Dieffenbachia*." *HortScience* 22:965–966.

Henny, R. J., C. A. Conover, and R. T. Poole. 1987. "Victory' *Dieffenbachia*." *HortScience* 22:967–968.

Henny, R. J., R. T. Poole, and C. A. Conover, 1988. "Tropic Star' *Dieffenbachia*." *HortScience* 23:919–920.

Henny, R. J., R. T. Poole, and C. A. Conover, 1989. "Starry Nights' *Dieffenbachia*." *HortScience* 24:522–524.

Table 1. Size and quality of *Dieffenbachia* ‘Sparkles’ grown for 24 weeks in 6 in. diameter pots.

| Fertilizer (grams per pot) | Plant height (inches) | Leaf length (inches) | Leaf width (inches) | Number of basal shoots | Final plant quality ^a |
|-------------------------------|--------------------------|----------------------------|---------------------------|---------------------------|-------------------------------------|
| 2.5 (low) | 13.0 | 7.9 | 3.8 | 18.1 | 4.8 |
| 4.2 (medium) | 13.2 | 7.8 | 3.8 | 16.9 | 4.9 |
| 5.9 ^b (high) | 13.0 | 7.8 | 3.7 | 17.5 | 5.0 |
| Significance ^c | None | None | None | None | None |

^a Quality = a visual quality rating, where 1 = *poor*, 3 = *acceptable* (saleable), and 5 = *excellent*.

^b One level tsp. equals approximately 5.5 g.

^c Data were analyzed using ANOVA procedures of the SAS program.