

Preventing Foodborne Illness: *Bacillus cereus*¹

Karuna Kharel, Kiara S. Bookman, Renée Goodrich Schneider, Rachael Silverberg, Ploy Kurdmongkoltham, Bruna Bertoldi, and Keith R. Schneider²

This is one in a series of fact sheets discussing common foodborne pathogens of interest to food handlers, processors, and retailers.

What is *Bacillus cereus*?

Bacillus cereus is a Gram-positive, facultative anaerobic bacterium characterized by large, rod-shaped cells and an ability to form heat-resistant endospores. Since this bacterium is commonly widespread in the environment and is often found in soil, it is naturally present in a wide range of food products of both plant and animal origin. *B. cereus* grows best in a temperature range of 39°F (4°C) to 118°F (48°C). Optimal growth occurs within the narrower temperature range of 82°F (28°C) to 95°F (35°C) and a pH range of 4.9 to 9.3 (FDA 2012).

While there are numerous known species in the genus *Bacillus*, only two, *B. anthracis* and *B. cereus*, are associated with human diseases. *Bacillus anthracis*, though pathogenic, is rarely linked to foodborne illness. However, *Bacillus cereus* is the known source of two distinct types of foodborne illness. Both illnesses are associated with the ingestion of a distinct toxin produced by the bacteria. The first form is emetic, which is characterized by nausea, vomiting, and abdominal cramps. The emetic form has a short onset time of about one to six hours after consumption of contaminated foods and is caused by the ingestion of a

pre-formed toxin that contaminates before it is eaten. The temperature for emetic toxin (cereulide) production ranges from 77°F (25°C) to 86°F (30°C) (FDA 2017b). The second form causes diarrhea, has a longer onset time of about 6 to 15 hours, and can last approximately 24 hours. This illness is caused by enterotoxins (toxins that specifically affect the intestinal mucosa) produced by *B. cereus* inside the host after ingestion (FSANZ 2013; Naranjo et al. 2011).



Figure 1. A colony of *Bacillus cereus*.
Credits: Pete Seidel [PHIL #12378]

1. This document is FSHN15-06, one of a series of the Department of Food Science and Human Nutrition, UF/IFAS Extension. Original publication date August 2015 and November 2024. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Karuna Kharel, postdoctoral associate; Kiara S. Bookman, former technician; Renée Goodrich-Schneider, professor, Department of Food Science and Human Nutrition; Rachael Silverberg, former lab technician (former author), Department of Food Science and Human Nutrition; Ploy Kurdmongkoltham, former technician (former author), Doctor of Plant Medicine program; Bruna Bertoldi, former technician (former author); and Keith R. Schneider, professor, Department of Food Science and Human Nutrition; UF/IFAS Extension, Gainesville, FL 32611.

Transmission of Foodborne Illness

Due to its ubiquity in the environment, *B. cereus* is easily spread to many types of fresh and processed food products. It is found in the environment and on many foods, including meat, cereal dishes, vegetables, milk products, puddings, and soups, but it does not usually pose a health risk. Illness usually occurs when food is improperly cooked and stored in the danger zone (41°F to 135°F) (FDA 2017b) for an extended period of time (Marriot and Gravani 2006).

Transmission of the diarrheal form of the illness is caused by ingestion of viable cells or spores. These cells grow in the body and secrete toxins (whereas the emetic form produces a toxin on the food and is ingested preformed) (Bottone 2010). The diarrheal form typically requires a concentration of 100,000 cells/g or greater to cause illness (FDA 2012). In most instances, illnesses associated with *B. cereus* are almost exclusively the result of improper food handling, storage, and cooling that allow for the growth of *B. cereus* and/or the production of emetic toxins.

Cooking contaminated food at recommended temperatures destroys the vegetative cells of *B. cereus*. Heat-resistant endospores of this bacteria are more likely to survive cooking and start growing again (germinate) when the food is cooled gradually over an extended period of time. The emetic toxin (cereulide) is heat stable up to 250°F (121°C) (Rajkovic 2014). If food is stored at improper temperatures for several hours before serving (> 41°F [5°C] for cold food; < 135°F [57°C] for hot food), hazardous concentrations of bacterial cells or toxins could develop before the food is eaten (HPSC 2012). The temperature range between 135°F (57°C) and 41°F (5°C) is known as the danger zone for food storage and thus should be avoided (FDA 2017b).

Additional sources of foodborne illness outbreaks could be improper hygiene during food canning and inadequate reheating of food in congregate settings, such as restaurants and schools (HPSC 2012).

What foods have been commonly associated with *Bacillus cereus*?

A broad range of foods have been implicated as vehicles of *B. cereus*. These include boiled or fried rice, cooked vegetables and meats, pasta, vanilla sauce, custards, casseroles, pastries, salads, soups, ice cream, and herbs and spices (FDA 2012). The emetic form of illness (vomiting) is most frequently associated with improperly refrigerated starch dishes, such as fried rice, whereas the diarrheal form

of illness is associated with foods containing meat and vegetables (Todar 2012).

There are many examples of outbreaks caused by this organism. In 2014 in China, 139 people reported nausea, vomiting, and diarrhea after consuming fermented black beans. Three strains of *B. cereus* were isolated from the outbreak, including two strains known to produce the emetic toxin and one known to produce the diarrheal enterotoxin (Zhou et al. 2014). Another severe outbreak occurred in 2003 in Leuven, Belgium, where five children in one family became ill after consuming pasta salad that contained the emetic toxin. The pasta was stored in a refrigerator set at 57.2°F (14°C), which is much higher than the proper storage temperature of 41°F (5°C). The youngest child, a 7-year-old girl, died in the hospital only 13 hours after consuming the pasta (Dierick et al. 2005). The Centers for Disease Control and Prevention (CDC) estimates that *B. cereus* was responsible for 63,400 cases of foodborne illness and 20 hospitalizations in the United States each year between 2000 and 2008 (Scallan et al. 2011; CDC 2012). In addition, a 2012 CDC surveillance annual report lists two outbreaks and 24 illnesses involving *B. cereus* in the United States (Bennett et al. 2014). However, many cases are never reported or diagnosed in a clinical setting because the associated symptoms are usually mild and/or short-lasting. Table 1 outlines recent *B. cereus* foodborne outbreaks.

Preventing Contamination by *B. cereus*

Because *B. cereus* endospores are extremely heat resistant, they will likely survive cooking at temperatures that would otherwise destroy foodborne pathogen cells. Heat resistance increases with increasing salinity (presence of salt) and decreases with increasing acidity. *Bacillus cereus* spores can germinate when exposed to heat or improper handling; therefore, the 2017 Food Code recommends that hot foods be maintained at a temperature of 135°F (57°C) or above and cold foods be maintained at a temperature of 41°F (5°C) or below (FDA 2017b).

According to the National Institutes of Health (NIH), the National Institute of Allergy and Infectious Diseases (NIAID), the National Food Processors Association (NFPA), and the FDA 2017 Food Code, the suggestions below are good methods and practices that destroy *B. cereus*:

- Steaming under pressure, roasting, frying, and grilling foods will destroy the vegetative cells and spores if temperatures within foods equal 145°F (63°C).

- Since foods containing the emetic toxin need to be heated to 249°F (121°C) for more than 80 minutes, reheating foods until they are steaming is not enough to inactivate the emetic toxin, and efforts should be devoted to prevention (Rajkovic 2014).
- Heating (i.e., cooking) to 145°F (63°C) and reheating to 165°F (74°C) for 15 seconds will destroy the vegetative (actively growing) cells. Once cooked, the rapid cooling of the product will prevent any present spores from germinating. However, note that foods cannot be made safe to eat if toxins have already been formed.
- Keep hot foods above 135°F (57°C) and cold foods below 41°F (5°C) to prevent the formation of spores.
- Refrigerate leftovers properly by cooling rapidly to 41°F (5°C) or below. Make sure leftover containers are not stacked close together; this will allow for adequate airflow around the food and ensure rapid cooling.

Good Practices for Food Product Receiving, Handling, Processing, and Storage

The FDA defines current Good Manufacturing Practices (GMPs) in the Code of Federal Regulations, Title 21, Part 117 (FDA 2017a). These GMPs outline minimal sanitation requirements in FDA-inspected food handling and processing facilities. It is recommended that more specific and stringent standard operating procedures (SOPs) be developed for individual facilities. In addition, the sanitation recommendations for food service and retail food facilities outlined in the FDA 2017 Food Code were adopted as law by sixteen states, including Florida, as of December 2022 (FDA 2022). The FDA updated the 2013 Food Code with the release of the 2017 Food Code. Since the adoption of the new versions of the Food Code varies by state, it is important that each facility checks with the appropriate state or local regulatory authority to make sure they are using the guidelines required by their jurisdiction. The Florida statutes can be found at <http://www.leg.state.fl.us/statutes>, Title 33: Chapter 509.

In addition to setting and adhering to strict sanitation requirements in the facility, a retail establishment should also develop SOPs for the receiving and storing of food products and ingredients. If food is being processed, appropriate controls and requirements should be established and strictly adhered to. The FDA 2017 Food Code outlines appropriate processing and cooking requirements for food service.

Processing

One of the easiest ways to prevent foodborne illness associated with *B. cereus* is by ensuring that foods are cooked thoroughly and cooled rapidly. One of the leading causes of foodborne infections and intoxications by *B. cereus* is the improper holding of cooked foods. Refer to sections 3-401.11 to 3-401.14 and 3-403.11 of the 2017 Food Code.

- Raw animal meats should be cooked to an internal temperature of 145°F (63°C) or above and be held for a minimum of 15 seconds at that temperature.
- Fruits and vegetables prepared for hot holding should be cooked to an internal temperature of 135°F (57°C).
- Hold all hot food at a temperature of 135°F (57°C) or higher.
- Chill cooked foods promptly. Cooked food, especially meat, should not be left out at room temperature for more than two hours.
- Reheating previously cooked food so that all parts of the food reach an internal temperature of at least 165°F (74°C) for 15 seconds will kill most harmful bacteria but may not be adequate to inactivate existing toxins (Rajkovic 2014). (Note: Contaminated food should never knowingly be served, even if reheated properly. If a food is thought to be contaminated, it should be disposed of. When in doubt, throw it out!)

Storage

Once a product has been received or processed, it should be properly displayed or stored. There are some general guidelines governing these practices as well. Refer to sections 3-501.13 and 3-501.14 of the 2017 Food Code.

- Frozen food should remain frozen until used.
- If frozen food is displayed in a refrigerated case and allowed to thaw, the food should remain at 41°F (5°C) or below.
- Frozen food should be thawed at a temperature of 41°F (5°C) or below or under running water at a temperature of 70°F (21°C) or below.
- The product can be thawed as part of the cooking process.
- Food should be cooled from 135°F (57°C) to 70°F (21°C) within a period of two hours.
- Overall, the cooling process from 135°F (57°C) to 41°F (5°C) should take no more than six hours.
- Cooked product should be maintained above 135°F (57°C) while displayed and stored at or under 41°F (5°C) for no more than seven days.

- Properly label all stored food products.
- Always remember, when in doubt, throw it out.

For more specific recommendations, consult the 2017 Food Code: <https://www.fda.gov/media/110822/download>.

References

Bennett, S. D., K. Manikonda, E. Mungai, D. Dewy-Mattia, and L. H. Gould. 2014. *Surveillance for Foodborne Disease Outbreaks United States, 2012: Annual Report*. Centers for Disease and Control Prevention. https://stacks.cdc.gov/view/cdc/23299/cdc_23299_DS1.pdf

Bottone, E. J. 2010. “*Bacillus cereus*, a Volatile Human Pathogen.” *Clinical Microbiology Reviews* 23 (2): 382–398. <https://doi.org/10.1128/CMR.00073-09>

British Columbia Centre for Disease Control (BCCDC). 2002. “Foodborne Illness Outbreaks: *Bacillus cereus*.” *Milk Safety Notes*. Accessed February 14, 2024. http://www.bccdc.ca/resource-gallery/Documents/Educational%20Materials/EH/FPS/Dairy/Outbreak_Bacillus_Milk.pdf

Carroll, L. M., M. Wiedmann, M. Mukherjee, D. C. Nicholas, L. A. Mingle, and J. Kovac. 2019. “Characterization of Emetic and Diarrheal *Bacillus cereus* Strains from a 2016 Foodborne Outbreak Using Whole-Genome Sequencing: Addressing the Microbiological, Epidemiological, and Bioinformatic Challenges.” *Frontiers in Microbiology* 10: 420842. <https://doi.org/10.3389/fmicb.2019.00144>

Centers for Disease Control and Prevention (CDC). 2012. *Pathogens Causing US Foodborne Illnesses, Hospitalizations, and Deaths, 2000–2008*. Archived at Wayback Machine, April 28, 2024. <https://web.archive.org/web/20240428062156/http://www.cdc.gov/foodborneburden/PDFs/pathogens-complete-list-01-12.pdf>

Centers for Disease Control and Prevention (CDC). 1986. “*Bacillus cereus* – Maine.” *Morbidity and Mortality Weekly Report* 35 (25): 408–410. <https://www.cdc.gov/mmwr/preview/mmwrhtml/00000754.htm>

Centers for Disease Control and Prevention (CDC). 1994. “Epidemiologic Notes and Reports: *Bacillus cereus* Food Poisoning Associated with Fried Rice at Two Child Day Care Centers—Virginia, 1993.” *Morbidity and Mortality Weekly Report* 43 (10): 177–178. <http://www.cdc.gov/mmwr/PDF/wk/mm4310.pdf>

Centers for Disease Control and Prevention (CDC). 2024. “BEAM (Bacteria, Enterics, Ameba, and Mycotics) Dashboard.” National Center for Emerging Zoonotic Infectious Diseases (NCEZID). Accessed November 14, 2024. <https://www.cdc.gov/ncezid/dfwed/BEAM-dashboard.html>

Dierick, K., E. Van Coillie, I. Swiecicka, G. Meyfroidt, H. Devlieger, A. Meulemans, G. Hoedemaekers, L. Fourie, M. Heyndrickx, and J. Mahillon. 2005. “Fatal Family Outbreak of *Bacillus cereus*-Associated Food Poisoning.” *Journal of Clinical Microbiology* 43 (8): 4277–4279. <https://doi.org/10.1128/JCM.43.8.4277-4279.2005>

Delbrassinne, L., N. Botteldoorn, M. Andjelkovic, K. Dierick, and S. Denayer. 2015. “An Emetic *Bacillus cereus* Outbreak in a Kindergarten: Detection and Quantification of Critical Levels of Cereulide Toxin.” *Foodborne Pathogens and Disease* 12 (1): 84–87. <https://doi.org/10.1089/fpd.2014.1788>

Food and Drug Administration (FDA). 2012. “*Bacillus cereus* and Other *Bacillus* spp.” *Bad Bug Book: Handbook of Foodborne Pathogenic Microorganisms and Natural Toxins*. <http://www.fda.gov/downloads/Food/FoodborneIllness-Contaminants/UCM297627.pdf>

Food and Drug Administration (FDA). 2017a. Code of Federal Regulations Title 21-Part 117—*Current Good Manufacturing Practice, Hazard Analysis, and Risk-Based Preventive Controls for Human Food*. Accessed February 28, 2024: <https://www.ecfr.gov/on/2017-12-31/title-21/chapter-I/subchapter-B/part-117>

Food and Drug Administration (FDA). 2017b. *Food Code 2017*. US Public Health Service. Accessed March 15, 2024. <https://www.fda.gov/media/110822/download>

Food and Drug Administration (FDA). 2022. *Adoption of the FDA Food Code by State and Territorial Agencies Responsible for the Oversight of Restaurants and/or Retail Food Stores*. Archived at Wayback Machine, October 5, 2023. <https://web.archive.org/web/20231005112241/https://www.fda.gov/media/107543/download>

Food Standards Australia New Zealand (FSANZ). 2013. *Bacillus cereus*. Last updated May 2013. <https://www.foodstandards.gov.au/sites/default/files/publications/Documents/Bacillus%20cereus.pdf>

Health Protection Surveillance Center (HPSC). 2012. “*Bacillus cereus* Food-borne Illness.” *Infectious Intestinal Disease: Public Health and Clinical Guidance*. <http://www.hpsc.ie/A-Z/Gastroenteric/GastroenteritisorIID/Guidance/File,13492,en.pdf>

Marriot, N. G., and R. B. Gravani. 2006. “*The Relationship of Microorganisms Sanitation*.” *Principles of Food Sanitation*. 5th ed. Springer Science+Business Media, Inc.

Naranjo, M., S. Denayer, N. Botteldoorn, L. Delbrassinne, J. Veys, J. Waegenaere, N. Sirtaine, et al. 2011. “Sudden Death of a Young Adult Associated with *Bacillus cereus* Food Poisoning.” *Journal of Clinical Microbiology* 49 (12): 4379–4381. <https://doi.org/10.1128/JCM.05129-11>

Rajkovic, A. 2014. “Microbial Toxins and Low Level of Foodborne Exposure.” *Trends in Food Science & Technology* 38 (2): 149–157. <https://doi.org/10.1016/j.tifs.2014.04.006>

Scallan, E., R. B. Hoekstra, F. J. Angulo, R. V. Tauxe, M. A. Widdowson, S. L. Roy, J. L. Jones, and P. M. Griffin. 2011. “Foodborne Illness Acquired in the United States—Major Pathogens.” *Emerging Infectious Diseases* 17 (1): 7–15. <https://doi.org/10.3201/eid1701.P11101>

Todar, K. 2012. “*Bacillus cereus* Food Poisoning.” In *Todar’s Online Textbook of Bacteriology*. Accessed February 28, 2024.

Williams, J. 2011. “Outbreak at Okeechobee Correctional Institute.” *Food for Thought: An E-Bulletin from the Food and Waterborne Disease Program 1*. Florida Department of Public Health. Accessed March 15, 2024. https://outbreakdatabase-com.s3.amazonaws.com/Florida_Dept._of_health_Publication_Food_for_Thought_Issue_1_February_2011_See_page_2.pdf

Zhou, G., K. Bester, B. Liao, Z. Yang, R. Jiang, and N. B. Hendriksen. 2014. “Characterization of Three *Bacillus cereus* Strains Involved in a Major Outbreak of Food Poisoning After Consumption of Fermented Black Beans (*Douchi*) in Yunan, China.” *Foodborne Pathogens and Disease* 11 (10): 769–774. <https://doi.org/10.1089/fpd.2014.1768>

Table 1. Significant *Bacillus cereus* Foodborne Outbreaks.¹

Year	Location	Cases	Deaths	Toxin	Food/Source
1985	US	11	0	ND ²	Hibachi steak
1988	Canada	37	0	ND	Milkshake
1989	US	55	0	ND	Cornish game hens
1989	Canada	74	0	ND	Milk
1991	US	139	0	Diarrheal	Barbecued pork
1993	US	14	0	Emetic	Fried rice
1998	France	44	3	Diarrheal	Vegetable puree
2000	Italy	173	0	Diarrheal	Cake
2003	Belgium	5	1	Emetic	Pasta salad
2005	US	26	1	ND	Turkey
2006	US	26	0	Emetic	Fried rice
2007	Australia	1	1	ND	Asparagus sauce
2008	Belgium	1	1	Emetic	Spaghetti
2010	US	103	0	Emetic	Rice
2010	US	17	0	ND	Pork/chicken
2011	US	58	0	ND	ND
2012	Belgium	20	0	Emetic	Rice, cucumber, chicory
2013	UK	93	0	ND	Rice
2014	England	14	1	ND	Intravenous liquid (TPN ³)
2014	Canada	44	0	Emetic	Fried rice
2014	China	139	0	Emetic	Fermented black beans
2015	US	22	0	ND	Enchilada, cheese, salsa
2015	US	3	0	ND	Rice
2016	US	179	0	ND	Refried beans
2017	US	105	0	ND	Tortilla
2018	US	106	90	ND	Ham
2019	US	116	0	ND	Rice
2020	US	280	0	ND	Coleslaw
2021	US	324	0	ND	Chicken, beans, chili

¹ Sources: Delbrassinne et al. (2015); Zhou et al. (2014); FSANZ (2013); Williams (2011); Naranjo et al. (2011); Dierick et al. (2005); BCCDC (2002); CDC (1994); CDC (1986); CDC (2017); Carroll et al. (2019); CDC (2024)

² ND—No data.