

2025–2026 Quick Reference Guide to Postemergence Herbicides for Citrus Weed Control¹

Products recommended in the *Florida Citrus Production Guide* and their effects on weed management.

	Herbicide ^a	MOA ^b	REI ^c Hours	PHI ^d Day(s)	Weeds Controlled		Comments	Suggested Rate per Acre			
					Grasses	Broadleaf					
Nonselective Systemic Herbicides	Glyphosate —Undertree	G (9)	Varies ^e	1	X	X	Avoid contact with citrus fruit, foliage, and green bark. Rainfall within 1–6 hours after application may reduce effectiveness.	Annual weeds: 0.75–1.5 lb A.E. ^f Perennial weeds: 1.5–3.75 lb A.E. See product label for annual maximum rate			
	Glyphosate —Chemical mowing	G (9)	Varies	1	X	X	Do NOT mow within 1 week before or after treatment.	Bahia grass: 0.125 lb A.E. followed by 2nd application 45 days later Bermudagrass: 0.125–0.37 lb A.E.			
	Glyphosate —Wiping	G (9)	Varies	1	X	X	Use wipers to remove tall growing and difficult weeds.	5%–10% solution—carpet wiper 50%–100% solution—panel wiper			
	Glyphosate —Spot treatment	G (9)	Varies	1	X	X	Avoid contact with citrus fruit, foliage, and green bark.	1%–2% solution			
	Glyphosate plus 2,4-D Landmaster II	G, O (9, 4)	48	7	X	X	Apply with shielded and hooded sprayers in citrus middles or under the trees. User must have supplemental labeling at time of application. See label for min. distance from susceptible crops, record-keeping requirements, and additional restrictions. Rainfall or irrigation within 4 hours may reduce effectiveness. Heavy rainfall or irrigation within 2 hours will wash product off foliage, requiring repeat treatment.	Use recommended rate in 10–40 GPA Annual weeds: 1–8 qt Perennial weeds: 4–8 qt Maximum of 8 qt/year			
Nonselective Contact Herbicides	Carfentrazone-ethyl Aim EC	E (14)	12	3	X	X	Adjuvant is required such as nonionic surfactant or crop oil concentrate. Avoid contact with green tissue or fruit. Finished spray volume of at least 20 GPA required.	Rate per application	Maximum rate/year	Max. # appl./yr	Min. time btwn. appl.
								Max 2.0 fl oz	7.9 fl oz		14 days
	Glufosinate-ammonium Rely 280	H (10)	12	14	X	X	Warm temperatures, high humidity, and bright sunlight improve performance. Avoid contact or spray drift with green bark, stems, or foliage. Spot treatment: 1.7 fl oz per gallon of water. Apply to undesirable vegetation foliage until wet but prior to runoff.	48–82 fl oz	246 fl oz (4.5 lb a.i.)	3 at max rate	14 days
	Paraquat Gramoxone SL 2.0	D (22)	24	—	X	X	Addition of surfactant or crop oil concentrate is essential for maximum contact activity. Avoid contact with citrus fruit, foliage, and green bark. Per new labeling requirement, applicators must complete mandatory training program and be certified applicators of restricted-use pesticides.	2.5–4.0 pt	20 pt	5	
Selective Systemic Herbicides	Fluazifop Fusilade DX 2 E	A (1)	12	14	X		Do not apply to grasses under stressed conditions. For spot treatment use 1% v/v solution with 1% crop oil concentrate or 0.25% nonionic surfactant in 30–40 GPA.	24 fl oz	72 fl oz	3	21 days
	Mesotrione Broadworks 4 L	F2 (27)	12	1		X	Allow at least 12 weeks between applications at 6 fl oz/A and at least 6 weeks between applications of 6 fl oz/A and subsequent applications of 3 fl oz/A. Use of a crop oil at 1% v/v or nonionic surfactant at 0.25% is recommended. Addition of ammonium sulfate is suggested.	Max 6 fl oz at 1st appl.	12 fl oz	3	6 weeks
	Sethoxydim Poast Plus 1.0 EC	A (1)	12	15	X		Repeated applications at 3–4 week intervals may be required to control some species. Carrier volume should not exceed 20 GPA. Spot treatment use 1.5%–2.25% v/v solution of Poast Plus w/1% crop oil concentrate.	2.25–3.75 pt	15 pt		
	Saflufenacil Treevix	E (14)	12	—		X	Use methylated seed oil and ammonium sulfate to optimize burndown activity. Avoid contact with citrus fruit, foliage, flowers, buds, and green bark. Increased efficacy at water volumes of 20 to 40 GPA. *Additional Dormant Period Application: May be applied sequentially up to 4 times per year only if 1 or 2 applications occur from postharvest to the beginning of bloom.	Max 1.0 oz	3.0 oz (4.0 oz*)	3 (4*)	21 days
	2,4-D Embed Extra	O (4)	48	7		X	Controls annual and perennial broadleaf weeds. Apply when weeds are in their early and active growth stages, and limit application to groves established for at least one year. Only apply when there is no risk of spray drift. Avoid contact with tree trunks and foliage.	Max 4.0 pt	n/a	2 at max rate	75 days

Always read individual product labels for complete directions for use, restrictions and limitations, and tank-mix and additive recommendations.

^a All listed pesticides are registered and trademarked products.

^b Mode of action class (MOA) for citrus pesticides from the Herbicide Resistance Action Committee (HRAC). Weed Science Society of America (WSSA) classification is provided in the parentheses.

^c Restricted Entry Interval

^d Preharvest Interval

^e REIs vary from 4–12 hours depending upon glyphosate formulation chosen. Check the individual product label to confirm REI.

^f Acid Equivalence—**conversion factors on reverse**

Nonselective Postemergence Systemic Herbicides: Glyphosate Conversions

Products recommended in the *Florida Citrus Production Guide* and their effects on weed management.

<ul style="list-style-type: none"> • Glyphosate rates are often stated in pounds acid equivalent (A.E.) per acre instead of active ingredient (a.i.) per acre because only the “glyphosate acid” portion of product formulation is herbicidally active. • The glyphosate concentration in acid equivalent (A.E.) is specified normally on the product label’s front panel in the “ingredient statement” as pounds per gallon (lb/gal) of the glyphosate acid in the product. • Use this table to calculate the amount of the product needed to achieve the recommended rate of A.E. per acre. 	Acid Equivalence Conversions for Glyphosate							
	Acid Equivalence (A.E.) (lb/gal)	Rate per Treated Acre in A.E.						
		0.094 lb	0.188 lb	0.282 lb	0.37 lb	0.75 lb	1.5 lb	2.25 lb
		Amount of product to equal the above pounds of A.E.						
	3.0	4 oz	8 oz	12 oz	16 oz	32 oz (1 qt)	64 oz (2 qt)	96 oz (3 qt)
	3.7	3.25 oz	6.5 oz	9.75 oz	13 oz	26 oz	52 oz	78 oz
	4.0	3 oz	6 oz	9 oz	12 oz	24 oz	48 oz	72 oz
	4.5	2.7 oz	5.4 oz	8 oz	10.5 oz	21.3 oz	43 oz	64 oz
	5.0	2.4 oz	4.8 oz	7.2 oz	9.5 oz	19.2 oz	38.4 oz	57.6 oz

Various formulations of glyphosate are currently registered for use in Florida citrus. It is important to adjust the application rate used according to the product concentration. A product concentration is stated in pounds per gallon of acid equivalent (A.E.) on the label.

1. This document is HS1410, one of a series of the Department of Horticultural Sciences, UF/IFAS Extension. Original publication date March 2021. Revised August 2023 and August 2024. Reviewed August 2025. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Ramdas Kanissery, associate professor, Department of Horticultural Sciences, UF/IFAS Southwest Florida Research and Education Center; Camille E. McAvoy, former laboratory technician, UF/IFAS SWFREC; Jamie D. Burrow, Extension program manager, UF/IFAS Citrus Research and Education Center; Stephen H. Futch, Extension agent (retired), UF/IFAS CREC; Brent A. Sellers, professor, Department of Agronomy, and center director, UF/IFAS Range Cattle Research and Education Center; and S. Shea Teems, lab manager, UF/IFAS SWFREC; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition. Use herbicides safely. Read and follow directions on the manufacturer’s label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county’s UF/IFAS Extension office.