

Yellow-Legged Hornet (suggested common name), *Vespa velutina* (Lepeletier 1836) (Insecta: Hymenoptera: Vespidae)¹

Krystal Ashman, Oliver Keller, and Cameron Jack²

Introduction

The yellow-legged hornet, *Vespa velutina* (Lepeletier) (Figure 1), is a pest of concern outside of its native range. *Vespa velutina* is native to Southeast Asia (Monceau et al. 2014) and has invaded several regions in Europe, first appearing in France in 2004 (Monceau et al. 2014). As a generalist predator, they are a pest of honey bees and a major concern to many beekeepers. *Vespa velutina* has not been intercepted in North America, but it is believed to have high invasion potential.

Distribution

Vespa velutina is native to Southeast Asia, occurring in Korea (Monceau et al. 2014), Japan (Kishi and Goka 2017), and China (Robinet et al. 2016). It was introduced to France in 2004, most likely through the importation of bonsai pots (Robinet et al. 2016). This species has very high invasion potential and has easily spread to other European regions (Figure 2) such as Italy (Bertolino et al. 2016), Great Britain (Budge et al. 2017), the Balearic Islands (Leza et al. 2017) and northern Germany (Husemann et al. 2020).



Figure 1. *Vespa velutina* (Lepeletier) feeding on nectar.
Credits: Monceau et al. (2014), Karine Monceau




Figure 2. Distribution of *Vespa velutina* (Lepeletier) in Europe up to the year 2018.
Credits: Daniela Laurino. Used with permission

1. This document is EENY-755, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date May 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ufl.edu/creatures>.
2. Krystal Ashman; Oliver Keller; and Cameron Jack, lecturer; Entomology and Nematology Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

Description

Vespa velutina (Figures 3 and 4) adults are approximately 22 mm in length, roughly the length of a US nickel. According to Monceau et al. (2014), the black and yellow coloration of *Vespa velutina* can be used to differentiate it from similar wasps, such as the European species *Vespa crabro* (Linnaeus). Monceau et al. (2014) also notes that males and females of *Vespa velutina* can be differentiated from each other by their antennae, with female antennae appearing thinner and shorter in length compared to those of male wasps. As with all Hymenoptera, females have a stinger and males do not. Distinguishing between queens and workers proves more difficult, with wing characteristics being the best morphological characters to use (Monceau et al. 2014).



Figure 3. Adult female *Vespa velutina* (Lepeletier), dorsal view.
Credits: Oliver Keller and Krystal Ashman, UF/IFAS



Figure 4. Adult female *Vespa velutina* (Lepeletier), dorsal view.
Credits: Oliver Keller and Krystal Ashman, UF/IFAS



Figure 5. Adult female *Vespa mandarinia* (Smith), dorsal view.
Credits: Oliver Keller and Krystal Ashman, UF/IFAS



Figure 6. Adult female *Vespa mandarinia* (Smith), lateral view.
Credits: Oliver Keller and Krystal Ashman, UF/IFAS



Figure 7. Adult *Sphecus speciosus* (Drury), dorsal view.
Credits: Oliver Keller and Krystal Ashman, UF/IFAS

Life Cycle and Biology

The life cycle of *Vespa velutina* begins with a lone/single queen forming a nest (Figures 8 and 9), laying eggs, and waiting for workers to emerge (Monceau et al. 2014). Nests may be built in a variety of locations, including (but not limited to) bushes, shrubs, treetops, and building rooftops. The nest grows larger during the growing seasons (from spring to autumn), reaching around 6,000 individuals on average (Monceau et al. 2014). Colony growth is usually achieved by the construction of secondary nests (Herrera et al. 2019). Queens become active around March, laying eggs around April, and workers become active in June. Increased predation of honey bee colonies occurs by the wasps in the summer months, and mating of queens and males ensues in the later months of the year (Monceau et al. 2014) (Figure 10). The *Vespa velutina* life cycle is annual, and all workers and males die at the end of the season. New nests are constructed by founder queens at the beginning of each year (Herrera et al. 2019).



Figure 8. Primary nests of *Vespa velutina* (Lepeletier) on shed ceiling represented by numbers 2 and 3. Numbers 1 and 4 represent nests of *Polistes dominula* (Christ).

Credits: Monceau et al. (2014), Jacques Tardits




Figure 9. Secondary nest of *Vespa velutina* (Lepeletier). Credits: Karine Monceau, from Monceau et al. (2014)




Figure 10. Life cycle of *Vespa velutina* (Lepeletier) in France. Credits: Monceau et al. (2014)

Predatory Strategies

Vespa velutina is a predatory wasp that feeds on a variety of arthropods as a source of protein. It is known to be an opportunistic feeder, even feeding on decaying animals (Monceau et al. 2014). This species happens to prefer honey bees, with *Apis mellifera* (Linnaeus) proving to be an easier target compared to other *Apis* species. *Apis cerana* (Fabricius) for example, is more successful in defending against *Vespa velutina*. In response to pheromones produced by *Apis cerana* and its predator, many individuals of *Apis cerana* form a ball around the invader (Figure 11) and use thermoregulation and carbon dioxide to overheat the *Vespa velutina* wasp, resulting in its death (Dong et al. 2018).



Figure 11. *Apis cerana* (Fabricius), forming a heat ball. Credits: Takahashi via GFDL, CC BY-SA-2.1 JP

Economic Importance

Vespa velutina poses a major threat to the beekeeping industry, particularly that of *Apis mellifera*, because it reduces honey bee productivity by preying on individuals (Robinet et al. 2016). So far only few estimates are available, but some reports from Europe mention up to 30% of honey bee hives being weakened by attacks and approximately 5% can be completely destroyed (Monceau et al. 2014). The damage *Vespa velutina* inflicts on the beekeeping industry can be economically important because honey bees produce honey and several other important resources, such as pollen, propolis, and wax. From a human health perspective, it has been noted that wasp stings in general have associated health risks because they can incite allergic reactions (Robinet et al. 2016).

Management

Many management tactics have been proposed to control *Vespa velutina*, including using broad-spectrum insecticides (Poidatz et al. 2018), hunting for nests and trapping adult wasps (Turchi and Derijard 2018), and installing baited traps (Rojas-Nossa et al. 2018). Some have proposed the use of natural enemies such the conopid fly *Conops vesicularis* (Darrouzet et al. 2015) and the mermithid nematode *Pheromermis* sp. (Villemant et al. 2015) as biological control agents. Unfortunately, many of these efforts have not proved successful due to impracticality, high-risk association, or several other factors.

Acknowledgements

The authors would like to acknowledge Dr. Morgan Byron and two other anonymous reviewers for reviewing this article while providing meaningful feedback and constructive suggestions.

Selected References

Bertolino, S., S. Lioy, D. Laurino, A. Manino, and M. Porporato. 2016. "Spread of the invasive yellow-legged hornet *Vespa velutina* (Hymenoptera: Vespidae) in Italy." *Applied Entomology and Zoology* 51: 589–597.

Budge, G., J. Hodgetts, E. P. Jones, J. C. Ostojka-Starzewski, J. Hall, V. Tomkies, N. Semmence, M. Brown, M. Wakefield, and K. Stainton. 2017. "The invasion, provenance and diversity of *Vespa velutina* Lepeletier (Hymenoptera: Vespidae) in Great Britain." *PLoS ONE* 12: e0185172.

Darrouzet, E., J. Gévar, and S. Dupont. 2015. "A scientific note about a parasitoid that can parasitize the yellow-legged hornet, *Vespa velutina nigrithorax*, in Europe." *Apidologie* 46: 130–132.

Dong, S., P. Wen, Q. Zhang, Y. Wang, Y. Cheng, K. Tan, and J. Nieh. 2018. "Olfactory eavesdropping of predator alarm pheromone by sympatric but not allopatric prey." *Animal Behaviour* 141: 115–125.

Herrera, C., A. Marqués, and M. M. Leza. 2019. "Analysis of the secondary nest of the yellow-legged hornet found in the Balearic Islands reveals its high adaptability to Mediterranean isolated ecosystems." In Veitch CR, Clout MN, Martin AR, Russell JC, West CJ. (eds.) *Island invasives: Scaling up to meet the challenge*, Occasional Paper SSC 62. IUCN, Gland, Switzerland. pp. 375–380.

Husemann, M., A. Sterr, S. Maack, and R. Abraham. 2020. "The northernmost record of the Asian hornet *Vespa velutina nigrithorax* (Hymenoptera, Vespidae)." *Evolutionary Systematics* 4: 1–4.

Kishi, K., and K. Goka. 2017. "Review of the invasive yellow-legged hornet, *Vespa velutina* (Hymenoptera: Vespidae), in Japan and its possible chemical control." *Applied Entomology and Zoology* 52: 361–368.

Leza, M., M. A. Miranda, and V. Colomar. 2017. "First detection of *Vespa velutina nigrithorax* (Hymenoptera: Vespidae) in the Balearic Islands (Western Mediterranean): A challenging study case." *Biological Invasions* 20: 1643–1649.

Monceau, K., O. Bonnard, and D. Thiéry. 2014. "*Vespa velutina*: A new invasive predator of honeybees in Europe." *Journal of Pest Science* 87: 1–16.

Poidatz, J., R. L. Plantey, and D. Thiéry. 2018. "Indigenous strains of *Beauveria* and *Metharizium* as potential biological control agents against the invasive hornet *Vespa velutina*." *Journal of Invertebrate Pathology* 153: 180–185.

Robinet, C., C. Suppo, and E. Darrouzet. 2016. "Rapid spread of the invasive yellow-legged hornet in France: The role of human-mediated dispersal and the effects of control measures." *Journal of Applied Ecology* 54: 205–215.

Rojas-Nossa, S. V., N. Novoa, A. Serrano, and M. Calviño-Cancela. 2018. "Performance of baited traps used as control tools for the invasive hornet *Vespa velutina* and their impact on non-target insects." *Apidologie* 49: 872–885.

Turchi, L., and B. Derijard. 2018. "Options for the biological and physical control of *Vespa velutina nigrithorax* (Hym.: Vespidae) in Europe: A review." *Journal of Applied Entomology* 142: 553–562.

Villemant, C., D. Zuccon, Q. Rome, F. Muller, G. O. Poinar Jr., and J. L. Justine. 2015. "Can parasites halt the invader? Mermithid nematodes parasitizing the yellow-legged Asian hornet in France." *PeerJ* 3: e947.