

White Grubs, *Phyllophaga* spp. (Insecta: Coleoptera: Scarabaeidae)¹

H. Lane Selman²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

True white grubs are the larvae of May beetles (also called June Beetles) found in the genus *Phyllophaga*, of which there are over 100 different species. *Phyllophaga* larvae and other larvae of the family Scarabaeidae are often referred to as “white grubs”, including larvae of the **Japanese beetle** (*Popillia japonica* Newman), annual white grubs (*Cyclocephala* spp.), and the green June beetle (*Cotinis nitida* Linnaeus).

Distribution

Phyllophaga spp. and related insects are distributed throughout the United States and Canada. However, the distribution of individual species usually is more restricted.

Description

Adult

May beetles are about 12 to 25 mm (½ to 1 in) long. Most adults are yellow to dark reddish-brown to black, robust, oblong, glossy beetles. Some, such as the green June beetle, are more brightly colored.



Figure 1. Adult Japanese beetle, *Popillia japonica* Newman.
Credits: James Castner, University of Florida



Figure 2. Adult green June beetle, *Cotinis nitida* Linnaeus.
Credits: John L. Capinera, University of Florida

1. This document is EENY-045, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date July 1998. Revised June 2014, April 2017, and December 2023. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.
2. H. Lane Selman; Entomology and Nematology Department, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.



Figure 3. Typical June beetle, adult of “white grub”, of the genus *Phyllophaga*.

Credits: James Castner, University of Florida

Larva

The length of the larvae varies from 20 to 45 mm ($\frac{3}{4}$ –1 $\frac{3}{4}$ in). Larvae are white with a C-shaped body, brown head, and three pairs of legs. The hind portion of the abdomen is slightly enlarged and appears darker due to the soil particles showing through the body wall. Two parallel rows of spines seen on the underside of the last abdominal segment distinguish true white grubs from similar-looking larvae.



Figure 4. Typical white grub of the genus *Phyllophaga*.

Credits: John L. Capinera, University of Florida

Pupa

The length of the pupae varies from 20 to 24 mm ($\frac{3}{4}$ –1 in) long. The pupa is usually white, faint yellow, or dark brown in color.

Egg

Eggs are usually 1.5 to 3 mm ($\sim\frac{1}{16}$ to $\frac{1}{8}$ in) in diameter and found encased in soil aggregates. They are small, spherical, pearly white eggs that darken just before hatching.

Life Cycle

The *Phyllophaga* life cycles vary somewhat because some species complete their growth in one year, while others require up to four years. The common life cycle of the more destructive and abundant of these beetles extends over three years. The adults mate in the evening and, at dawn, females return to the ground to deposit 15 to 20 eggs, 1 to 8 inches deep in the soil. Since the adults are attracted to trees to feed, they tend to lay most eggs in the higher portions of sod near wooded areas.

Eggs hatch about three weeks later into young larvae that feed upon roots and decaying vegetation throughout the summer and, in autumn, migrate downward (to a depth of up to 1.5 meters (5 ft)) and remain inactive until the following spring. The greatest amount of damage occurs as the larvae move near the soil surface to feed on the roots of the plants. The next autumn the larvae again migrate deep into the soil to overwinter, returning near the soil surface the following spring to feed on plant roots until they are fully grown in late spring. These grubs then form oval earthen cells and pupate. The adult beetles emerge from the pupal stage a few weeks later, but they do not leave the ground. The beetles overwinter in the soil, emerging the following year in May or June when feeding, mating, and egg-laying take place.

Management

Although white grubs can be a problem every year, the most serious damage occurs in regular three year cycles. The greatest damage to crops occurs the year after the appearance of the adults. During the years of heavy May beetle infestation, deep-rooted legumes, such as alfalfa or clovers, should be planted. If corn or small grains are present, every effort should be made to keep the field free of grass and weed growth to reduce the number of beetle eggs laid. The year following heavy flights of May beetles, planting corn or potatoes should be avoided in fields that were previously under sod or grass.

Late spring or early autumn plowing destroys many larvae, pupae, and adults in the soil and also exposes the insects to predators, such as birds and skunks. For this cultural practice to be effective, plowing must occur before the grubs migrate below the plow depth. No-tillage or reduced tillage crop management encourages grub populations.

Natural enemies that control these white grubs include parasitic wasps and flies in the genera *Tiphia* and *Myzinum* (Hymenoptera: Tiphidae), and *Pelecinus polyturator* Drury (Hymenoptera: Pelecinidae), and the fly, *Pyrgota*

undata (Diptera: Pyrgotidae). Also, *Cordyceps* fungus infects the grubs.

Inoculating the soil with bacterial spores of *Bacillus popilliae* Dutky and *B. lentimorbus* Dutky aids in reducing populations. These products are available commercially.




Figure 5. Parasitic wasp feeding externally on white grub, *Phyllophaga* spp.



Figure 6. A peleciniid wasp, a natural enemy of white grubs, *Phyllophaga* spp.

Credits: Paul M. Choate, University of Florida



Figure 7. A pyrgotid fly, a natural enemy of white grubs, *Phyllophaga* spp.



Figure 8. White grub, *Phyllophaga* spp., infected with *Cordyceps* fungus.

Credits: Jim Kalisch, University of Nebraska - Lincoln

Selected References

Buss EA. (2006). White grub management in turf.

Davidson RH, Lyon WF. 1979. Insect Pests of Farm, Garden, and Orchard. John Wiley and Sons. New York. pp. 126–128.

Jaques HE. 1951. How to Know the Beetles. WM. C. Brown Company. Dubuque, Iowa. pp. 241.

Luginbill P Sr, Painter HR. 1953. Rev. *Phyllophaga*. U.S. Department of Agriculture Technical Bulletin 1060. 102 pp.

Metcalf RL, Metcalf RA. 1993. Destructive and Useful Insects: Their Habits and Control. McGraw-Hill Book Company, Inc. New York.

North Carolina Corn Information. (1998). White grubs. (no longer available online).

Ritcher PO. 1966. White grubs and their allies: a study of North American scarabaeoid larvae. Oregon State Monograph Studies in Entomology 4: 1–219.

Rolston LH, Barlow T. 1980. Insecticide control of a white grub, *Phyllophaga ephilida* Say, (Coleoptera: Scarabaeidae) on sweet potato. Journal of the Georgia Entomological Society 15: 445–449.

Stehr FW. 1991. Immature Insects (Volume 2). Kendall/Hunt Publishing Co. Dubuque, Iowa. pp. 378–385.

Tashiro H. 1973. Bionomics and control of root feeding insect pests: Grubs and billbugs. Bulletin of the Entomological Society of America 19: 92–94.

Woodruff RE, Beck BM. 1989. The scarab beetles of Florida (USA): (Coleoptera: Scarabaeidae) Part II. The May or June beetles (genus *Phyllophaga*). Arthropods of Florida and Neighboring Land Areas 13: 1–226.