

Bayberry Whitefly, *Parabemisia myricae* (Kuwana) (Insecta: Hemiptera: Aleyrodidae: Aleyrodinae)¹

Avas B. Hamon, Ru Nguyen, and Harold Browning²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

The whitefly, *Bemisia myricae*, was described by Kuwana (1927) from Japan on the hosts *Myrica rubra*, *Morus alba*, and *Citrus*. Takahashi (1952) transferred this whitefly to the genus *Parabemisia* because of long marginal setae on the so-called “pupal” case and blunt lateral tubercles at the base of the lingula.

Bayberry whitefly was first discovered in the United States by California agriculture officials in 1978 (Rose et al. 1981), and in Florida by agriculture officials in early 1984 (Hamon 1986). Early dense populations caused defoliation in California citrus (Rose et al. 1981), but this has not happened in Florida. The early finds in Florida were under natural biological control by hymenopterous parasites, including *Eretmocerus* sp. Apparently the parasites were introduced with the whitefly. According to Mike Rose (personal communication), the *Eretmocerus* species present in Florida is the same as that found in California.

It was not until 1989 that any population in Florida was large enough to cause damage. This occurred in a nursery/greenhouse of the United States Sugar Corporation, Clewiston, Florida, on small citrus seedlings. A chemical control program was being carried out which eliminated the *Eretmocerus* sp., but not the bayberry whitefly. Outdoors, this whitefly is under biological control in Florida, and is only a problem in situations where natural balances are disturbed by the use of chemicals.

Distribution

This whitefly is only known from California and Florida in the United States. Foreign distribution includes China, Hong Kong, Israel, Japan, Taiwan, Malaysia, and Venezuela.

Description

The fourth nymphal skin has 30 to 32 marginal setae including the caudal setae. The anterior spiracular furrows are scarcely visible, but the caudal furrow is slightly ridged longitudinally. The vasiform orifice is elongate triangular with the lingula included. The lingula has two blunt lateral tubercles and two long caudal lingular setae. The operculum covers only the anterior one-third of the vasiform orifice. In vivo, the nymphs are surrounded by a marginal fringe of clear wax.

1. This document is EENY-138, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date June 2000. Revised August 2014 and June 2021. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. Avas B. Hamon; Ru Nguyen, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville; and Harold Browning, professor emeritus & center director, Citrus REC, UF/IFAS Extension, Gainesville, FL 32611.




Figure 1. Fourth nymphal skin "pupal case" of the bayberry whitefly, *Parabemisia myricae* (Kuwana), v=ventral view, d= dorsal view
Credits: Division of Plant Industry




Figure 2. Vasiform orifice [a - lingual, b - operculum] of the nymph of the bayberry whitefly, *Parabemisia myricae* (Kuwana)
Credits: Division of Plant Industry



Figure 3. Nymph of the bayberry whitefly, *Parabemisia myricae* (Kuwana).
Credits: Jeffrey Lotz, Division of Plant Industry

The adult is a small whitish-yellow moth-like insect that flits about when disturbed. The adults have a strong ovipositional preference for very young foliage in the "feather" stage (Walker & Aitken 1985). The adult will frequently place eggs along the leaf margin. At first, the eggs are white, but turn black in a few days.



Figure 4. Adult of the bayberry whitefly, *Parabemisia myricae* (Kuwana).
Credits: Harold Browning, UF/IFAS

Hosts

Camellia sinensis, *Chiococca alba*, *Citrus* spp., *Diospyros kaki*, *Elaeocarpus serratus*, *Ficus carica*, *Gardenia jasminoides*, *Machilus* sp., *Maesa japonica*, *Morus alba*, *Myrica rubra*, *Prunus mume*, *Prunus persica*, *Prunus triflora*, *Psidium guajava*, *Quercus serrata*, *Rhododendron* sp., *Salix*

babylonica, and *Salix gracilistyla*. The most favored hosts are *Citrus* spp. and *gardenia*.

Selected References

Hamon, A.B. 1986. *Parabemisia myricae* (Kuwana). p. 33. In 35th Biennial Report, Div. of Plant Industry. Florida Dept. Agric. and Cons. Serv.

Kuwana, I. 1927. On the genus *Bemisia* (Family Aleyrodidae) found in Japan, with description of new species. *Annotnes Zool. Japan*. 11: 245–253. (Not seen)

Mound, L.A., and S.H. Halsey. 1978. Whitefly of the world. A systematic catalogue of the Aleyrodidae (Homoptera) with host plant and natural enemy data. BM (NH) and John Wiley & Sons. Chichester. 340 p.

Rose, M., P. DeBach, and J. Woolley. 1981. Potential new citrus pest: Japanese bayberry whitefly. *California Agric*. 35: 22–24, illus.

Rose, M., and P. DeBach. 1982. A native parasite of bayberry whitefly. *Citrography*- October: 272–276.

Takahashi, R. 1952. *Aleurotuberculatus* and *Parabemisia* of Japan (Aleyrodidae: Homoptera). *Misc. Rep. Res. Inst. Nat. Resour. Tokyo* 25: 17–24.

Walker, G.P., and D.C.G. Aitken. 1985. Oviposition and survival of bayberry whitefly, *Parabemisia myricae* (Homoptera: Aleyrodidae) on lemons as a function of leaf age. *Environ. Entomol.* 14: 254–257.

Walker, G.P. 1987. Probing and oviposition behavior of the bayberry whitefly (Homoptera: Aleyrodidae) on young and mature lemon leaves. *Ann. Entomol. Soc. America* 80: 524–529.