

Florida Predatory Stink Bug (unofficial common name), *Euthyrhynchus floridanus* (Linnaeus) (Insecta: Hemiptera: Pentatomidae)¹

Frank W. Mead and David B. Richman²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

The predatory stink bug, *Euthyrhynchus floridanus* (Linnaeus) (Figure 1), is considered a beneficial insect because most of its prey consists of plant-damaging bugs, beetles, and caterpillars. It seldom plays a major role in the natural control of insects in Florida, but its prey includes a number of economically important species.



Figure 1. Adult of the Florida predatory stink bug, *Euthyrhynchus floridanus* (L.), feeding on a beetle.

Credits: Lyle J. Buss, University of Florida

Distribution

Euthyrhynchus floridanus is primarily a Neotropical species that ranges within the southeastern quarter of the United States.

Description

Adults

The length of males is approximately 12 mm, with a head width of 2.3 mm and a humeral width of 6.4 mm. The length of females is 12 to 17 mm, with a head width of 2.4 mm and a humeral width of 7.2 mm. *Euthyrhynchus floridanus* (Figure 2) normally can be distinguished from all other stink bugs in the southeastern United States by a reddish spot at each corner of the scutellum outlined against a blue-black to purplish-brown ground color. Variations occur that might cause confusion with somewhat similar stink bugs in several genera, such as *Stiretrus*, *Oplomus*, and *Perillus*, but these other bugs have obtuse humeri, or at least lack the distinct humeral spine that is present in adults of *Euthyrhynchus*. In addition, species of these genera known to occur in Florida have a short spine or tubercle situated on the lower surface of the front femur behind the apex 1/4 to 1/3 of the femoral length. Hayslip et al. (1953) illustrated species in *Stiretrus*, *Euthyrhynchus*, and *Podisus*,

1. This document is EENY157, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date October 2000. Revised April 2016 and July 2022. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. Frank W. Mead, Florida Department of Agriculture and Consumer Services, Division of Plant Industry; and David B. Richman; UF/IFAS Extension, Gainesville, FL 32611.

and a prominently marked *Euthyrhynchus floridanus* was illustrated by Chittenden (1911).



Figure 2. Adult predatory stink bug *Euthyrhynchus floridanus* (L.).
Credits: Lyle J. Buss, University of Florida

The adults of *Euthyrhynchus floridanus* are often confused with the banded form of *Stiretrus anchorago* (F.), from which it differs in size as *Stiretrus anchorago* is less than 10 mm long. Also, the shape of *Stiretrus anchorago* is more globose and resembles a leaf beetle. There are also differences in scutellum size, as *Stiretrus anchorago* has a very large scutellum covering much of the dorsum of the abdomen.

The basic characters of the predatory subfamily Asopinae serve to eliminate the much more common plant-feeding stink bugs:

- Asopines have the first segment of the beak short and thick, free, only its base being between the bucculae, which converge and unite behind or beneath the beak. The base of the beak is close to the end of the tylus.
- Other groups of stink bugs have the first segment of beak slender and embedded between the bucculae, which are wide and parallel. The base of the beak is distinctly separated from the end of the tylus.

Blatchley (1926) and Torre-Bueno (1939) remain the basic keys to stink bugs in the eastern United States.

Eggs

The females lay egg masses with individual eggs somewhat barrel shaped. Eggs and all five nymphal instars were illustrated and described in detail by Oetting and Yonke (1975). Eggs (Figure 3) of *Euthyrhynchus floridanus* are approximately 1 mm in diameter, with short projections around the operculum, and are laid 20 to 90 at a time.



Figure 3. Eggs of *Euthyrhynchus floridanus* (L.), a predatory stink bug.
Credits: Lyle J. Buss, University of Florida

Nymphs

Identification of the nymphs (Figure 4) is less certain, particularly the earlier instars. The available keys are based on the last (fifth) instar, but key characteristics often apply to the fourth instar as well. Hart (1919) included a key to nymphs in the various stink bug subfamilies. DeCoursey and Allen (1968) published a key to the fifth instar nymphs of 25 genera of eastern United States stink bugs. However, four asopine genera (*Alcaeorrhynchus*, *Andrallus*, *Mineus*, and *Oplomus*) reported from Florida were not included. Also listed were field recognition characters of mature nymphs, with *Euthyrhynchus* summarized as having head, thorax, and lateral and medial plates unicolorous metallic green, abdomen bright red; length 8 to 9 mm. The mature nymphs reared by Oetting and Yonke (1975) were 10 to 12.5 mm in length. An occasional mistake of beginners is to confuse *Euthyrhynchus* nymphs with beetles. The latter would have elytra forming a suture dorsally, and the mouthparts would be of the chewing type. Also, the young stink bugs lack wings and have tube-like piercing-sucking mouthparts.



Figure 4. Recently molted nymphs of a predatory stink bug, *Euthyrhynchus floridanus* (L.), with one (center) almost totally emerged from exuviae (cast skin).

Credits: Lynette Schimming

FIRST INSTAR

The first instar is approximately 1.5 mm long. The head width, including eyes, is 0.7 mm, and the humeral width is 1.0 mm. These are difficult to distinguish from the first instar nymphs of *Alcaeorrhynchus grandis* (Dallas), as both have a blue-black head and thorax and red abdomen with dark central and lateral “stripes” composed of dorsal and lateral dark-colored plates. Nymphs of this age do not stray far from the egg mass and may be distinguished by association with the form of the mass and the numbers of eggs. *Alcaeorrhynchus grandis* masses are in multiple rows and contain 100 to 200 eggs.

SECOND INSTAR

The second instar is 2 to 2.5 mm long. The head width is 0.9 mm, and the humeral width 1.1 mm. As in other asopine stink bugs, *Euthyrhynchus floridanus* begins to capture insect prey as a second instar nymph. It looks much like the first instar and retains the same coloration.

THIRD INSTAR

The third instar is approximately 4 mm long. The head width is 1.2 mm, and the humeral width is 1.7 mm. This instar is colored much as in the first instar. The iridescence of the blue-black markings is more noticeable because of the larger size of the nymph.

FOURTH INSTAR

The fourth instar (Figure 5) is 6 to 7 mm long. The head width is 1.7 mm and the humeral width is 2.9 mm. It is much like the earlier stages in appearance.



Figure 5. Fourth instar nymph of the predatory stink bug, *Euthyrhynchus floridanus* (L.).

Credits: Lyle J. Buss, University of Florida

FIFTH INSTAR

The fifth instar (Figure 6) is 8 to 9 mm long. The head width is 2.1 mm, and the humeral width is 4.8 mm. The wing pads, which are blue-black, are prominent, but otherwise the nymph is much like the preceding stages.



Figure 6. A fifth instar nymph of the Florida predatory stink bug, *Euthyrhynchus floridanus* (L.). Notice the prominent wing pads that are characteristic of this stage.

Credits: Lyle J. Buss, University of Florida



Figure 7. Nymphs of the Florida predatory stink bug, *Euthyrhynchus floridanus* (L.) feeding on an earwig.

Credits: Anthony Ditro, Elizabeth City, NC

Life Cycle and Biology

Euthyrhynchus floridanus has been reared in the laboratory by Ables (1975), Oetting and Yonke (1975), and Richman and Whitcomb (1978). At 26 to 27°C, and with a photoperiod of 14:10, both Ables (1975) and Richman and Whitcomb (1978) found that the length of time from egg to adult was 58 days. The egg stage lasted 18 to 19 days.

Ables (1975) reared South Carolina examples of *Euthyrhynchus floridanus* in the laboratory, providing prey of greater wax moth larvae, *Galleria mellonella* (L.); all stages of the Mexican bean beetle, *Epilachna varivestis* Mulsant; and larvae of tobacco budworm, *Heliothis virescens* (F.). The life cycle was about 89 days under the conditions provided. Oetting and Yonke (1975) reared Missouri stock of *Euthyrhynchus floridanus* in the laboratory, providing black cutworms, *Agrotis ipsilon* (Hufnagel), as a food source. Oetting and Yonke (using a temperature of 24°C and a photoperiod of 13:11) found the length of time from egg to adult was 100 days, with the egg stage lasting 33 days.

These papers (Ables 1975, Oetting and Yonke 1975) included comments on the gregarious nature of the nymphs and even the adults tended to aggregate at night in the cages. Ables (1975) commented that capture of prey and feeding by early instars often appeared to be a group effort. He added that such behavior would be of a selective advantage because it would allow feeding on large prey unavailable to a single nymph.

Records in the Florida State Collection of Arthropods show that *Euthyrhynchus floridanus* has been collected during all months of the year in Florida and in a wide variety of habitats, as its prey is highly variable. There is a peak in the spring and again in the fall. Of 140 habitat records, 25% have been ornamentals, 17% citrus, 8% in traps, 6% in

weeds and turf, and the remainder in miscellaneous fruits, trees, and random habitats.

Host

Prey records include:

- *Anticarsia gemmatilis* Hübner, velvetbean caterpillar
- *Bassareus brunnipes* (Olivier), a leaf beetle
- *Cosmophilla erosa* Hübner, “okra” or “mallow” caterpillar
- *Datana integerrima* Grote & Robinson, walnut caterpillar
- *Diaprepes abbreviatus* (L.), diaprepes root weevil
- *Hypera postica* (Gyllenhal), alfalfa weevil
- *Leptinotarsa decemlineata* (Say), Colorado potato beetle
- *Lymire edwardsii* (Grote), Edwards wasp moth
- *Nezara viridula* (L.), southern green stink bug
- *Ormenaria* (= *Monoflata*) *rufifascia* (Walker), a flatid planthopper
- *Papilio cressphontes* Cramer, orangedog

Selected References

- Ables JR. 1975. Notes on the biology of the predacious pentatomid *Euthyrhynchus floridanus* (L.). Journal of the Georgia Entomological Society 10: 353-356.
- Blatchley WS. 1926. Heteroptera or True Bugs of North America with Special Reference to the Faunas of Indiana and Florida. Nature Publishing Company, Indianapolis, USA. 1116 pp.
- Chittenden FH. 1911. Notes on various truck-crop insects. U.S. Department of Agriculture Bureau of Entomology Bulletin 82, Part VII: 85-93.
- DeCoursey RM, Allen RC. 1968. A generic key to the nymphs of the Pentatomidae of the eastern United States (Hemiptera: Heteroptera). University of Connecticut Occasional Papers 1: 141-151.
- Distant WL. 1880-1893. Insecta. Rhynchota. Hemiptera: Heteroptera. Vol. 1. In Biologia Centrali-Americana. London. 462 pp.
- Hart CA. 1919. The Pentatomoidea of Illinois with keys to the Nearctic genera. Bulletin of the Illinois Natural History Survey 13: 157-223.

Hayslip NC, Genung WG, Kelsheimer EG, Wilson JW. 1953. Insects attacking cabbage and other crucifers in Florida. Florida Agricultural Experiment Station Bulletin 534: 1-57.

Hoffman RL. 1971. The insects of Virginia: No. 4. Shield bugs (Hemiptera: Scutelleroidea: Scutelleridae, Coreidae, Cydnidae, Pentatomidae). Virginia Polytechnic Institute and State University, Research Division Bulletin 67: 1-61.

McDonald FJD. 1966. The genitalia of North American Pentatomoidea (Hemiptera: Heteroptera). Quaestiones Entomologicae 2: 7-150.

Oetting RD, Yonke TR. 1975. Immature stages and notes on the biology of *Euthyrhynchus floridanus* (L.) (Hemiptera: Pentatomidae). Annals of the Entomological Society of America 68: 659-662.

Richman DB, Whitcomb WH. 1978. Comparative life cycles of four species of predatory stink bugs (Hemiptera: Pentatomidae). Florida Entomologist 61: 113-119.

Torre-Bueno JR. de la. 1939. A synopsis of the Hemiptera: Heteroptera of America north of Mexico. Part 1. Families Scutelleridae, Cydnidae, Pentatomidae, Aradidae, Dyso-diidae and Temitaphididae. Entomologia Americana 19: 141-304.