

Bee Killers, *Mallophora bomboides* (Wiedemann), *Mallophora orcina* (Wiedemann), and *Mallophora nigra* Williston (Insecta: Diptera: Asilidae)¹

Gary J. Steck²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

Three large predaceous **robber flies** of the genus *Mallophora* occur in Florida (Bromley 1950).

- *Mallophora bomboides* (Wiedemann) is known as the Florida bee killer
- *Mallophora orcina* (Wiedemann) is known as the southern bee killer
- *Mallophora nigra* Williston is known as the black bee killer

Two other species are recorded from Florida: *Mallophora rex* Bromley and *Mallophora chrysomela* Bromley. Morphologically, there is nothing to distinguish them from *Mallophora bomboides* (Wiedemann), and they are considered to be color variants of the latter. Also, *Mallophora nigra* Williston may be merely a wholly black color variant of *Mallophora bomboides* (Cole and Pritchard 1964).

Distribution and Phenology

At least 20 species in seven genera of asilid flies in the southeastern United States prey on various Hymenoptera, including honeybees (*Apis mellifera* L.). Of these, *Mallophora orcina* is the most prominent, with 80% or more of its diet composed of honeybees (Bromley 1946).

Mallophora bomboides occurs in the southeastern United States including North Carolina, South Carolina, Georgia, Florida, Alabama, and Mississippi. Flight time in Florida is from April through December, though they are most common in late summer.

Mallophora orcina occurs in the eastern US from Florida north to Ohio and west to Missouri and Arkansas. Flight time in Florida is from April through August, though they are most common in mid-summer.

Mallophora nigra is uncommon. It is apparently even rarer outside of Florida, having been recorded only from “North Carolina” (Cole and Pritchard 1960), presumably North Carolina. Flight time in Florida is from July through September.

1. This document is EENY315 (originally published as DPI Entomology Circular No. 366), one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date March 2004. Revised September 2009, August 2012, March 2013, October 2016, and January 2023. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entnemdept.ufl.edu/creatures/>.

2. Gary J. Steck, Florida Department of Agriculture and Consumer Services, Division of Plant Industry; UF/IFAS Extension Gainesville, FL 32611.



Figure 1. The Florida bee killer, *Mallophora bomboides* (Wiedemann), with honey bee prey.
Credits: Nancy West, University of Florida



Figure 2. The southern bee killer, *Mallophora orcina* Williston.
Credits: Gary Steck, FDOACS-DPI

Description

Members of this genus in Florida are large, robust flies. The wings are smoky brown and the body has dense patches of black and yellow or white hairs. Bee killers resemble [bumblebees](#) and [carpenter bees](#) in shape and coloration. They generally mimic all of the five bumblebee species occurring in Florida (Stange 1992). They make a beelike hum or buzz when flying (Linsley 1960).

Biology

Adults typically occur in open habitats, often in the vicinity of apiaries. They perch on stalks of weeds or on tips of shrubs from which they launch their attack (Brower et al. 1960). Prey are primarily social bees and wasps, including honey bees, bumble bees, carpenter bees, *Polistes* and *Vespa* wasps. Honey bees may be particularly suitable as prey because of their slow flight, local abundance, and appropriate body size (Poulton 1906). Eggs are laid into the soil. Specific feeding habits of larvae of the species discussed here are unknown, but larvae of other *Mallophora* species are ectoparasites on scarabaeid beetle grubs in the soil (Knutson 1972).

Key to the of Florida

1. Abdomen and scutellum wholly black-haired..... *Mallophora nigra*




Figure 3. Lateral view of the black bee killer, *Mallophora nigra* (Wiedemann).
Credits: Jeff Hollenbeck

- 1'. Abdomen and scutellum with white or yellow hairs..... 2
2. Basal four or five abdominal tergites densely covered with yellow hairs; and remaining segments wholly with black hairs; ventrum of abdomen wholly black haired. Average body length about 20 mm..... *Mallophora orcina*
- 2'. Basal three abdominal tergites densely covered with yellow hairs, 4th and 5th tergites with black hairs, and final 2 segments with pale hairs; ventrum of abdomen with yellow hairs. Average body length about 25 mm..... *Mallophora bomboides*



Figure 4. The southern bee killer, *Mallophora orcina* Williston.
Credits: Gary Steck, FDACS-DPI



Figure 5. Frontal view of the Florida bee killer, *Mallophora bomboides* (Wiedemann).
Credits: Jeff Hollenbeck



Figure 6. Dorsal view of the Florida bee killer, *Mallophora bomboides* (Wiedemann).
Credits: Jeff Hollenbeck



Figure 7. Ventral view of the Florida bee killer, *Mallophora bomboides* (Wiedemann).
Credits: Jeff Hollenbeck

Economic Importance

Instances of economic losses to beekeepers due to the depredations of asilid bee killers are sporadic, but Florida is one of the few states where such losses have been reported (Bromley 1950). During July 2008, hundreds of these insects were seen attacking beehives in a watermelon field in Little Lake City, Gilchrist County, Florida (Halbert 2008).

Selected References

Bromley SW. 1946. Bee-killing Asilidae of the southeastern states (Diptera). Proceedings of the Entomological Society of Washington 48: 16–17.

Bromley SW. 1950. Florida Asilidae (Diptera) with description of one new species. Annals of the Entomological Society of America 43: 227–239.

Brower LP, Brower JVZ, Westcott PW. 1960. Experimental studies in mimicry. 5. The reactions of toads (*Bufo terrestris*) to bumblebees (*Bombus americanorum*) and their robberfly mimics (*Mallophora bomboides*), with a discussion of aggressive mimicry. The American Naturalist 94: 343–355.

Cole FR, Pritchard AE. 1964. The genus *Mallophora* and related asilid genera in North America (Diptera: Asilidae). University of California Publications in Entomology 36: 43–100.

Halbert SE. (July–August 2008). Tri-Ology Report: Entomology section: Arthropod detection. FDACS-Division of Plant Industry. (no longer available online)

Knutson LV. 1972. Pupa of *Neomochtherus angustipennis* (Hine), with notes on feeding habits of robber flies and a review of publications on morphology of immature stages (Diptera: Asilidae). Proceedings of the Biological Society of Washington 85: 163–178.

Linsley EG. 1960. Ethology of some bee- and wasp-killing robber flies of southeastern Arizona and western New Mexico (Diptera: Asilidae). University of California Publications in Entomology 16: 357–381.

Poulton EB. 1906. Predacious insects and their prey. Transaction of the Entomological Society of London 1906: 323–409.

Stange LA. 1992. The bumble bees of Florida (Hymenoptera: Apidae). Florida Department of Agriculture & Consumer Services, Division of Plant Industry, Gainesville. Entomology Circular 353. 2 pp.