

Predatory Mite, *Neoseiulus californicus* (McGregor) (Arachnida: Acari: Phytoseiidae)¹

Elena M. Rhodes and Oscar E. Liburd²

The Featured Creatures collection provides in-depth profiles of insects, nematodes, arachnids and other organisms relevant to Florida. These profiles are intended for the use of interested laypersons with some knowledge of biology as well as academic audiences.

Introduction

The predatory mite *Neoseiulus californicus* (McGregor) has characteristics of both type II specialist predatory mites and type III generalist predatory mites. *Neoseiulus californicus* prefers Tetranychid mites as food, but will also consume other mite species, small insects, such as thrips, and even pollen when the primary prey is unavailable.

Neoseiulus californicus is often used to control the **twospotted spider mite**, *Tetranychus urticae* Koch, and other phytophagous mites on various crops in temperate and subtropical regions around the world.

Synonymy

Neoseiulus californicus has a very complex taxonomic history. It was first described by McGregor in 1954 from lemon in California as *Typhlodromus californicus*. After 1954, it was moved to the genus *Amblyseius* and later to the genus *Neoseiulus* or *Cydnodromus*, and now *Neoseiulus chilensis* (Dosse) is considered a synonym of *Neoseiulus californicus*.



Figure 1. Adult *Neoseiulus californicus* (McGregor) feeding on a twospotted spider mite, *Tetranychus urticae* Koch.

Credits: Lyle Buss, UF/IFAS

Distribution

Natural populations of *Neoseiulus californicus*, are found in Argentina, California, Chile, Florida, Japan, South Africa, Texas, parts of southern Europe, and all along the border of the Mediterranean Sea. *Neoseiulus californicus* has been found on many crops including avocado, citrus, and other fruit trees. They are also found on cassava, corn, grapes, strawberries, and several vegetable crops and ornamental plants. They prefer warm 10°C–33°C (50°F–91°F) temperatures, but they can tolerate much colder temperatures for short periods of time. For example, they can survive the winters in north Florida where temperatures can fall below

1. This document is EENY-359, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date November 2005. Revised May 2009, August 2015, and December 2023. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>.
2. Elena M. Rhodes; and Oscar E. Liburd, professor, Entomology and Nematology Department; UF/IFAS Extension, Gainesville, FL 32611.

freezing at night. They can tolerate a wide range of humidity (40%–80% relative humidity) but prefer humidity at the upper end of this range.

Description

Neoseiulus californicus eggs are football shaped, approximately 0.04 mm (0.00016 in) in length, and are pale whitish in color. Larvae have only six legs and are translucent in color. Both nymphal stages, the protonymph and the deutonymph, resemble the adults except that they are smaller and cannot reproduce. Adult females are approximately 0.1 mm (0.00039 in) in length and oval in shape. Males are slightly smaller than females. Both males and females are translucent and can be pale orange, peach, or pink in color.



Figure 2. Two eggs of *Neoseiulus californicus* (McGregor) attached to leaf hairs on a strawberry leaf.

Credits: Elena M. Rhodes, UF/IFAS

Life Cycle

Neoseiulus californicus females can lay up to four eggs a day. However, two eggs per day is the average. Eggs take from 1.5 to 4.0 days to hatch depending on the temperature. Eggs hatch into six-legged larvae, which can progress to the protonymphal stage without feeding. The larval stage can last from 0.5 to 1.0 day. *Neoseiulus californicus* then passes through two nymphal stages: protonymph and deutonymph. Both stages (protonymph and deutonymph) are active feeders. Each nymphal stage can last from 1.0 to 3.0 days. Total developmental time can be as short as 4.0 days or as long as 12.0 days depending on the temperature. *Neoseiulus californicus* develops more quickly at higher temperatures. Adults live for approximately 20 days.

Neoseiulus californicus develops faster when consuming the twospotted spider mite, *Tetranychus urticae* Kotch, than when consuming other prey sources. However, it will also

successfully develop and reproduce when consuming other mite species including *Aculus schlenchtendali* (Nalepa), *Oligonychus pratensis* (Banks), *Oligonychus perseae* Tuttle, *Oligonychus ilicis* (McGregor), *Panonychus ulmi* (Koch), *Phytonemus pallidus* (Banks), *Polyphagotarsonemus (Stenotarsonemus) latus* Banks (the **broad mite**), and *Phytonemus pallidus* L. (the **cyclamen mite**). Many of these mites are crop pests. It can also survive and reproduce by consuming thrips and other small insects, but reproduction is very low. *Neoseiulus californicus* can even survive for a short period of time by consuming only pollen.



Figure 3. Dorsal view of a *Neoseiulus californicus* (McGregor) larva indicated by pointer. An adult *Neoseiulus californicus* and a cluster of spider mite eggs are shown for size comparison.

Credits: Jack Kelly Clark, University of California



Figure 4. Dorsal view of adult female *Neoseiulus californicus* (McGregor) on a strawberry leaf.

Credits: Elena M. Rhodes, UF/IFAS



Figure 5. A strawberry leaf infested with twospotted spider mite, *Tetranychus urticae* Koch, adults and their eggs.

Credits: Lyle Buss, UF/IFAS

Economic Importance

Neoseiulus californicus is used commercially around the world to control the twospotted spider mite and several other economically important mites on avocado, citrus, dwarf hops, grapes, raspberries, roses and other ornamentals, strawberries, and several vegetable crops. It has been shown to effectively control the twospotted spider mite and *Phytonemus pallidus* on strawberries in glasshouses in the UK and in greenhouses in Argentina. In the United States, *Neoseiulus californicus* has been used successfully to control twospotted spider mites on field grown strawberries in southern California and Florida. *Neoseiulus californicus* has also been used to control the twospotted spider mite on dwarf hops and to control *Oligonychus perseae* on avocado, to mention a few examples.

The recommended release rate depends on pest species, pest density, and crop. In strawberry, a release rate of one female *Neoseiulus californicus* per 10 twospotted spider mites is recommended once mites are present. A predator in first technique, where *Neoseiulus californicus* is released at a low rate before twospotted spider mite infestations appear, has become popular in California strawberries. If twospotted spider mite populations are already high, it may be necessary to spray a miticide before releasing *Neoseiulus californicus*. The predatory mite can tolerate applications of some miticides.



Figure 6. Commercial presentation of *Neoseiulus californicus* (McGregor) to be released on the field (courtesy of Koppert Biological Supply Co.).

Credits: Elena Rhodes, UF/IFAS

Selected References

Barber A, Campbell CAM, Crane H, Lilley R, Tregidga E. 2003. "Biocontrol of two-spotted spider mite *Tetranychus urticae* on dwarf hops by the phytoseiid mites *Phytoseiulus persimilis* and *Neoseiulus californicus*." *Biocontrol Science and Technology* 13: 275–284.

Castagnoli M, Liguori M. 1991. Laboratory observations on duration of copulation and egg production of three Phytoseiid species fed on pollen. *In: The Acari*. New York, NY: Chapman and Hall.

Castagnoli M, Simoni S. 2003. "*Neoseiulus californicus* (McGregor) (Acari: Phytoseiidae): Survey of biological and behavioral traits of a versatile predator." *Redia* 86: 153–164.

Croft BA, Monetti LN, Pratt PD. 1998. "Comparative life histories and predation types: Are *Neoseiulus californicus* and *N. fallacies* (Acari: Phytoseiidae) similar type II selective predators of spider mites." *Environmental Entomology* 27: 531–538.

Easterbrook MA, Fitzgerald JD, Solomon MG. 2001. "Biological control of strawberry tarsonemid mite *Phytonemus pallidus* and two-spotted spider mite *Tetranychus urticae* on strawberry in the UK using species of *Neoseiulus* (*Amblyseius*) (Acari: Phytoseiidae)." *Experimental & Applied Acarology* 25: 25–36.

- Fraulo AB, McSorley R, Liburd OE. 2008. "Effects of the biological control agent *Neoseiulus californicus* (Acari: Phytoseiidae) on arthropod community structures in north Florida strawberry fields." *Florida Entomologist* 91: 336–345.
- Gerson U, Smiley RL, Ochoa R. 2003. *Mites (Acari) for Pest Control*. Ames, IA: Blackwell Publishing.
- Greco NM, Liljestrom GG, Sanchez NE. 1999. "Spatial distribution and coincidence of *Neoseiulus californicus* and *Tetranychus urticae* (Acari: Phytoseiidae, Tetranychidae) on strawberry." *Experimental and Applied Acarology* 23: 567–580.
- Hoddle MS, Aponte O, Kerguelen V, Heraty J. 1999. "Biological control of *Oligonychus perseae* (Acari: Tetranychidae) on avocado: Evaluating release timing, recovery, and efficacy of six commercially available phytoseiids." *International Journal of Acarology* 25: 211–219.
- Johnson WT, Lyon HH. 1991. *Insects that feed on trees and shrubs*. 2nd edition. Comstock Publishing Associates. 560p.
- Krantz GW. 1978. *A manual of acarology*. Corvallis. Oregon State University, OR. 509p.
- Liburd OE, Seferina GG, Dinkins DA. 2003. Suppression of twospotted spider mites. In: UF/IFAS Berry/Vegetable Times. November 2003.
- Ma W-L, Laing JE. "Biology, potential for increase and prey consumption of *Amblyseius chilenesis* (Dosse) (Acarina: Phytoseiidae)." *Entomophaga* 18: 47–60.
- McMurtry JA, Croft BA. 1997. "Life-styles of Phytoseiid mites and their roles in biological control." *Annual Review of Entomology* 42: 291–321.
- Oatman, ER, McMurtry JA, Gilstrap FE, Voth V. 1977. "Effect of releases of *Amblyseius californicus* on the twospotted spider mite on strawberry in Southern California." *Journal of Economic Entomology* 70: 638–640.
- Rondon SI, Price JF, Liburd OE, Francis R, Cantliff DJ. (2004). *Neoseiulus californicus* McGregor: A Predatory Mite Species for Controlling Twospotted Spider Mites in Strawberries. HS1001. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <https://ufdc.ufl.edu/IR00002733/00001> (July 2019).