

Chikungunya¹

C. Roxanne Connelly, Christopher N. Mores, Chelsea T. Smartt, and Walter J. Tabachnick²

What is Chikungunya?

Chikungunya is a virus that can be transmitted to humans by mosquitoes. The resulting illness is also called chikungunya. It was originally described in the early 1950s after an outbreak in a Swahili village on the Makonde plateau that lies between Tanzania and Mozambique. The translation of chikungunya from Makonde means “illness of the bended walker,” “that which bends up,” stooped over, walking bent over, or “bended walker.” In India, it is known as Aakyda, meaning “stiff man” and Maakyda meaning “monkey-like”. These words refer to the arthritic condition that occurs in some patients which gives rise to a stooped posture. The virus was first found in Asia, isolated in Bangkok, Thailand in 1958.

The Virus

Chikungunya virus, or CHIKV, is a member of the Alphavirus genus in the family Togaviridae. CHIKV is related to other alphaviruses like Ross River, Sindbis, and Venezuelan equine encephalitis viruses.

What are the symptoms?

Reported symptoms include fever, chills, headache, rash, and severe joint pain with or without swelling. Pain in the joints is a major feature of chikungunya symptoms. The ankles and wrists are most commonly affected. Intense pain due to pressure placed on a wrist is commonly used to help diagnose the disease.

Symptoms begin to appear within 2–12 days after being infected from the bite of an infected female mosquito. Initial symptoms include a sudden onset of flu-like symptoms including severe headache, chills, insomnia, fever, joint pain, nausea, and vomiting. A rash may occur, first as a flush over the face and chest, followed by a rash that can have lesions. There can be mild hemorrhaging in children. Joint pain can persist for many months or even years after the other symptoms have subsided. All but a few patients recover.

How is it transmitted to humans?

The cycle of transmission is from mosquito to human and back to mosquito. Transmission to humans may occur when infected female mosquitoes attempt to feed on a human host. The primary species of mosquitoes that may transmit the virus are *Aedes aegypti* (Figure 1) and *Aedes albopictus* (Figure 2). The eggs of these mosquitoes are laid just above the waterline in water-holding containers. When the water level rises, the eggs hatch into larvae, where they continue to develop through the pupal stage, then to the adult that will fly to seek a blood meal. During one outbreak, a nurse caring for an infected patient in France came down with chikungunya fever, suggesting that transmission may also occur from person to person, without the involvement of a mosquito vector.

1. This document is ENY-736, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date April 2007. Revised July 2010, September 2013, and January 2017. Reviewed July 2020. Visit the EDIS website at <https://edis.ifas.ufl.edu>.
2. C. Roxanne Connelly, former professor; Christopher N. Mores, former assistant professor; Chelsea T. Smartt, associate professor; and Walter J. Tabachnick, former professor; Department of Entomology and Nematology, Florida Medical Entomology Laboratory, UF/IFAS Extension, Vero Beach, FL.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 1. *Aedes aegypti* female.
Credits: James M. Newman, UF/IFAS/FMEL



Figure 2. *Aedes albopictus* female.
Credits: Sean McCann

How is it diagnosed?

Chikungunya is suspected when there is epidemic disease with the characteristic triad of symptoms of fever, rash, and joint pain. Chikungunya can easily be confused with another mosquito-borne human pathogen called dengue. The definitive diagnosis of chikungunya requires laboratory testing. The virus can be detected during the first 48 hours of disease, and may be detected as late as day 4 in some patients. Usually chikungunya is diagnosed by detecting patient antibodies produced by the immune response to the virus and found in the blood. Antibodies to CHIKV persist in the blood in excess of 6 months.

What is the cure?

As with the other mosquito-borne viruses, there is no vaccine, nor is there a specific “cure.” The fever and aches can be treated with traditional medicines like analgesics and anticonvulsants. Aspirin should be avoided. There is no medication that will kill the virus. Supportive care with rest is advised to deal with acute joint pain. Movement and mild exercise tend to improve stiffness. However, heavy exercise may exacerbate symptoms.

Chikungunya is usually self-limiting and recovery was thought to be universal, however, in January 2006, a high number of deaths (>200), either directly or indirectly related to infection with the virus, were reported from Reunion Island in the Indian Ocean. Excessive deaths were seen in patients who were 75 years of age or older. Once infected, immunity to further infection is long-lasting.

Where does it occur?

Chikungunya outbreaks have been reported from India, Africa, and Southeast Asia. Between March 1, 2005 and April 30, 2006, there were 255,000 cases reported from Reunion Island in the Indian Ocean, with an estimated fatality rate of 1 per 1,000 cases. The epidemic spread throughout the Indian Ocean, reaching India which had thousands of cases. In 2009, there were 4,430 cases in Malaysia, almost 50,000 cases reported from Thailand. During 2006 through 2011, 117 cases were identified among US travelers (CDC 2012).

Infections may occur in areas that are not considered endemic, or where the virus does not normally occur. Travel and globalization increase the possibility of epidemic outbreaks in other regions around the globe, wherever the mosquitoes capable of transmitting CHIKV are found.

Are Floridians at risk?

Transportation and globalization are important factors in assessing the risk of exposure to CHIKV for Floridians. The mosquito species that have been shown to be good vectors of CHIKV are present and well-established in Florida. It is believed that the CHIKV circulates at high levels in the blood of the human host, and therefore can directly infect capable mosquitoes. Travelers to endemic regions who become infected and then return to the United States, or visitors to the United States who are residents of those regions, may serve as a source of spreading the virus to local mosquitoes. This would initiate a transmission cycle in Florida capable of producing more cases and a potential chikungunya epidemic.

In 2014, there were 11 cases of local transmission of Chikungunya reported in Florida from Miami-Dade (2), Palm Beach (4), St. Lucie (4) and Broward (1) counties. By the end of 2016, there were no further reports of local transmission.

How can it be prevented?

Mosquito control and public health agencies in Florida will need knowledge of *Ae. aegypti* and/or *Ae. albopictus* populations, where they are produced, and hot spots with large populations that could be easily targeted for control.

What can or should a mosquito control district do to control these populations? There is a great deal of information and experience clearly demonstrating that effective *Ae. aegypti* and *Ae. albopictus* control must involve **public support** and **public participation** to effectively reduce larval habitats. Plans must be in place on how to utilize citizens for these important surveillance and control efforts.

Prevent mosquito infection. It is clear that anyone with a chikungunya infection should limit all exposure to mosquito bites. This can be done with use of repellents, or more appropriately staying indoors or under mosquito netting. Doing so during the first 4–6 days after infection when there is virus in the blood will avoid spreading the infection.

Reduce the mosquito vectors. *Ae. albopictus* and *Ae. aegypti* lay their eggs in containers to complete larval development. This includes containers from bucket-sized to something as small as a bottle cap that holds water. *Ae. albopictus* in particular is able to utilize an array of both natural and artificial containers. To reduce the mosquito vectors, source reduction is the best method.

- Remove all container sources that can hold water including bottles, cans, flower pots with saucer bases, clogged gutters, tires, lawn equipment such as wheel barrows, and items that are covered with tarps.
- Other types of containers can be modified as follows: flush and replace water in bird baths and pet dishes several times a week; cover water collection containers, such as rain barrels, with netting to prevent female mosquitoes from entering to lay eggs.
- Start a neighborhood watch program and sponsor container clean-up days to remove or modify water-holding containers in the area.

Be an informed traveler. Consult the Centers for Disease Control website for information on the current status of mosquito-borne disease transmission, as well as other diseases across the globe prior to travel: <http://www.cdc.gov/travel/>.

References

CDC. 2012. “Chikungunya distribution and global map.” <http://www.cdc.gov/chikungunya/map/index.html>

CDC. 2014. “Notes from the Field: Transmission of Chikungunya Virus in the Continental United States - Florida, 2014.” <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6348a4.htm>

Grandadam, M., V. Caro, S. Plumet, J-M Thiberge, Y. Souares, A-B Failloux, et al. 2011. “Chikungunya virus, southeastern France.” *Emerging Infectious Diseases*. <http://www.cdc.gov/EID/content/17/5/910.htm>

Josseran, L., C. Paquet, A. Zehgnoun, N. Caillere, A. Le Terte, J. Solet, and M. Ledrans. 2006. “Chikungunya disease outbreak, Reunion Island.” *Emerging Infectious Diseases*. 12:1994–1995.

Parola, P., X. deLamballerie, J. Jourdan, C. Rovey, V. Vaillant, P. Minodier, P. Broqui, A. Flahault, D. Raoult, and R. Charrel. 2006. “Novel Chikungunya virus variant in travelers returning from Indian Ocean islands.” *Emerging Infectious Diseases*. 12:1493–1499.